ERRATUM

Erratum to: Preclinical Comparison of the Amyloid- β Radioligands [^{11}C]Pittsburgh compound B and [^{18}F]florbetaben in Aged APPPS1-21 and BRI1-42 Mouse Models of Cerebral Amyloidosis

Ann-Marie Waldron,¹ Jeroen Verhaeghe,¹ Leonie wyffels,^{1,2} Mark Schmidt,³
Xavier Langlois,³ Annemie Van Der Linden,⁴ Sigrid Stroobants,² Steven Staelens¹

¹Molecular Imaging Center Antwerp, University of Antwerp, Campus Drie Eiken – UC, Universiteitsplein 1, 2610, Wilrijk, Antwerp, Belgium

²Nuclear Medicine, University Hospital Antwerp, Antwerp, Belgium

³Department of Neuroscience, Janssen Pharmaceutica NV, Beerse, Belgium

⁴Bio-Imaging Laboratory, University of Antwerp, Antwerp, Belgium

Mol Imaging Biol (2015) 17:688–696

DOI: 10.1007/s11307-015-0833-9

Figure 1c was in error. The correct Figure 1 is shown below.

Figure 1: Ex vivo quantification of total amyloid- β in APPPS1-21 and BRI 1–42 mice. Amyloid- β load was quantified as the percent marker stain (JRF/A β N/25 positive) per region and is depicted in a max and min boxplot in graph for **a** APPPS1-21 mice and **d** for BRI1-42 mice. Representative immunohistochemical images show the dorsal hippocampus and neighboring cortical region (scale bar=200 μm) and the cerebellum (scale bar=20 μm) from APPPS1-21 mice (**b**, **c**) and BRI1-42 mice (**e**, **f**).

The online version of the original article can be found under doi:10.1007/s11307-015-0833-9.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s11307-015-0833-9>.

Correspondence to: Steven Staelens; e-mail: steven.staelens@uantwerpen.be