Reforestation of abandoned pasture on Hokkaido, northern Japan: effect of plantations on the recovery of conifer-broadleaved mixed forest

Kikuko Shoyama

Published online: 16 April 2008

© International Consortium of Landscape and Ecological Engineering and Springer 2008

Erratum to: Landscape Ecol Eng DOI 10.1007/s11355-008-0034-7

The original version of this article unfortunately contained a mistake. The legend to Fig. 4 was incorrect. The corrected legend is given below.

Fig. 4 DCA ordination diagram showing the plot scores. The first and second axes accounted for 34 and 28% of the

total variance, respectively. Vegetation type a, *Sasa* community, *open triangle*; type b, meadow community, *open square*; type A, conifer-broadleaved forest, *filled circle*; type B, broadleaved forest, *filled triangle*; type C, birch plantation, *filled square*; type D, larch plantation, *filled diamond*; Control, *cross + circle*; gray years 1986–1988; black years 2005–2006

The online version of the original article can be found under doi:[10.1007/s11355-008-0034-7](https://doi.org/10.1007/s11355-008-0034-7).

K. Shoyama (✉)
Graduate School of Environmental Science,
Hokkaido University, Sapporo 060-0810, Japan
e-mail: shoyama@ees.hokudai.ac.jp