Correction to: Influence of pyrolysis temperature and production unit on formation of selected PAHs, oxy-PAHs, N-PACs, PCDDs, and PCDFs in biochar—a screening study

Eva Weidemann¹ · Wolfram Buss² · Mar Edo¹ · Ondřej Mašek² · Stina Jansson¹

Published online: 5 January 2018

© Springer-Verlag GmbH Germany, part of Springer Nature 2017

Correction to: Environ Sci Pollut Res

<https://doi.org/10.1007/s11356-017-0612-z>

The correct presentation of Figs. 1, 3 and Table 3 are shown in this paper.

Fig. 1 xxxx

The online version of the original article can be found at
<https://doi.org/10.1007/s11356-017-0612-z>

✉ Stina Jansson
stina.jansson@umu.se

¹ Department of Chemistry, Umeå University, SE-901
87 Umeå, Sweden

² UK Biochar Research Centre, School of GeoSciences, University of
Edinburgh, Edinburgh, UK

Fig. 3 xxxx

Table 3 Overview of pyrolysis experiments. “X” denotes the samples that were analyzed for contaminants. In addition, the product yields for the Stage I unit are shown with the number of replicates in parentheses

		Stage I				Stage II		Stage III	
Feedstock		550 °C	Yield (%) 550 °C	700 °C	Yield (%) 700 °C	550 °C	700 °C	550 °C	700 °C
Softwood	Biochar	X	21 ± 1.1 (4)	X	19 ± 0.5 (4)	X	X	X	X
	Liquid	X	46 ± 1.1	X	45 ± 0.9				
	Gas		33 ± 2.1		36 ± 0.8				
Wheat Straw	Biochar	X	25 ± 0.5 (3)	X	23 ± 0.8 (3)	X	X	X	X
	Liquid	X	44 ± 0.3	X	44 ± 0.8				
	Gas		31 ± 0.7		33 ± 0.8				
Anaerobic Digestate	Biochar	X	25 ± 1.5 (7)						
	Liquid	X	49 ± 1.3						
	Gas		26 ± 1.2						