

# Role of cation structure in the phytotoxicity of ionic liquids: growth inhibition and oxidative stress in spring barley and common radish

Robert Biczak<sup>1</sup>  · Barbara Pawłowska<sup>1</sup> · Arkadiusz Telesiński<sup>2</sup> · Janusz Kapuśniak<sup>1</sup>

Received: 24 January 2017 / Accepted: 1 June 2017 / Published online: 22 June 2017  
© The Author(s) 2017. This article is an open access publication

**Abstract** The present study determines the influence of three ionic liquids (ILs) containing cations with diversified structure on the growth and development of spring barley seedlings and common radish leaves. Increasing amounts of 1-butyl-1-methylpyrrolidinium hexafluorophosphate [Pyrrol][PF<sub>6</sub>], 1-butyl-1-methylpiperidinium hexafluorophosphate [Piper][PF<sub>6</sub>], and 1-butyl-4-methylpyridinium hexafluorophosphate [Pyrid][PF<sub>6</sub>] were added to the soil on which both plants were cultivated. The results of this studies showed that the applied ILs were highly toxic for plants, demonstrated by the inhibition of length of plant shoots and roots, decrease of fresh mass, and increase of dry weight content. Common radish turned out to be the plant with higher resistance to the used ILs. The differences in the cation structure did not influence phytotoxicity of ILs for spring barley. Furthermore, all ILs led to a decrease of photosynthetic pigments, which was directly followed by decreased primary production in plants. Oxidative stress in plants occurred due to the presence of ILs in the soil, which was demonstrated by the increase of malondialdehyde (MDA) content, changes in the H<sub>2</sub>O<sub>2</sub> level, and antioxidant enzymes such as superoxide

dismutase (SOD), catalase (CAT), and peroxidase (POD). The changes in the chlorophyll contents and the increase of POD activity turned out to be the most significant oxidative stress biomarkers in spring barley and common radish. Both spring barley and radish exposed to ILs accumulated a large amount of fluoride ion.

**Keywords** Ionic liquids · Phytotoxicity · Oxidative stress · Antioxidant enzyme activity · Photosynthetic pigments

## Introduction

In the last decade, interest on the so-called clean technologies in chemical industry has been increased due to technological and environmental reasons. It gives rise to a continuous search for alternatives to the conventional organic solvents. Ionic liquids may become such an alternative. Their basic features, such as low melting temperatures, low vapor pressure, polarity, high thermal and electrochemical stability, high ionic conductivity, and non-flammability, prevent solvent loss and increase the safety for persons employed in the processes of industrial chemical synthesis. However, the study conducted on the influence of ionic liquids (ILs) on the aquatic and soil environment gives rise to a serious concern about the possibility of a permanent contamination of these ecosystems with the discussed compounds. IL toxicity has been observed in microorganisms, fungi, algae, terrestrial plants, invertebrates, and vertebrates (Pham et al. 2010; Petkovic et al. 2012; Peric et al. 2013; Cvjetko Bubalo et al. 2014b; Feder-Kubis and Tomczuk 2013; Messali et al. 2013).

When large amounts of ILs become available in the market and are used in numerous processes, it should be borne in mind that they will penetrate into the soil environment, where the phenomenon of soil sorption related to the presence of

Responsible editor: Philippe Garrigues

**Electronic supplementary material** The online version of this article (doi:10.1007/s11356-017-9439-x) contains supplementary material, which is available to authorized users.

✉ Robert Biczak  
r.biczak@ajd.czest.pl

<sup>1</sup> The Faculty of Mathematics and Natural Sciences, Jan Długosz University in Częstochowa, 13/15 Armii Krajowej Av, 42-200 Częstochowa, Poland

<sup>2</sup> The Faculty of Environmental Management and Agriculture, West Pomeranian University of Technology, Juliusza Słowackiego st. 17, 71-434 Szczecin, Poland

humus and inorganic colloids may be limited in the top layer of the soil, close to the roots, posing direct threat to plants. Therefore, numerous publications have been published, evaluating the level of ILs' influence on the growth and development of terrestrial plants (Biczak et al. 2010, 2013, 2014a, 2015; Matzke et al. 2008a, b; Studzińska and Buszewski 2009). In the abovementioned papers, ILs' phytotoxicity was determined primarily on the basis of plant growth inhibition, but as mentioned by Cvjetko Bubalo et al. (2014a), the IL toxicity mechanism has not yet been fully understood. Therefore, a view gains increasing popularity in scientific studies in which ILs' phytotoxicity is related to the generation of oxidative stress in plants by these compounds (Liu et al. 2013, 2014, 2015a, b, 2016a, b; Biczak 2016; Biczak et al. 2016a, 2016b; Pawłowska and Biczak 2016).

An opinion prevails in the scientific literature in which the IL phytotoxicity largely depends on the cation and the length of substituent (Cvjetko Bubalo et al. 2014b; Biczak et al. 2014a; Matzke et al. 2008a, b; Studzińska and Buszewski 2009), while to a lesser extent, it depends on the anion type (Cvjetko Bubalo et al. 2014b; Studzińska and Buszewski 2009; Liu et al. 2016a, b; Biczak et al. 2014b). However, scientific reports on the correlation of IL phytotoxicity and their cations are limited. Only Pham et al. (2016) demonstrated the higher toxicity of ILs with pyridinium cations compared to pyrrolidinium and imidazolium cations in *Pseudokirchneriella subcapitata* algae. Therefore, the objective of the present study was to determine and compare the toxic effects of ILs with different cation structures to spring barley and common radish. The study focused on relatively popular ILs, which are used in chemical synthesis, electrochemistry, and biotechnological processes, containing hexafluorophosphate anion and the following cations: 1-butyl-1-methylpyrrolidinium, 1-butyl-1-methylpiperidinium, and 1-butyl-4-methylpyridinium (Bae et al. 2013; Atta et al. 2016; Elgharbawy et al. 2016). In order to compare the phytotoxicity of these ILs and at the same time demonstrate the effects of cations on IL toxicity, apart from traditional phytotoxicity biomarkers including shoot length, root length, and fresh and dry weight yield, the present study also examined the oxidative stress indicators in the seedlings of spring barley and radish such as the malondialdehyde (MDA) content, H<sub>2</sub>O<sub>2</sub>, photosynthetic pigments, and activity of superoxide dismutase (SOD), peroxidase (POD), and catalase (CAT). Since the used ILs contain PF<sub>6</sub><sup>-</sup> anions, the effect of fluoride ions was evaluated for both spring barley and common radish as fluoroacetate, a harmful compound to plants, may be formed when fluoride is absorbed from the soil (Baunthiyal and Pandey 2012). The choice of spring barley for the study was dictated by the evidence that it is the fourth most common cereal species in production and acreage, and radish is a popular vegetable, enriching the human diet with a number of microelements and macroelements and vitamins (Schubert and Jähren 2011; Dragišić Maksimović et al. 2013; Arias-Baldrich et al. 2015).

## Materials and methods

### Ionic liquids

All ILs used in these studies were purchased from Sigma-Aldrich Chemical Co. Chemical structures and abbreviations of ILs 1-butyl-1-methylpyrrolidinium hexafluorophosphate (≥97.5% purity), 1-butyl-1-methylpiperidinium hexafluorophosphate (≥97.5% purity), and 1-butyl-4-methylpyridinium hexafluorophosphate (≥97.0% purity) are presented in Fig. 1.

### Evaluation of ILs' phytotoxicity

Determination of phytotoxicity of ILs with different cation structures was performed according to the OECD/OCDE guidelines (2006). The following concentrations of ILs were used: 0, 1, 10, 50, 100, 400, 700, and 1000 mg of compound per 1 kg of dry weight (DW) of the soil. When preparing the abovementioned concentrations, appropriate amount of ILs was dissolved in acetone and subsequently mixed with quartz sand. After overhead evaporation of acetone, quartz sand containing ILs was mixed with the soil. Three independent samples were prepared for each concentration of IL.

Following the above procedure, control samples were prepared by adding acetone to the sand, but without ILs. In this study, loam was used as soil. It contained about 11% of fraction with a diameter of <0.02 mm and organic carbon of 9.5 g kg<sup>-1</sup>. The pH was 6.1. Plastic pots were filled with the prepared medium; thereafter, 20 seeds of spring barley (*Hordeum vulgare*) and common radish (*Raphanus sativus* L. *radicula* Pers.) derived from the same source were added. Seed germination and seedlings' growth (14 days) were carried out under strictly controlled conditions: soil moisture, 70% ppw; temperature, 20 ± 2 °C; and constant illumination, 170 μmol m<sup>-2</sup> s<sup>-1</sup> for 16-h day/8-h night. The experiments were conducted in a vegetation hall, which belongs to the Department of Biochemistry and Ecotoxicology at Jan Długosz University in Częstochowa.

Phytotoxicity of the studied ILs for spring barley and common radish was estimated based on, among others, the yield of fresh weight of seedlings, dry weight content, and length of




Fig. 1 Structures of the analyzed ILs

the shoots and roots. The inhibition factor of fresh weight and lengths of shoots and roots were calculated according to the study published by Wang et al. (2009). Using non-linear regression analysis,  $EC_{50}$  was estimated based on the calculated inhibition, with the GraphPad Prism software (GraphPad Software, Inc., La Jolla, CA, USA). Furthermore, both germination potential (GP) and germination rate (GR) of spring barley and common radish seeds were determined. The seeds with longer than 2-mm germ were considered germinated (Liu et al. 2014).

In fresh plant material (spring barley seedlings and common radish leaves), the content of all assimilation pigments, malondialdehyde,  $H_2O_2$ , and enzymes' activity was determined. In the dried material, the analysis of total fluoride content was performed. Samples treated with high concentrations of ILs were not included in some analyses because the growth inhibition of spring barley and radish shoots was extremely strong at these concentrations.

#### Determination of total fluoride content

Based on the method described by Eyde (1982), the total fluoride content in spring barley seedlings and common radish leaves was assessed. Dried and ground samples were fused in nickel crucibles with sodium hydroxide. Water, diluted hydrochloric acid, and citrate buffer solution were added to the melt. Fluoride concentration was determined in the presence of TISAB III buffer using the potentiometric method with an Orion Research ion-selective electrode. The total fluoride content was expressed in  $mg\ kg^{-1}$  of DW.

#### Determination of assimilation pigment content

According to the methodology presented by Oren et al. (1993), the content of assimilation pigments was determined by spectrophotometry. With the addition of 80% acetone solution cooled to 4 °C, weighed portion (200 mg) of fresh spring barley seedlings and common radish leaves was homogenized. The resulting extract was transferred into centrifuge tubes and left in the dark for 24 h. The extract was then centrifuged, and in the filtrate, the content of assimilation dyes was determined by measuring the absorbance at 470, 647, and 664 nm. The contents of chlorophylls and carotenoids were expressed in  $mg\ g^{-1}$  of dry plant weight (DW).

#### Determination of malondialdehyde and hydrogen peroxide ( $H_2O_2$ )

With the addition of 0.1% trichloroacetic acid solution cooled to 4 °C, 500 mg of fresh spring barley seedlings and common

radish leaves were homogenized. After centrifugation, MDA and  $H_2O_2$  content were determined in the obtained supernatant according to the procedures described by Hodges et al. (1999) and Singh et al. (2007), respectively. As a substrate to determine MDA, thiobarbituric acid was used and the MDA content was determined by measuring the absorbance at 532 and 600 nm. In order to determine the content of  $H_2O_2$ , the absorbance was measured at a wavelength of 390 nm for the reaction mixture consisting of supernatant, phosphate buffer (pH 7.0), and potassium iodide. The contents of MDA and  $H_2O_2$  were calculated using the extinction coefficient equaling  $155\ mM^{-1}\ cm^{-1}$  and expressed in  $\mu mol\ g^{-1}$  fresh weight (FW).

#### Determination of superoxide dismutase, catalase, and peroxidase activity

For the determination of antioxidant enzyme activity, enzymatic extracts were obtained by homogenizing the weighed portion (500 mg) of fresh spring barley seedlings and common radish leaves with the addition of cooled (4 °C) extraction mixture. The mixture contained phosphate buffer (pH 7.4), 1 mM EDTA solution, and 0.1% polyvinylpyrrolidone solution. After centrifugation, the obtained supernatant was also used to determine the protein content.

SOD [EC 1.15.1.1] activity was determined spectrophotometrically by measuring the degree of reduction of nitroblue tetrazolium (NBT) by superoxide anion formed as a result of photochemical reduction of riboflavin in the presence of light (Giannopolitis and Ries 1977). NBT reduction is inhibited by SOD. For determining the activity of SOD, measurement of absorbance of reaction mixture at a wavelength of 560 nm was performed. SOD activity was expressed as  $U\ mg^{-1}$  protein, where 1 U is the amount of enzyme that induces 50% inhibition of NBT reduction.

CAT activity [EC 1.11.1.6] was determined by the decomposition of  $H_2O_2$  by this enzyme during 15 min, and  $H_2O_2$  remaining in the reaction mixture was titrated with 0.01 N  $KMnO_4$  (Kar and Mishra 1976). CAT activity was expressed as  $U\ mg^{-1}$  protein  $min^{-1}$ .

The activity of POD [1.11.1.7] was determined spectrophotometrically by the rate of guaiacol oxidation in the presence of  $H_2O_2$  via enzyme occurring in a given volume of extract during 1 min (Abassi et al. 1998). Absorbance measurement of the reaction mixture at a wavelength of 470 nm was performed, in order to measure the activity of POD. POD activity was expressed as  $U\ mg^{-1}$  protein  $min^{-1}$ .

The total protein content necessary to calculate the activity of SOD, CAT, and POD was determined by the Bradford (1976) assay.


### Statistical analysis

Using two-way analysis of variance, results were subjected to statistical analysis. Tukey’s test with  $p < 0.05$  was used to determine the significance of differences. Data present in tables and figures are expressed as means  $\pm$  standard deviation obtained from three measurement replicates.

### Results and discussion

#### Phytotoxicity assay

The analysis of the obtained results on the influence of the used ILs on the growth and development of spring barley and common radish revealed that all the tested compounds



**Fig. 2** The inhibition rate (%) for shoot length, root length, and fresh weight of spring barley and common radish after exposure to ILs

**Table 1** The EC<sub>50</sub> values and 95% confidence intervals for spring barley and common radish following exposure to ILs

Biomarkers	[Pyrrol][PF <sub>6</sub> ]		[Piper][PF <sub>6</sub> ]		[Pyrid][PF <sub>6</sub> ]	
	EC <sub>50</sub> (mg kg <sup>-1</sup> of soil DW)	95% Confidence intervals	EC <sub>50</sub> (mg kg <sup>-1</sup> of soil DW)	95% Confidence intervals	EC <sub>50</sub> (mg kg <sup>-1</sup> of soil DW)	95% Confidence intervals
Spring barley						
Root length	88.97	71.24–111.10	114.9	74.64–177.00	97.78	69.63–137.30
Shoot length	10.16	7.96–12.98	19.86	14.14–27.88	14.49	12.08–19.87
Fresh weight	10.04	7.55–11.67	11.40	8.66–16.68	10.31	9.01–12.10
Common radish						
Root length	137.20	72.07–156.10	120.40	28.64–255.80	101.40	67.41–152.70
Shoot length	228.50	102.80–387.20	149.60	35.51–300.60	90.87	50.92–162.10
Fresh weight	172.00	84.83–248.70	115.20	79.07–168.00	129.60	82.20–204.20

exhibited phytotoxic effect. This is demonstrated by the decrease of seedlings' fresh weight yield, decrease of the length of their shoots and roots, and the calculated EC<sub>50</sub> values on the basis of inhibition of these toxicity biomarkers (Fig. 2, Table 1).

The observed phytotoxicity was significantly influenced by ILs' concentrations. This is confirmed by the reports of other authors, who also point out to the strict dependency of the level of phytotoxicity on the used ILs' concentration. Numerous papers (Biczak et al. 2010, 2013, 2014a, 2015; Cvjetko Bubalo et al. 2014a; Liu et al. 2013, 2014, 2015a, b; Biczak 2016; Pawłowska and Biczak 2016; Wang et al. 2009) demonstrate the linear relationship between phytotoxicity of quaternary ammonium salts (QASs) and ILs and the concentration of the compound in soil. Such strong correlations are observed mainly for the high concentrations of these chemicals, whereas ILs used in low concentrations may have a stimulatory effect on the growth and development of plants (Liu et al. 2013, 2015b, 2016b).

Spring barley was very sensitive to the used ILs, and the plant material could only be obtained in the case of ILs' concentrations of 1 and 10 mg kg<sup>-1</sup> of the soil DW. Common radish exhibited higher resistance to the tested ILs (Fig. 2, Table 1). The difference in the sensitivity of spring barley and common radish observed in the present study may stem from the reaction of the root systems of these plants to the tested salts. At higher IL concentrations (50–1000 mg kg<sup>-1</sup> of soil DW), over 90% spring barley root growth inhibition was observed, which effectively disabled the plants to absorb and transport water and nutrients. As reported by Chapman et al. (2012), proper root development determined the optimum growth and development of each plant. Such root inhibition was not observed for common radish.

It is shown that there is no significant effect of cation structures on the phytotoxicity of the tested ILs to spring barley and common radish. The results of shoot length, root length, fresh weight, and EC<sub>50</sub> values of both plants showed no remarkable

differences (Fig. 2, Table 1). This is further illustrated by the photographs of the seedlings cultivated on the soil with increasing ILs' concentration, presented in Supplementary Materials (Suppl. Figs 1 and 2). Currently, the obtained results cannot be compared to the other studies describing the IL effect on terrestrial plants. Only Pham et al. (2016) compared the effect of pyridinium, pyrrolidinium, and imidazolium bromides on the growth of the *P. subcapitata* algae and stated that the IL with the pyridinium cation was more toxic to the algae than the other compounds. Cho et al. (2008) evaluated toxicity of imidazolium, pyridinium, pyrrolidinium, phosphonium, and ammonium ILs on the algae, *Selenastrum capricornutum*, and the obtained results showed that the pyrrolidinium cation was the least toxic. However, Stolte et al. (2007) proved higher toxicity of ILs containing aromatic cation for duckweed (*Lemna minor*) and microorganisms. The study carried out by Couling et al. (2006) pointed at the fact that the higher amount of nitrogen atoms in a cation ring caused the higher toxicity of ionic liquids for aquatic organisms. However, these studies were conducted in an aquatic environment, whereas our study was conducted in the soil, which is a very complex environment for toxicity study. The obtained results can be influenced by numerous factors, such as pH of the soil, content of humus substances, soil colloids, and sorption level (Cvjetko Bubalo et al. 2014b; Matzke et al. 2008a, 2009; Mrozik et al. 2009; Studzińska et al. 2009).

Moreover, the present study determined the influence of ILs with different cation on the dry weight content in spring barley seedlings and common radish leaves. In both plants, an increase of dry weight was observed, correlated to the increase of ILs' concentration in the soil (Table 2).

High ionic liquid concentrations lead to soil salinity, which, in turn, disturbs the water metabolism in plants, leading to the observed increased level of dry weight in both experimental plants. Biczak et al. (2013, 2014a, 2015, 2016b), Biczak (2016), Matusiak et al. (2013), and Pawłowska and Biczak (2016) came to analogous conclusions in their studies on the

**Table 2** Effect of ILs on the dry weight ( $\text{g g}^{-1}$  FW) in spring barley seedlings and common radish leaves

Dry weight content ( $\text{g g}^{-1}$ FW) in spring barley seedlings and common radish leaves			
IL concentration ( $\text{mg kg}^{-1}$ of soil DW)	[Pyrrol][PF <sub>6</sub> ]	[Piper][PF <sub>6</sub> ]	[Pyrid][PF <sub>6</sub> ]
Spring barley			
0	0.0787 ± 0.0013c	0.0757 ± 0.0032c	0.0763 ± 0.0014c
1	0.0793 ± 0.0041c	0.0762 ± 0.0019c	0.0760 ± 0.0008c
10	0.1047 ± 0.0049a	0.0868 ± 0.0020ba	0.0971 ± 0.0105ab
50	–	–	–
100	–	–	–
400	–	–	–
700	–	–	–
1000	–	–	–
Common radish			
0	0.0609 ± 0.0025ijkl	0.0668 ± 0.0122hijk	0.0508 ± 0.0005kl
1	0.0533 ± 0.0017kl	0.0594 ± 0.0043jkl	0.0494 ± 0.0018l
10	0.0622 ± 0.0041ijkl	0.0594 ± 0.0047jkl	0.0541 ± 0.0024kl
50	0.0765 ± 0.0064hc	0.0720 ± 0.0010hc	0.0719 ± 0.0057hij
100	0.0952 ± 0.0090g	0.0766 ± 0.0040g	0.0804 ± 0.0057gh
400	0.1786 ± 0.0033f	0.1748 ± 0.0130f	0.2052 ± 0.0037dc
700	0.2444 ± 0.0010c	0.2045 ± 0.0066c	0.2209 ± 0.0051d
1000	0.2596 ± 0.0045bc	0.2878 ± 0.0012a	0.2675 ± 0.0101b

Data are expressed as a mean ± SD of three replicates for each concentration. Values denoted by the same letters in the columns do not differ statistically at  $p < 0.05$

determination of toxicity of different chemicals for terrestrial plants.

Increase in the concentration of [Pyrrol][PF<sub>6</sub>], [Piper][PF<sub>6</sub>], and [Pyrid][PF<sub>6</sub>] in the soil led to a systematic decrease of GP

**Table 3** Effect of ILs on the fluoride ( $\text{mg kg}^{-1}$  DW) content in spring barley seedlings and common radish leaves

The fluoride content ( $\text{mg kg}^{-1}$ FW) in spring barley seedlings and common radish leaves			
IL concentration ( $\text{mg kg}^{-1}$ of soil DW)	[Pyrrol][PF <sub>6</sub> ]	[Piper][PF <sub>6</sub> ]	[Pyrid][PF <sub>6</sub> ]
Spring barley			
0	4.306 ± 0.120a	4.338 ± 0.037a	4.279 ± 0.228a
1	4.359 ± 0.170a	4.369 ± 0.076a	4.351 ± 0.151a
10	4.287 ± 0.090a	4.396 ± 0.220a	4.477 ± 0.175a
50	–	–	–
100	–	–	–
400	–	–	–
700	–	–	–
1000	–	–	–
Common radish			
0	2.094 ± 0.130g	2.073 ± 0.052g	2.109 ± 0.090f
1	2.132 ± 0.158g	2.102 ± 0.100g	2.279 ± 0.165f
10	2.699 ± 0.306fg	3.274 ± 0.240f	2.731 ± 0.077fg
50	5.253 ± 0.207e	5.863 ± 0.126e	5.140 ± 0.129e
100	7.671 ± 0.211d	8.010 ± 0.144d	7.616 ± 0.012d
400	9.510 ± 0.179c	9.805 ± 0.452c	9.218 ± 0.096c
700	12.106 ± 0.172b	12.298 ± 0.678b	12.104 ± 0.198b
1000	14.386 ± 0.318a	15.103 ± 0.396a	14.584 ± 0.256a

Data are expressed as a mean ± SD of three replicates for each concentration. Values denoted by the same letters in the columns do not differ statistically at  $p < 0.05$

**Table 4** Effect of ILs on photosynthetic pigment content ( $\text{mg g}^{-1}$  DW) in spring barley seedlings and common radish leaves

Concentration of ILs ( $\text{mg kg}^{-1}$ of soil DW)	Pigments ( $\text{mg g}^{-1}$ DW)							
	Chla	Chlb	Car	Chla + Chlb	Chla/Chlb	Chl(a + b)/Car		
<b>Spring barley</b>								
[Pyrrol][PF <sub>6</sub> ]	0	13.196 ± 0.341de	3.520 ± 0.042c	3.189 ± 0.058de	16.716 ± 0.377d	3.749 ± 0.064a	5.242 ± 0.024b	
	1	13.923 ± 0.665bcd	3.498 ± 0.194c	3.486 ± 0.115b	17.420 ± 0.512cd	3.995 ± 0.409a	4.998 ± 0.025e	
	10	10.374 ± 0.157f	2.684 ± 0.005e	2.568 ± 0.035f	13.058 ± 0.156f	3.865 ± 0.059a	5.086 ± 0.009cdc	
	50	–	–	–	–	–	–	
	100	–	–	–	–	–	–	
	400	–	–	–	–	–	–	
	700	–	–	–	–	–	–	
	1000	–	–	–	–	–	–	
	[Piper][PF <sub>6</sub> ]	0	15.092 ± 0.110a	3.989 ± 0.051a	3.461 ± 0.011b	19.201 ± 0.147a	3.814 ± 0.038a	5.548 ± 0.025a
		1	14.381 ± 0.017b	3.847 ± 0.013ab	3.463 ± 0.015b	18.228 ± 0.027b	3.738 ± 0.010a	5.264 ± 0.015b
10		13.522 ± 0.283cd	3.626 ± 0.051bc	3.294 ± 0.058cd	17.148 ± 0.329cd	3.729 ± 0.038a	5.205 ± 0.018bc	
50		–	–	–	–	–	–	
100		–	–	–	–	–	–	
400		–	–	–	–	–	–	
700		–	–	–	–	–	–	
1000		–	–	–	–	–	–	
[Pyrid][PF <sub>6</sub> ]		0	15.420 ± 0.094a	4.055 ± 0.014a	3.942 ± 0.057a	19.476 ± 0.082a	3.803 ± 0.035a	4.941 ± 0.087e
		1	14.018 ± 0.151bc	3.654 ± 0.083bc	3.426 ± 0.033bc	17.672 ± 0.223bc	3.837 ± 0.060a	5.158 ± 0.112bcd
	10	12.547 ± 0.102e	3.116 ± 0.051d	3.115 ± 0.005e	15.663 ± 0.124e	4.027 ± 0.067a	5.028 ± 0.045de	
	50	–	–	–	–	–	–	
	100	–	–	–	–	–	–	
	400	–	–	–	–	–	–	
	700	–	–	–	–	–	–	
	1000	–	–	–	–	–	–	
	<b>Common radish</b>							
	[Pyrrol][PF <sub>6</sub> ]	0	11.732 ± 0.065a	4.001 ± 0.027ab	3.193 ± 0.081a	15.732 ± 0.090a	2.932 ± 0.010def	4.930 ± 0.151abcde
1		9.012 ± 0.294def	3.118 ± 0.093de	2.374 ± 0.128cde	12.130 ± 0.385ef	2.890 ± 0.021ef	5.116 ± 0.180abcde	
10		8.702 ± 0.029fg	2.773 ± 0.031fg	2.338 ± 0.013def	11.475 ± 0.047gh	3.139 ± 0.033cdef	4.909 ± 0.032abcde	
50		7.521 ± 0.027i	2.570 ± 0.092g	2.011 ± 0.163fg	10.091 ± 0.108i	2.929 ± 0.102def	5.038 ± 0.384abcde	
100		7.665 ± 0.120i	2.642 ± 0.030g	1.934 ± 0.015g	10.308 ± 0.094i	2.902 ± 0.076ef	5.331 ± 0.066abc	
400		5.051 ± 0.119j	1.598 ± 0.047h	1.302 ± 0.030h	6.648 ± 0.164j	3.162 ± 0.033cdef	5.106 ± 0.060abcde	
700		2.187 ± 0.145m	0.595 ± 0.078i	0.641 ± 0.025jk	2.782 ± 0.067m	3.745 ± 0.192bcdef	4.343 ± 0.270efg	
1000		1.737 ± 0.068no	0.401 ± 0.026j	0.522 ± 0.006k	2.137 ± 0.045no	4.356 ± 0.462bcd	4.096 ± 0.120fg	
[Piper][PF <sub>6</sub> ]		0	10.069 ± 0.039c	3.282 ± 0.029cde	2.591 ± 0.016cd	13.351 ± 0.062d	3.068 ± 0.022cdef	5.153 ± 0.019abcde
		1	10.589 ± 0.260b	3.502 ± 0.058c	2.716 ± 0.068bc	14.091 ± 0.317c	3.033 ± 0.026cdef	5.188 ± 0.016abcde
	10	10.174 ± 0.028c	3.335 ± 0.011cd	2.613 ± 0.005cd	13.508 ± 0.017d	3.051 ± 0.019cdef	5.171 ± 0.009abcde	
	50	9.253 ± 0.079de	2.600 ± 0.016a	2.147 ± 0.004efg	11.854 ± 0.064fg	3.559 ± 0.052bcdef	5.520 ± 0.023ab	
	100	8.908 ± 0.152ef	2.550 ± 0.003g	2.172 ± 0.037efg	11.458 ± 0.155gh	3.493 ± 0.055bcdef	5.276 ± 0.029abcd	
	400	4.850 ± 0.209j	1.170 ± 0.044i	1.243 ± 0.049hi	6.020 ± 0.253k	4.145 ± 0.025bcdef	4.842 ± 0.013abcde	
	700	3.909 ± 0.079k	0.929 ± 0.019i	1.056 ± 0.022hi	4.838 ± 0.096l	4.208 ± 0.048bcde	4.581 ± 0.013cdefg	
	1000	2.064 ± 0.027mn	0.330 ± 0.088j	0.626 ± 0.079jk	2.394 ± 0.075mno	6.589 ± 0.922a	3.871 ± 0.537g	
	[Pyrid][PF <sub>6</sub> ]	0	11.601 ± 0.113a	4.140 ± 0.149a	3.161 ± 0.066a	15.741 ± 0.258a	2.804 ± 0.075ef	4.982 ± 0.183abcde
		1	10.893 ± 0.083b	3.813 ± 0.121b	3.013 ± 0.075ab	14.705 ± 0.168b	2.859 ± 0.077ef	4.882 ± 0.140abcde
10		9.337 ± 0.018d	3.121 ± 0.155de	2.480 ± 0.078cde	12.459 ± 0.167e	2.996 ± 0.142cdef	5.028 ± 0.229abcde	
50		8.313 ± 0.022gh	3.018 ± 0.256ef	2.267 ± 0.460defg	11.331 ± 0.263gh	2.768 ± 0.233f	5.113 ± 0.849abcde	
100		8.222 ± 0.222h	2.778 ± 0.071fg	1.943 ± 0.025g	11.001 ± 0.217h	2.961 ± 0.120def	5.663 ± 0.163a	

**Table 4** (continued)

Concentration of ILs (mg kg <sup>-1</sup> of soil DW)	Pigments (mg g <sup>-1</sup> DW)					
	Chla	Chlb	Car	Chla + Chlb	Chla/Chlb	Chl(a + b)/Car
400	3.330 ± 0.126l	0.989 ± 0.054i	0.913 ± 0.056ij	4.319 ± 0.180l	3.369 ± 0.059bcdef	4.747 ± 0.399bcdef
700	2.139 ± 0.050m	0.489 ± 0.047j	0.548 ± 0.013jk	2.629 ± 0.003mn	4.403 ± 0.500bc	4.504 ± 0.095cdefg
1000	1.612 ± 0.021o	0.336 ± 0.008j	0.441 ± 0.048k	1.948 ± 0.015o	4.796 ± 0.177b	4.452 ± 0.491defg

Data are expressed as a mean ± SD of three replicates for each concentration. Values denoted by the same letters in the columns do not differ statistically at *p* < 0.05

and GR values of spring barley seeds. In the case of common radish grown on the soil with the addition of ILs, a small decrease in the germination potential and germination rate of this plant seeds was noted, especially after the application of the highest concentrations of [Pyrrol][PF<sub>6</sub>] (Suppl. Table 1). Liu et al. (2014) described a similar decrease in germination capacity of wheat and spring barley seeds under the influence of imidazolium ILs.

**Effect of ILs on fluoride content**

In the discussed studies, the content of fluoride ions in seedlings of spring barley and radish leaves was determined (Table 3).

Common radish accumulated large amounts of fluoride ions in its leaves, and the observed increase in the level of these anions was directly proportional to the increase in concentration of all ILs in the soil. In case of application of all studied ILs in a dosage of 1000 mg kg<sup>-1</sup> of soil DW, the amount of fluoride in plants was approximately seven times higher than in the control groups. Such changes were not established for spring barley, probably because the analyses were possible to conduct only for the objects with the concentration of 10 mg ILs per 1 kg of soil DW. The effect of toxic fluoride on plants is visible in the form of chlorosis, peripheral necrosis, leaf distortion and malformation, and abnormal fruit development (Pandey et al. 2014). Telesiński and Śnioszek (2009) determined that fluorine has a negative influence on the assimilation and photosynthesis processes in plants. These phenomena stem from the destructive influence of F<sup>-</sup> on chloroplasts.

**Effect of ILs on pigment content**


An analysis of the results on the effect of ILs on photosynthetic pigments’ level in spring barley seedlings and common radish leaves demonstrates the inhibitory influence of these salts on the level of chlorophylls and carotenoids. All the ILs used in the experiment led to a systematic decrease in the content of photosynthetic pigments in both plants, which was correlated to the increase of these substances in the soil (Table 4).

A similar decrease in the photosynthetic pigment content in spring barley and wheat seedlings and radish and broad bean leaves, duckweed, and algae under IL influence was also observed by Cvjetko Bubalo et al. (2014a), Biczak (2016), Biczak et al. (2016a), Pawłowska and Biczak (2016), Wang et al. (2009), Liu et al. (2014), Zhang et al. (2013), and Ma et al. (2010). The cited authors demonstrate that the presence of ILs in the environment generates a high level of oxidative stress in plants, which is linked to the elevated reactive oxygen species (ROS) production (Anjaneyulu et al. 2014; Di Baccio et al. 2014; Oukarroum et al. 2015). Some authors (Cvjetko Bubalo et al. 2014a; Biczak 2016; Biczak et al. 2016a, b) also report that chlorophyll content constitutes the most important biomarker of oxidative stress because its changes are directly correlated with the inhibition of the growth and yield of plants. Islam et al. (2014) and Herman et al. (1998) believe that chlorophyll level reflects the health of plant leaves.

Besides the content of photosynthetic pigments, the following indexes are also used to evaluate the physiological changes: Chla/Chlb ratio and Chl(a + b)/Car ratio. The increase of Chla/Chlb value and decrease of Chl(a + b)/Car value indicate the occurrence of oxidative stress in plants (Arias-Baldrich et al. 2015; Chen et al. 2014). In addition, the decrease of Chl(a + b)/Car value indicates antioxidant defense of the plant organism, by increasing carotenoid content. Carotenoids are efficient ROS scavengers, thus protecting PSI and PSII photosystems (Arias-Baldrich et al. 2015; Wang et al. 2009; Chen et al. 2014; Gengmao et al. 2015). No statistically significant differences in the values of both mentioned indexes were determined in the spring barley seedlings. In the case of common radish, value of the Chla/Chlb index increased on the samples with the highest ILs’ concentration; additionally, for the same samples, a decrease of the Chl(a + b)/Car value was observed in comparison with the control (Table 4). This indicates oxidative stress in radish leaves and suggests the attempt of antioxidant defense carried out by the plant.

**Effect of ILs on MDA and H<sub>2</sub>O<sub>2</sub> content**

The peroxidation level of protein-lipid membranes in plant organisms is commonly determined on the basis of



**Fig. 3** Effect of ILs on MDA ( $\mu\text{mol g}^{-1}$  FW) and  $\text{H}_2\text{O}_2$  ( $\mu\text{mol g}^{-1}$  FW) content in spring barley seedlings and common radish leaves. Data are expressed as a mean  $\pm$  SD of three replicates for each concentration. Values denoted by the same letters for each ILs do not differ statistically at  $p < 0.05$

malondialdehyde content. Thus, the MDA content is one of the most important indices, always determined in the study of oxidative stress level in plants subjected to the influence of different stress factors (Liu et al. 2013, 2014; Rosalie et al. 2015; Radošević et al. 2015). In the present paper, no considerable changes of MDA content of spring barley seedlings were observed between controls and samples cultivated on the soil with the addition of ILs. However, a significant increase of MDA content in the common radish leaves, which at the highest IL concentrations averaged 250–400% in comparison to the control, was observed (Fig. 3). The greatest changes of MDA level were caused by the ionic liquid containing the aromatic cation [Pyrid][PF<sub>6</sub>].

In the available literature, studies can be found describing the effect of ILs on the biochemical and physiological changes in plants, reporting increase of MDA content in plant cells (Cvjetko Bubalo et al. 2014a; Liu et al. 2013, 2014, 2015a, 2016a, b; Biczak et al. 2016a). The authors explain that the observed tendencies in the changes of MDA level were caused by oxidative stress generated by high concentrations of ILs.

Another biomarker indicating oxidative stress in plants is the  $\text{H}_2\text{O}_2$  accumulation in their cells. Hydrogen peroxide is the most stable chemical molecule of all ROS, characterized by the capability for a rapid penetration of cellular membranes. The increase of  $\text{H}_2\text{O}_2$  content in plant cells is observed in the conditions of intensified superoxide anion radical detoxification and in the situation when enzymatic mechanisms of  $\text{H}_2\text{O}_2$  detoxification fail in plants (Liu et al. 2013, 2015b; Sánchez-Rodríguez et al. 2010; Kumar et al. 2013; Demidchik 2015). Zhang et al. (2013), Liu et al. (2014), and Biczak (2016) found that  $\text{H}_2\text{O}_2$  accumulation depends on the ILs' concentration in the environment of plant vegetation.

The results revealed that extremely high accumulation of  $\text{H}_2\text{O}_2$  was found in common radish cells, and the observed increase was positively correlated with the concentration of the used ionic liquid. The highest increase in the level of hydrogen peroxide was observed when high concentrations of [Pyrro][PF<sub>6</sub>] and [Pyrid][PF<sub>6</sub>] were used (Fig. 3). On the contrary,  $\text{H}_2\text{O}_2$  content in spring barley seedlings was decreased, which was the result of elevated activity of the

**Table 5** Enzymatic activities of SOD (U mg<sup>-1</sup> protein), CAT (U mg<sup>-1</sup> protein min<sup>-1</sup>), and POD (U mg<sup>-1</sup> protein min<sup>-1</sup>) in spring barley seedlings and common radish leaves treated with ILs

Concentration of ILs (mg kg <sup>-1</sup> of soil DW)		The activity of enzymes			
		Superoxide dismutase (U mg <sup>-1</sup> protein)	Catalase (U mg <sup>-1</sup> protein min <sup>-1</sup> )	Peroxidase (U mg <sup>-1</sup> protein min <sup>-1</sup> )	
Spring barley					
[Pyrrol][PF <sub>6</sub> ]	0	9.901 ± 0.943bc	0.0483 ± 0.0014cd	14.868 ± 0.212cd	
	1	10.113 ± 0.168abc	0.0458 ± 0.0005c	16.115 ± 0.082c	
	10	8.950 ± 0.411c	0.0591 ± 0.0020a	38.649 ± 2.813a	
	50	–	–	–	
	100	–	–	–	
	400	–	–	–	
	700	–	–	–	
	1000	–	–	–	
	[Piper][PF <sub>6</sub> ]	0	11.223 ± 0.998ab	0.0434 ± 0.0014ef	9.897 ± 0.360ef
		1	11.769 ± 0.213a	0.0448 ± 0.0014de	12.215 ± 0.175de
10		11.781 ± 0.280a	0.0519 ± 0.0025b	26.976 ± 0.411b	
50		–	–	–	
100		–	–	–	
400		–	–	–	
700		–	–	–	
1000		–	–	–	
[Pyrid][PF <sub>6</sub> ]		0	11.174 ± 0.787ab	0.0286 ± 0.0016e	6.424 ± 0.100g
		1	11.373 ± 0.522ab	0.0293 ± 0.0016e	8.074 ± 0.288fg
	10	9.103 ± 0.225c	0.0448 ± 0.0013cd	13.118 ± 0.454d	
	50	–	–	–	
	100	–	–	–	
	400	–	–	–	
	700	–	–	–	
	1000	–	–	–	
	Common radish				
	[Pyrrol][PF <sub>6</sub> ]	0	10.915 ± 0.085ghijk	0.0255 ± 0.0014efghi	0.897 ± 0.019kl
1		11.415 ± 0.712ghij	0.0244 ± 0.001bghi	0.827 ± 0.085kl	
10		10.823 ± 0.455ghijk	0.0231 ± 0.0026ij	0.895 ± 0.005kl	
50		10.579 ± 0.207hijk	0.0281 ± 0.0023defgh	2.409 ± 0.056hi	
100		10.501 ± 0.398ijk	0.0320 ± 0.0306cd	2.739 ± 0.177gh	
400		19.271 ± 0.384bc	0.0370 ± 0.0009b	9.510 ± 0.227de	
700		17.931 ± 0.647cde	0.0334 ± 0.0008bc	9.901 ± 0.380cd	
1000		17.734 ± 0.175de	0.0333 ± 0.0003bc	10.521 ± 0.031c	
[Piper][PF <sub>6</sub> ]		0	12.197 ± 0.612fg	0.0164 ± 0.0001kl	1.161 ± 0.086jkl
		1	11.953 ± 0.379fgh	0.0148 ± 0.0010l	1.081 ± 0.167kl
	10	12.698 ± 0.490f	0.0147 ± 0.0014l	1.387 ± 0.059jk	
	50	12.340 ± 0.068f	0.0241 ± 0.0014hi	1.911 ± 0.123ij	
	100	12.548 ± 0.165f	0.0295 ± 0.0014cde	3.208 ± 0.083g	
	400	22.380 ± 0.574a	0.0417 ± 0.0009a	12.551 ± 0.358b	
	700	19.489 ± 0.800b	0.0283 ± 0.0009defg	14.211 ± 0.206a	
	1000	18.654 ± 0.496bcd	0.0191 ± 0.0009jk	14.666 ± 0.370a	
	[Pyrid][PF <sub>6</sub> ]	0	10.058 ± 0.263jk	0.0148 ± 0.0016l	0.588 ± 0.021l
		1	11.722 ± 0.721fghi	0.0142 ± 0.0005l	0.667 ± 0.039kl
10		10.011 ± 0.050jk	0.0140 ± 0.0001l	0.710 ± 0.015kl	

**Table 5** (continued)

Concentration of ILs (mg kg <sup>-1</sup> of soil DW)	The activity of enzymes		
	Superoxide dismutase (U mg <sup>-1</sup> protein)	Catalase (U mg <sup>-1</sup> protein min <sup>-1</sup> )	Peroxidase (U mg <sup>-1</sup> protein min <sup>-1</sup> )
50	9.732 ± 0.134k	0.0226 ± 0.0023ij	1.126 ± 0.079kl
100	10.489 ± 0.387ijk	0.0254 ± 0.0014fghi	2.426 ± 0.094hi
400	17.112 ± 0.515e	0.0279 ± 0.0016defgh	7.301 ± 0.102f
700	17.264 ± 0.490de	0.0289 ± 0.0009def	7.860 ± 0.527f
1000	17.670 ± 0.195de	0.0251 ± 0.0009fghi	8.808 ± 0.736e

Data are expressed as a mean ± SD of three replicates for each concentration. Values denoted by the same letters in the columns do not differ statistically at  $p < 0.05$

enzymes responsible for decomposition of the ROS in comparison to the observed increased enzymatic activity of peroxidase and catalase in common radish leaves.

### Effect of ILs on antioxidant enzyme activities

Terrestrial plants have developed a system of antioxidant enzymes, which enables them to remove the excess ROS from the organism. The activities of SOD, CAT, POD, and glutathione reductase (GR) are correlated, since normally, the product of action of one of them constitutes an activator and substrate influencing another (Anjaneyulu et al. 2014; Gengmao et al. 2015; Sánchez-Rodríguez et al. 2010; Noqueiro et al. 2015).

In the scientific literature, a view prevails that the first line of defense against ROS is SOD, which decomposes the superoxide anion radical to H<sub>2</sub>O<sub>2</sub> and O<sub>2</sub>. Therefore, the activity of the enzyme is always determined in scientific studies on the verification of the effect of oxidative stress on the biochemical and physiological changes in plants (Liu et al. 2013, 2015b, 2016a, b; Biczak 2016; Biczak et al. 2016a, b; Pawłowska and Biczak 2016; Zhang et al. 2013). Results of studies presented in the literature do not allow for reaching firm conclusions on the direction of changes in the activity of SOD in plants subject to the conditions of oxidative stress generated by ILs. The analysis of the present study results demonstrated that in spring barley seedlings, no statistically significant changes of the activity of superoxide dismutase between the control and plants cultivated in the soil with ILs' addition were observed. The observed lack of changes in SOD activity in spring barley seedlings was because the analysis was performed only at the lowest concentrations of applied ILs (1 and 10 mg kg<sup>-1</sup> of soil DW). On the contrary, a considerable increase in activity of the enzyme after the use of ILs was observed in common radish leaves. The observed increase of SOD activity occurred only to certain ILs' concentration. In the highest concentrations of these salts used in the experiment, SOD activity decreased slightly; however, it always remained on the level higher than in the control (Table 5).

Similar conclusions were drawn by other authors (Cvjetko Bubalo et al. 2014a; Liu et al. 2013, 2015a, b, 2016b; Biczak 2016) who proved that the increase of SOD activity determined at lower ILs' concentrations indicates the defense of the plants against oxidative stress. On the contrary, high ionic liquid concentrations may lead to a significant damage of plant cells, which disables the introduction of antioxidant enzymes. The situation continues to reduce the possibility for antioxidant defense, eventually leading to the death of the cells and the entire plant organism.

H<sub>2</sub>O<sub>2</sub> formed as a result of superoxide anion radical dismutation is digested by CAT and POD. The reaction of direct disproportionation of H<sub>2</sub>O<sub>2</sub> to H<sub>2</sub>O and O<sub>2</sub> is conducted by catalase. Despite the fact that some authors (Anjaneyulu et al. 2014; Chen et al. 2014) consider CAT as the basic enzyme responsible for the removal of H<sub>2</sub>O<sub>2</sub> from plant cells, the data present in the scientific literature on the changes of activity of this enzyme are not sufficient to conclude the direction of such changes. Some authors (Cvjetko Bubalo et al. 2014a; Liu et al. 2014, 2016a, b; Pawłowska and Biczak 2016; Gengmao et al. 2015) demonstrated that CAT activity always increases under ILs' induced oxidative stress. However, in some studies (Liu et al. 2014; Sánchez-Rodríguez et al. 2010), a considerable decrease of the enzyme's activity was observed in plants subjected to oxidative stress. In spring barley seedlings, a slight increase of CAT activity was observed at the highest IL concentration (10 mg kg<sup>-1</sup> of soil DW) (Table 5).

The changes of POD activity are considered to be the most reliable biomarker indicating the occurrence of oxidative stress in plants. Regardless of the cause for oxidative stress, the activity of the enzyme always increases under the excess of H<sub>2</sub>O<sub>2</sub> (Liu et al. 2013; Biczak 2016; Biczak et al. 2016a, b; Anjaneyulu et al. 2014). According to Zhang et al. (2013), the observed increase of POD activity indicates stronger affinity of the enzyme to H<sub>2</sub>O<sub>2</sub> than CAT. In the discussed study, a systematic increase of POD activity was determined in spring barley seedlings and common radish leaves, growing on the soil with increasing concentration of ILs. In the highest dosages of all applied ionic liquids, the observed increase of

peroxidase activity in common radish leaves was very high and reached several hundred percent compared to the control (Table 5). However, some studies (Arias-Baldrich et al. 2015; Wang et al. 2009) have increasingly paid attention to the fact that the increase of POD activity is not entirely favorable for plant organisms. The increased POD activity may disable plant metabolism via removal of H<sub>2</sub>O<sub>2</sub> molecules, which are responsible for the cellular signaling, and lead to the damage of chlorophyll particles.

## Conclusion

The analysis of the results on the influence of various ILs [Pyrrol][PF<sub>6</sub>], [Piper][PF<sub>6</sub>], and [Piryd][PF<sub>6</sub>] on the growth and development of spring barley and radish and the oxidative stress level allowed for drawing the following conclusions including

1. The used ILs, particularly in high concentrations, were clearly phytotoxic. Phytotoxicity depended on the plant species. Spring barley turned out to be more sensitive to the tested salts than common radish.
2. There were no significant differences in phytotoxicity of ILs with differentiated structure of cations for spring barley. In the case of common radish, modification of IL cation structure affected the size of the changes observed in selected parameters of phytotoxicity and oxidative stress. It was evident especially after the introduction of high concentrations of these salts into the soil.
3. All ILs caused a decrease of photosynthetic pigments' content in spring barley seedlings and common radish leaves, which, as a consequence, caused the decrease of yield of both plants.
4. ILs led to oxidative stress, which was followed by the MDA accumulation in both plants and increase of H<sub>2</sub>O<sub>2</sub> in common radish leaves.
5. In response to stress conditions, the plants activated a system of antioxidant enzymes—SOD, CAT, and POD. The most significant enhancement of activity was observed for peroxidase, which can be considered the basic biomarker of oxidative stress in spring barley and common radish.

**Acknowledgments** The work was financed by a statutory activity subsidy from the Polish Ministry of Science and Higher Education for The Faculty of Environmental Management and Agriculture of West Pomeranian University of Technology in Szczecin and The Faculty of Mathematics and Natural Sciences of Jan Długosz University in Częstochowa.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Abassi NA, Kushad MM, Endress AG (1998) Active oxygen-scavenging enzymes activities in developing apple flowers and fruits. *Sci Horti* 74:183–194
- Anjaneyulu E, Reddy PS, Sunita MS, Kishor PBK, Meriga B (2014) Salt tolerance and activity of antioxidative enzymes of transgenic finger millet overexpressing a vacuolar H<sup>+</sup>-pyrophosphatase gene (SbVPPase) from *Sorghum bicolor*. *J Plant Physiol* 171:789–798. doi:10.1016/j.jplph.2014.02.001
- Arias-Baldrich C, Bosch N, Begines D, Feria AB, Monreal JA, García-Mauriño S (2015) Proline synthesis in barley under iron deficiency and salinity. *J Plant Physiol* 183:121–129. doi:10.1016/j.jplph.2015.05.016
- Atta NF, El-Ads EH, Ahmed YM, Galal A (2016) Determination of some neurotransmitters at cyclodextrin/ionic liquid crystal/graphene composite electrode. *Electrochim Acta* 199:319–331. doi:10.1016/j.electacta.2016.02.078
- Bae S-Y, Shim E-G, Kim D-W (2013) Effect of ionic liquid as a flame-retarding additive on the cycling performance and thermal stability of lithium-ion batteries. *J Power Sources* 244:266–271. doi:10.1016/j.jpowsour.2013.01.100
- Baunthiyal M, Pandey A (2012) Organofluorine metabolism in plants. *Fluoride* 45:78–85
- Biczak R (2016) Quaternary ammonium salts with tetrafluoroborate anion: phytotoxicity and oxidative stress in terrestrial plants. *J Hazard Mater* 304:173–185. doi:10.1016/j.hazmat.2015.10.055
- Biczak R, Bachowska B, Bałczewski P (2010) Study of phytotoxicity of ionic liquid 1-(methylthiomethylene)-3-butylimidazolium chloride. *Proc ECoPol* 4:105–114 (in Polish)
- Biczak R, Bałczewski P, Bachowska B, Pawłowska B, Kaźmierczak-Barańska J, Cieślak M, Nawrot B (2013) Phytotoxicity and cytotoxicity of imidazolium ionic liquids containing sulfur atom. *Phosphorus Sulfur Silicon Relat Elem* 188:459–461. doi:10.1080/10426507.2012.737880
- Biczak R, Bałczewski P, Pawłowska B, Bachowska B, Rychter P (2014a) Comparison of phytotoxicity of selected phosphonium ionic liquid. *Ecol Chem Eng S* 21:281–295. doi:10.2478/eces-2014-0022
- Biczak R, Pawłowska B, Bałczewski P, Rychter P (2014b) The role anion in the toxicity of imidazolium ionic liquids. *J Hazard Mater* 274:181–190. doi:10.1016/j.hazmat.2014.03.021
- Biczak R, Pawłowska B, Feder-Kubis J (2015) The phytotoxicity of ionic liquids from natural pool of (–)-menthol with tetrafluoroborate anion. *Environ Sci Pollut Res* 22:11740–11754. doi:10.1007/s11356-015-4327-8
- Biczak R, Telesiński A, Pawłowska B (2016a) Oxidative stress in spring barley and common radish exposed to quaternary ammonium salts with hexafluorophosphate anion. *Plant Physiol Biochem* 107:248–256. doi:10.1016/j.plaphy.2016.05.016
- Biczak R, Pawłowska B, Telesiński A, Ciesielski W (2016b) The effect of the number of alkyl substituents on imidazolium ionic liquids phytotoxicity and oxidative stress in spring barley and common radish seedlings. *Chemosphere* 165:519–528. doi:10.1016/j.chemosphere.2016.09.074
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities utilizing the principle of protein-dye binding. *Anal Biochem* 72:248–254
- Chapman N, Miller AJ, Lindsey K, Whalley WR (2012) Roots, water, and nutrient acquisition: let's get physical. *Trends Plant Sci* 17:701–710. doi:10.1016/j.tplants.2012.08.001
- Chen Y, Lin F, Yang H, Yue L, Hu F, Wang J, Luo Y, Cao F (2014) Effect of varying NaCl doses on flavonoid production in suspension cells of *Ginkgo biloba*: relationship to chlorophyll fluorescence, ion homeostasis, antioxidant system and ultrastructure. *Acta Physiol Plant* 36:3173–3187. doi:10.1007/s11738-014-1684-8

- Cho C-W, Jean Y-C, Pham TPT, Vijayaraghaven K, Sun Y-S (2008) The ecotoxicity of ionic liquids and traditional organic solvents on microalga *Selenastrum capricornutum*. *Ecotoxicol Environ Saf* 71:166–171. doi:10.1016/j.ecoenv.2007.07.001
- Couling DJ, Bemot RJ, Docherty KM, Dixon JNK, Maginn EJ (2006) Assessing the factors responsible for ionic liquid toxicity to aquatic organisms via quantitative structure–property relationship modeling. *Green Chem* 8:82–90. doi:10.1039/b511333d
- Cvjetko Bubalo M, Hanousek K, Radošević K, Srček VG, Jakovljević T, Radojčić Redovniković I (2014a) Imidazolium based ionic liquids: effect of different anions and alkyl chains lengths on the barley seedlings. *Ecotox Environ Saf* 101:116–123. doi:10.1016/j.ecoenv.2013.12.022
- Cvjetko Bubalo M, Radošević K, Radojčić Redovniković I, Halambek J, Srček VG (2014b) A brief overview of the potential environmental hazards of ionic liquids. *Ecotoxicol Environ Saf* 99:1–12. doi:10.1016/j.ecoenv.2013.10.019
- Demidchik V (2015) Mechanisms of oxidative stress in plants: from classical chemistry to cell biology. *Environ Exp Bot* 109:212–228. doi:10.1016/j.envexpbot.2014.06.021
- Di Baccio D, Castagna A, Tognetti R, Ranieri A, Sebastiani L (2014) Early responses to cadmium of two poplar clones that differ in stress tolerance. *J Plant Physiol* 171:1693–1705. doi:10.1016/j.jplph.2014.08.007
- Dragišić Maksimović J, Zhang J, Zeng F, Živanović BD, Shabala L, Zhou M, Shabala S (2013) Linking oxidative and salinity stress tolerance in barley: can root antioxidant enzyme activity be used as a measure of stress tolerance? *Plant Soil* 365:141–155. doi:10.1007/s11104-012-1366-5
- Elgharabawy AA, Alam MZ, Moniruzzaman M, Goto M (2016) Ionic liquid pretreatment as emerging approaches for enhanced enzymatic hydrolysis of lignocellulosic biomass. *Biochim Eng J* 109:252–267. doi:10.1016/j.bej.2016.01.021
- Eyde B (1982) Determination of fluoride in plant material with ion-selective electrode. *Fresenius Z. Anal Chem* 311:19–22
- Feder-Kubis J, Tomczuk K (2013) The effect of the cationic structures of chiral ionic liquids on the their antimicrobial activities. *Tetrahedron* 69:4190–4198. doi:10.1016/j.tet.2013.03.107
- Gengmao Z, Yu H, Xing S, Shihui L, Quanmei S, Changhai W (2015) Salinity stress increases secondary metabolites and enzyme activity in safflower. *Ind Crop Prod* 64:175–181. doi:10.1016/j.indcrop.2014.10.058
- Giannopolitis CN, Ries SK (1977) Superoxide dismutase. I. Occurrence in higher plants. *Plant Physiol* 59:309–314
- Herman B, Biczak R, Gurgul E (1998) Effect of 1,10-phenanthroline on peroxidase and catalase activity and chlorophyll, sugar, and ascorbic acid contents. *Biol Plant* 41:607–611
- Hodges DM, DeLong JM, Forney CF, Prange RK (1999) Improving the thiobarbituric acid-reactive-substances assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other interfering compounds. *Planta* 207:604–611
- Islam MK, Khanam MS, Lee SY, Alam I, Huh MR (2014) The interaction of arsenic (As) and chromium (Cr) influences growth and antioxidant status in tossa jute (*Corchorus olitorius*). *POJ* 7:499–509
- Kar M, Mishra D (1976) Catalase, peroxidase, and polyphenoloxidase activities during rice leaf senescence. *Plant Physiol* 57:315–319
- Kumar M, Reddy CRK, Jha B (2013) The ameliorating effect of Acadian marine plant extract against ionic liquids-induced oxidative stress and DNA damage in marine macroalga *Ulva lactuca*. *J Appl Phycol* 25:369–378. doi:10.1007/s10811-012-9871-8
- Liu H, Zhang S, Hu X, Chen C (2013) Phytotoxicity and oxidative stress effect of 1-octyl-3-methylimidazolium chloride ionic liquid on rice seedlings. *Environ Pollut* 181:242–249. doi:10.1016/j.envpol.2013.06.007
- Liu T, Zhu L, Xie H, Wang J, Wang J, Sun F, Wang F (2014) Effects of the ionic liquid 1-octyl-3-methylimidazolium hexafluorophosphate on the growth of wheat seedlings. *Environ Sci Pollut Res* 21:3936–3945. doi:10.1007/s11356-013-2348-8
- Liu H, Zhang S, Zhang X, Chen C (2015a) Growth inhibition and effect on photosystem by three imidazolium ionic liquids in rice seedlings. *J Hazard Mater* 286:440–448. doi:10.1016/j.hazmat.2015.01.008
- Liu T, Zhu L, Wang J, Wang J, Xie H (2015b) The genotoxic and cytotoxic effects of 1-butyl-3-methylimidazolium chloride in soil on *Vicia faba* seedlings. *J Hazard Mater* 285:27–36. doi:10.1016/j.hazmat.2014.11.028
- Liu T, Wang J, Wang J, Zhu L, Zhang J, Sun X (2016a) Growth and physiological and biochemical responses of wheat seedlings to imidazolium-based ionic liquids 1-octyl-3-methylimidazolium chloride and 1-oktyl-3-methylimidazolium bromide. *Bull Rnviron Contam Toxicol* 96:544–549. doi:10.1007/s00128-016-1747-1
- Liu T, Zhu L, Wang J, Wang J, Tan M (2016b) Phytotoxicity of imidazolium-based ILs with different anions in soil on *Vicia faba* seedlings and the influence of anions on toxicity. *Chemosphere* 145:269–276. doi:10.1016/j.chemosphere.2015.11.055
- Ma J-M, Cai L-L, Zhang B-J, Hu L-W, Li X-Y, Wang J-J (2010) Acute toxicity and effects of 1-alkyl-3-methylimidazolium bromide ionic liquids on green algae. *Ecotox Environ Saf* 73:1465–1469. doi:10.1016/j.ecoenv.2009.10.004
- Matusiak A, Lewkowski J, Rychter P, Biczak R (2013) Phytotoxicity of new furan-derived aminophosphonic acid, N-aryl furaldimines and 5-nitrofuraldimine. *J Agric Food Chem* 61:7673–7678. doi:10.1021/jf402401z
- Matzke M, Stolte S, Arning J, Uebers U, Filser J (2008a) Imidazolium based ionic liquids in soil: effects of the side chain length on wheat (*Triticum aestivum*) and cress (*Lepidium sativum*) as affected by different clays and organic matter. *Green Chem* 10:584–591. doi:10.1039/b717811e
- Matzke M, Stolte S, Böschen A, Filser J (2008b) Mixture effects and predictability of combination effects of imidazolium based ionic liquids as well as imidazolium based ionic liquids and cadmium on terrestrial plants (*Triticum aestivum*) and limnic green algae (*Scenedesmus vacuolatus*). *Green Chem* 10:784–792. doi:10.1039/b802350f
- Matzke M, Stolte S, Arning J, Uebers U, Filser J (2009) Ionic liquids in soil: effects of different anion species of imidazolium based ionic liquids on wheat (*Triticum aestivum*) as affected by different clay minerals and clay concentrations. *Ecotoxicology* 18:197–203. doi:10.1007/s10646-008-0272-3
- Messali M, Moussa Z, Alzahrani AY, El-Naggar MY, El Douhaibi AS, Judeh ZMA, Hammouti B (2013) Synthesis, characterization and the antimicrobial activity of new eco-friendly ionic liquids. *Chemosphere* 91:1627–1634. doi:10.1016/j.chemosphere.2012.12.062
- Mrozik W, Jungnickel C, Ciborowski T, Pitner WR, Kumirska J, Kaczyński Z, Stepnowski P (2009) Predicting mobility of alkyimidazolium ionic liquids in soils. *J Soils Sediments* 9:237–245. doi:10.1007/s11368-009-0057-1
- Noqueiro RC, Monteiro FA, Gratão PL, Borgo L, Azevedo RA (2015) Tropical soil with high aluminium concentrations cause oxidative stress in two tomato genotypes. *Environ Monit Assess* 187:73. doi:10.1007/s10661-015-4282-3
- OECD (2006) Guidelines for the Testing of Chemicals, Section 2, Test No. 208: Terrestrial Plant Test: Seedling Emergence and Seedling Growth Test. doi:10.1787/20745761
- Oren R, Werk KS, Buchmann N, Zimmermann R (1993) Chlorophyll-nutrient relationships identify nutritionally caused decline in *Picea abies* stands. *Can J For Res* 23:1187–1195
- Oukarroum A, Bussotti F, Goltsev V, Kalaji HM (2015) Correlation between reactive oxygen species production and photochemistry of photosystems I and II in *Lemna gibba* L. plants under salt stress. *Environ Exp Bot* 109:80–88. doi:10.1016/j.envexpbot.2014.08.005

- Pandey PC, Kumar P, Rani M, Katiyar S, Tomar V (2014) Fluoride-induced impact of aluminium industrial power plant on plants and human inhabiting areas. *Geofizika* 31:151–168. doi:10.15233/gfz.2014.31.8
- Pawłowska B, Biczak R (2016) Evaluation of the effect of tetraethylammonium bromide and chloride on the growth and development of terrestrial plants. *Chemosphere* 149:24–33. doi:10.1016/j.chemosphere.2016.01.072
- Peric B, Sierra J, Martí E, Cruañas R, Garau MA, Arning J, Bottin-Weber U, Stolte S (2013) (Eco)toxicity and biodegradability of selected protic and aprotic ionic liquids. *J Hazard Mater* 261:99–105. doi:10.1016/j.jhazmat.2013.06.070
- Petkovic M, Hartmann DO, Adamová G, Seddon KR, Rebelo LPN, Silva Pereira C (2012) Unravelling the mechanism of toxicity of alkyltributylphosphonium chlorides in *Aspergillus nidulans* conidia. *New J Chem* 36:56–63. doi:10.1039/c1nj20470j
- Pham TPT, Cho C-W, Yun Y-S (2010) Environmental fate and toxicity of ionic liquids: a review. *Water Res* 44:352–372. doi:10.1016/j.watres.2009.09.030
- Pham TPT, Cho C-W, Yun Y-S (2016) Structural effects of ionic liquids on microalgal growth inhibition and microbial degradation. *Environ Sci Pollut Res* 23:4249–4300. doi:10.1007/s11356-015-5287-8
- Radošević K, Cvjetko Bubalo M, Srček VG, Grgas D, Dragičević TL, Radojčić Redovniković I (2015) Evaluation of toxicity and biodegradability of choline chloride based deep eutectic solvents. *Ecotoxicol Environ Saf* 112:46–53. doi:10.1016/j.ecoenv.2014.09.034
- Rosalie R, Joas J, Deytieu-Belleau C, Vulcain E, Payet B, Dufossé L, Léchaudel M (2015) Antioxidant and enzymatic responses to oxidative stress induced by pre-harvest water supply reduction and ripening on mango (*Mangifera indica* L. cv. 'Cogshall') in relation to carotenoid content. *J Plant Physiol* 184:68–78. doi:10.1016/j.jplph.2015.05.019
- Sánchez-Rodríguez E, Rubio-Wilhelmi MM, Cervilla LM, Blasco B, Rios JJ, Rosales MA, Romero L, Ruiz JM (2010) Genotypic differences in some physiological parameters symptomatic for oxidative stress under moderate drought in tomato plants. *Plant Sci* 178:30–40. doi:10.1016/j.plantsci.2009.10.001
- Schubert BA, Jahren AH (2011) Fertilization trajectory of the root crop *Raphanus sativus* across atmospheric pCO<sub>2</sub> estimates of the next 300 years. *Agric Ecosyst Environ* 140:174–181. doi:10.1016/j.agee.2010.11.024
- Singh HP, Batish DR, Kohli RK, Arora K (2007) Arsenic-induced root growth inhibition in mung bean (*Phaseolus aureus* Roxb.) is due to oxidative stress resulting from enhanced lipid peroxidation. *Plant Growth Regul* 53:65–73. doi:10.1007/s10725-007-9205-z
- Stolte S, Matzke M, Arning J, Bösch A, Pitner W-R, Welz-Biermann U, Jastorff B, Ranke J (2007) Effects of different head groups and functionalised sidechains on the aquatic toxicity of ionic liquids. *Green Chem* 9:1170–1179. doi:10.1039/b711119c
- Studzińska S, Buszewski B (2009) Study of toxicity of imidazolium ionic liquids to watercress (*Lepidium sativum* L.). *Anal Bioanal Chem* 393:983–990. doi:10.1007/s00216-008-2523-9
- Studzińska S, Kowalkowski T, Buszewski B (2009) Study of ionic liquid cations transport in soil. *J Hazard Mater* 168:1542–1547. doi:10.1016/j.hazmat.2009.03.029
- Telesiński A, Śnioszek M (2009) Bioindicators of environmental pollution with fluorine. *Bromatol Chem Toksykol* 4:1148–1154 In Pol.
- Wang L-S, Wang L, Wang L, Wang G, Li Z-H, Wang J-J (2009) Effect of 1-butyl-3-methylimidazolium tetrafluoroborate on the wheat (*Triticum aestivum* L.) seedlings. *Environ Toxicol* 24:296–303. doi:10.1002/tox.20435
- Zhang B, Li X, Chen D, Wang J (2013) Effects of 1-octyl-3-methylimidazolium bromide on the antioxidant system of *Lemna minor*. *Protoplasma* 250:103–110. doi:10.1007/s00709-012-0379-5