ERRATUM

Erratum to: Selenium nanoparticles with low-level ionizing radiation exposure ameliorate nicotine-induced inflammatory impairment in rat kidney

Walid E. Zahran¹ · Sawsan M. Elsonbaty² · Fatma S. M. Moawed³

Published online: 17 August 2017
© Springer-Verlag GmbH Germany 2017

Erratum to: Environ Sci Pollut Res
<https://doi.org/10.1007/s11356-017-9558-4>

The original publication of this paper contains adjustment errors.

The adjusted images for figures 2 and 4 are now shown.

In table 1, the accession no. for GAPDH was adjusted.

The original article was corrected.

The online version of the original article can be found at doi:[10.1007/s11356-017-9558-4](https://doi.org/10.1007/s11356-017-9558-4)

✉ Walid E. Zahran
wahid_zahran@sci.asu.edu.eg

¹ Biochemistry Department, Faculty of Science, Ain Shams University, Cairo, Egypt

² Radiation Microbiology Department, National Center for Radiation Research and Technology, Atomic Energy Authority, Cairo, Egypt

³ Health Radiation Research Department, National Center for Radiation Research and Technology, Atomic Energy Authority, Cairo, Egypt

Fig. 2 Comparative FT-IR spectrums of ginger *Zingiber officinale* rhizome extract (a) and prepared ginger rhizome SeNPs (b) towards functional groups analysis

Gr.1 Control	Gr.2 SeNPs	Gr.3 Nicotine	Gr.4 γ -R	Gr.5 SeNPs + γ -R	Gr.6 Nicotine + γ -R	Gr.7 Nicotine + SeNPs	Gr.8 Nicotine + SeNPs + γ -R
------------------------	----------------------	-------------------------	----------------------------	------------------------------------	---------------------------------------	---------------------------------	---

Fig. 4 Effect of nicotine upon COX-1 and COX-2 mRNA expression in treated rat groups. Data were expressed as relative quantification compared to control group using quantitative RT-PCR analysis. Each bar represents mean $\pm$ SD ($n = 10$). ^a $P < 0.05$ compared to control; ^b $P < 0.05$ compared to nicotine-treated group; ^c $P < 0.05$ compared to nicotine combined treatments