Prenylated flavonoids and related compounds of the Indonesian *Artocarpus* (Moraceae)

Euis H. Hakim · Sjamsul A. Achmad · Lia D. Juliawaty · Lukman Makmur ·
Yana M. Syah · Norio Aimi · Mariko Kitajima · Hiromitsu Takayama ·
Emilio L. Ghisalberti

Published online: 2 February 2007
© The Japanese Society of Pharmacognosy and Springer 2007

Erratum to: J Nat Med 60:161–184
DOI 10.007/s11418-006-0048-0

There were errors in the original version of this article. The structures in Figs. 3, 11, 17, and 23 were incorrect. The corrected figures are shown below.

Quinonoxanthone in Fig. 3

31

32

Artelastin (31) and Artelastochromene (32) in Fig. 11

49

Artonin O (49) in Fig. 17

Xanthonolide and Cyclopentenoxanthone in Fig. 23

The online version of the original article can be found at <http://dx.doi.org/10.007/s11418-006-0048-0>.

E. H. Hakim (✉) · S. A. Achmad · L. D. Juliawaty ·
L. Makmur · Y. M. Syah
Natural Products Research Group,
Department of Chemistry,
Bandung Institute of Technology,
Jalan Ganeca 10, Bandung 40132, Indonesia
e-mail: euis@chem.itb.ac.id

N. Aimi · M. Kitajima · H. Takayama
Graduate School of Pharmaceutical Sciences,
Chiba University, 1-33 Yayoi-cho, Inage-ku,
Chiba 263-8522, Japan

E. L. Ghisalberti
Chemistry, School of Biomedical and Chemical Sciences,
The University of Western Australia, Crawley
WA 6909, Australia