Erratum to: Xeno-repopulation of *Fah*^{-/-}*Nod/Scid* mice livers by human hepatocytes

SU BaoLiang^{1,2†}, LIU ChangCheng^{1†}, XIANG Dao¹, ZHANG HaiBin³, YUAN SiMing⁴,
WANG MinJun¹, CHEN Fei¹, ZHU HaiYing¹, HE ZhiYing^{1*},
WANG Xin⁵ & HU YiPing^{1*}

¹Department of Cell Biology, Second Military Medical University, Shanghai 200433, China;

²Center for Instrumental Analysis, China Pharmaceutical University, Nanjing 210009, China;

³Eastern Hepatobiliary Surgery Hospital, Second Military Medical University, Shanghai 200433, China;

⁴Department of Plastic Surgery, Nanjing General Hospital of Nanjing Military Command, Nanjing 210002, China;

⁵Institute of Biochemistry and Cell Biology, Shanghai Institute for Biological Sciences,
Chinese Academy of Sciences, Shanghai 200031, China

Erratum to: SCIENCE CHINA Life Sciences, March 2011 Vol.54 No.3: 227–234

doi: 10.1007/s11427-011-4140-7

The authors report the correction for their wrong information about *Fah*^{-/-}*Rag2*^{-/-}*Il2rg*^{-/-} mice in three parts of this paper: Abstract, Introduction and Discussion. We announce that the wrong information should be deleted in the following sections:

(i) In Abstract, sentence of “the high mortality and low breeding rate of this model may hinder its application” should be deleted.

(ii) In Introduction, paragraph of “there are some disadvantages to the *Fah*^{-/-}*Rag2*^{-/-}*Il2rg*^{-/-} mouse model, including high mortality rates during breeding, throughout life and after cell transplantation, which have presented a barrier to its larger scale application” should be deleted.

(iii) In Discussion, the paragraph of “maintenance of *Fah*^{-/-}*Rag2*^{-/-}*Il2rg*^{-/-} mice during colony breeding, growing and cell transplantation is difficult, with high mortality rates seen in our previous experiments. Also, the genotyping of the offspring is overly elaborate. Overall, there are still problems with the large-scale application of *Fah*^{-/-}*Rag2*^{-/-}*Il2rg*^{-/-} mice” should be deleted.

To remove these wrong information is to clear away our misunderstanding of the facts for *Fah*^{-/-}*Rag2*^{-/-}*Il2rg*^{-/-} mice. Generally, *Fah*^{-/-}*Rag2*^{-/-}*Il2rg*^{-/-} mice are good in large scale breeding and have no survival problem after cell transplantation surgery. Our paper is only to report the information of introducing an addition model used for the study on xeno-repopulation of human hepatocytes.

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s11427-011-4140-7>