Phototransduction in *Drosophila*

TIAN Yao, HU Wen, TONG HuaWei & HAN JunHai*

Institute of Life Science, the Key Laboratory of Developmental Genes and Human Disease, Southeast University, Nanjing 210096, China

Received July 8, 2011; accepted September 12, 2011

The *Drosophila* visual transduction is the fastest known G protein-coupled signaling cascade and has been served as a model for understanding the molecular mechanisms of other G protein-coupled signaling cascades. Numbers of components in visual transduction machinery have been identified. Based on the functional characterization of these genes, a model for *Drosophila* phototransduction has been outlined, including rhodopsin activation, phosphoinositide signaling, and the opening of TRP and TRPL channels. Recently, the characterization of mutants, showing slow termination, revealed the physiological significance and the mechanism of rapid termination of light response.

GPCR signaling, visual transduction, rhodopsin, TRP channels, *Drosophila*

Citation: Tian Y, Hu W, Tong H W, *et al.* Phototransduction in *Drosophila*. *Sci China Life Sci*, 2011, 54: 27–34, doi: 10.1007/s11427-012-4272-4

1 Overview and relationship of *Drosophila* to mammalian phototransduction

Drosophila vision is the first sensory system that has been detailed studied using genetic analyses in any animal [1]. Now we know that *Drosophila* vision is the fastest known G protein-coupled signaling cascade, which is activated in less than 20 milliseconds [2]. In *Drosophila*, visual transduction is initiated by the photon-induced isomerization of chromophore in rhodopsin, and subsequently activates the heterotrimeric G protein. The effector of the heterotrimeric GTP-binding protein is phospholipase C β (PLC) [3], which leads to the opening of the TRP and TRPL cation channels (Figure 1) [4–7]. In vertebrate rods and cones, visual transduction is also initiated by the activation of rhodopsins and subsequent interaction with heterotrimeric G proteins (transducin). However, in rods and cones, the downstream effector of transducin is a cyclic guanosine monophosphate (cGMP) phosphodiesterase, which leads to the reduction of cGMP levels, and subsequently closes the cGMP-gated

channels [8]. Thus, the consequences of light stimulation on the channels are opposite, opening the channels in fly photoreceptors and closing the channels in rods and cones.

Although the visual transduction cascades are different between fly photoreceptors and mammal rods/cones, mammalian ipRGCs phototransduction cascades show the similarity with fly visual transduction. ipRGCs are the small subset of intrinsically photosensitive retinal ganglion cells, which function in photoentrainment of circadian rhythm and in light-induced papillary constriction [9]. In the ipRGCs, the cascade is initiated through melanopsin, which shows

Figure 1 Model of the *Drosophila* phototransduction cascade.

*Corresponding author (email: junhaihan@seu.edu.cn)

higher similarities in sequence and biophysical properties to fly rhodopsins than the light receptors in rods and cones [10–16]. Some studies have demonstrated that ipRGC cascade functions through the activation of PLC and the opening of channels, which displays the features of TRP channels [10,17,18].

2 The anatomy of *Drosophila* visual system

Drosophila visual system consists of the retina and the optic lobe known as the lamina, the medulla, the lobula, and the lobula plate (Figure 2A). *Drosophila* compound eye comprises approximately 750 unit eyes, called ‘ommatidia’ (Figure 2B and C). One ommatidium contains 20 cells, eight of which are photoreceptor cells (R cells, R1–R8) (Figure 2C) [19]. The main cells surrounding the photoreceptor cells are secondary pigment cells. Peripheral photoreceptors (R1–6) arrange as a hexagon and extend the retina, while central photoreceptors (R7 and R8) reside in the centre and occupy the distal and proximal regions of the ommatidia, respectively (Figure 2A). R1–R6 expresses major Rhodopsin (Rh1), which is sensitive to a broad spectrum of light with a maximum sensitivity in 486 nm [20,21]. Peripheral photoreceptors function as the rod cells in vertebrate, and mediate motion detection. R7s express about 30% and 70% of the rhodopsin Rh3 and Rh4 respectively, which

are sensitive to ultraviolet light. While R8s express the blue light-sensitive Rh5 and green light-sensitive Rh6 opsin, respectively, with the same relative proportions as R7 photoreceptors in a seemingly random distribution pattern [22–25]. R7 and R8 neurons are thought of the equal function as vertebrate cone cells, and are specialized for color vision.

R-cell neuron axons from a single ommatidium form a bundle to innervate the lamina (R1–R6) and the medulla (R7 and R8) optic ganglia (Figure 2A). R1–R6 axons terminate in the lamina and connect with the lamina neurons to form synaptic units called ‘cartridges’ (Figure 2D). Each cartridge contains R1–R6 input terminal from six separate ommatidia, five lamina neurons (L1–L5), and several other cell types (Figure 2E) [26]. The projection pattern of the R1–R6 axons mirrors the arrangement of their rhabdomeres in the ommatidium. This kind of organization facilitates to pool the visual input from the R1–R6 neurons from six separate ommatidia.

The membrane of each photoreceptor cell forms a stack microvillar structure, called rhabdomere, which is the functional equivalent of the rod and cone outer segment in mammals (Figure 2C). Each R1–6 neuron contains about 50000 microvilli, whereas each R7/8 neuron contains about 17000 microvilli. Each microvilli provides a massive plasma membrane surface to anchor a high concentration of rhodopsin and the proteins that function in phototransduction.

Figure 2 The anatomy of the fly visual system.

(A) Connection patterns of the photoreceptor neurons. Each ommatidium contains eight photoreceptor neurons (R1–R8). R-cell axons from a single ommatidium form a bundle to innervate the lamina (R1–R6) and the medulla (R7 and R8) optic ganglia. Single ommatidium is shown. R, photoreceptor cells 1–6; R7, photoreceptor cell 7; R8, photoreceptor cell 8. (B) EM image of eye cross section through the distal regions of the ommatidia. The eye comprises about 750 ommatidia, each of which contains eight photoreceptor neurons termed R1–R8. (C) Enlarged EM image from B. Seven photoreceptor cells are present. N, nucleus. (D) EM image of cross section through the lamina. R1–R6 axons terminate in the lamina and connect with the lamina neurons to form synaptic units, cartridges. (E) Enlarged EM image from D. Each cartridge contains R1–R6 input terminal from six separate ommatidia, five lamina neurons (L1–L5). Two lamina neurons are labeled L, Lamina neuron.

3 Genetic approaches of *Drosophila* phototransduction

Early genetic screens for the mutations affecting visual response were performed on behavioral assays, such as phototaxis or optomotor response in adult flies [27–29]. Retinal degeneration screens also isolated some mutations that show defective phototransduction [28,30,31]. In the late of 1960s, light-evoked electrophysiological responses of the eye (electroretinogram, ERG) were used in the studies of phototransduction [32,33]. ERGs are extracellular recordings, which measure the summed responses of the compound eye to light [34]. Later, ERG based chemical mutagenesis screens turned highly effective and productive in the isolation of mutants that display defective phototransduction. The ERG assays not only identified all the mutants isolated in behavioral screens but also isolated many additional mutants that were not identified by behavioral screens [35]. From the late 1960s to the late 1970s, more than 200 mutants that display defective ERG response were isolated from approximately 10^5 flies.

For ERG recording, live flies are immobilized and two glass microelectrodes filled with ringer's solution are put separately on the eye surface and the thorax (as reference). High intensity light pulses (4000 lux) are used to stimulate the eye of fly. Once the light turns on, the phototransduction cascade is activated, TRP and TRPL channels are opened, the photoreceptor neurons depolarize in 20 milliseconds. The photoreceptor neurons maintain in the depolarization status during the light stimulation. When the light turns off, the phototransduction cascade is deactivated, TRP and TRPL channels are closed, and the photoreceptor neurons re-polarize in 100 milliseconds [34,36,37]. There are several primary features of the ERG response, the activation speed and amplitude that reflects the activation of phototransduction cascade, the maintenance of depolarization status, the termination speed that results from the deactivation of phototransduction cascade, and the on- and off- transients, which occur due to the feedback of postsynaptic activity to the photoreceptor cells in the lamina. We summarized the mutants display abnormal ERG responses with different features (Table 1).

Although the forward genetic screens identified many genes required for phototransduction (Table 1), mutagenesis was not carried out to saturation. Therefore, using reverse genetic approaches, several genes, encoding components of the phototransduction machinery, have been identified and well studied. For example, combined reverse genetics and western blotting analysis, *Drosophila arrestin* mutants were successfully isolated [38]. Some mutants were isolated in the related studies. One mutant was first isolated for the defects in following of visual cues in courtship. Finally, it was confirmed that this allele is a spontaneous mutant of the *transient receptor potential (trp)* gene, which encodes a

light-activated channel subunit [6].

It is possible that some of the eye-enriched genes might function in the phototransduction pathway, because the identified components of phototransduction machinery are usually highly expressed in the eye. Based on this hypothesis, nearly 100 uncharacterized eye-enriched genes were identified in the DNA microarray analysis [39]. Two eye-enriched genes have been characterized to function in the visual response [39–41].

4 Activation of the phototransduction cascade

Phototransduction cascade is initiated by the activation of rhodopsin, which is comprised of a seven transmembrane protein, opsin, and a chromophore. The chromophore is covalently linked to a lysine residue in the seventh transmembrane domain through a Schiff base. In vertebrates the chromophore is 11-*cis* retinal, while in *Drosophila* it is 11-*cis* 3-hydroxyretinal [42]. Photon absorption leads to the photoisomerization of chromophore, and results in the conformational change of opsin, forming the activated metarhodopsin. In vertebrates, *all-trans* retinal subsequently dissociates from metarhodopsin and re-isomerizes through a lengthy enzymatic pathway. In rods, it takes approximately 30 min for retinal to be re-isomerizes and absorbed by rhodopsin. In contrast, *Drosophila* metarhodopsin is usually thermostable, and retinal does not dissociate from metarhodopsin and can be directly re-isomerized back by absorption of 580 nm light [43].

Metarhodopsin in turn activates heterotrimeric G proteins by shifting α -subunit from an inactive GDP-bound form to active GTP-bound form [43–45]. Activated G protein α -subunit then dissociates from the receptor and opens the cascades. In vertebrates, the downstream effector for the G protein is a phosphodiesterase, which hydrolyzes 3'-5' cyclic guanosine monophosphate (cGMP) to 5' GMP, and leads to the closure of cGMP-gated ion channels to terminate the Na^+ and Ca^{2+} influx [46,47]. In contrast, the effector for the *drosophila* heterotrimeric G protein is PLC, which catalyzes phosphatidylinositol-4,5-bisphosphate (PIP_2) to form inositol-1,3,5-trisphosphate (IP_3) and diacylglycerol (DAG) [3,48–50]. Activation of PLC results in the opening of cation influx channels and leads to Na^+ and Ca^{2+} influx [2,6].

5 Deactivation of the phototransduction cascade

In the first decade, the study of phototransduction focused on the activation of visual signaling by isolating defective mutants and by identifying the components. Recently, people try to reveal the mechanisms of deactivation in visual response, because timely deactivation is required for the receiving of subsequent stimulation and preventing toxicity

induced by excessive Ca^{2+} influx [51].

The most important step in phototransduction termination is the deactivation of metarhodopsin. In this step, Arrestins plays a critical role by displacing Gq α subunit to bind with rhodopsin [38,52,53]. *Drosophila* photoreceptor cells contain two Arrestins, Arrestin1 and Arrestin2 (Arr 1 and Arr 2) [54–56]. Arr2, the major isoform of Arrestins, plays the predominant roles in the deactivation of rhodopsin [38], while Arr1 mediates light-dependent rhodopsin endocytosis [52]. Different from other GPCR signaling pathway, the phosphorylation of fly rhodopsin is not required for its deactivation [57], but is essential for its endocytosis [52]. However, the dephosphorylation of rhodopsin is critical for receptor deactivation [58].

A novel mechanism for rhodopsin deactivation, which involves a CaM binding transcription factor, dCAMTA, and its transcriptional target, *dFbx14*, has been demonstrated [36]. The role of FBXL4 in regulating the photoresponse termination remains to be investigated. It is possible that there are some additional mechanisms that function in rhodopsin deactivation, as photo response can still be terminated in *dcamta;arr2* double mutant flies [36].

G protein is deactivated by promoting the hydrolysis of GTP to GDP. As a signaling switch, PLC also deactivates G

protein through its C-terminal region in stimulating the intrinsic GTPase activity of Gq α [59, 60].

A couple of molecules participate in TRP deactivation. INAC has been shown to play critical roles in deactivating TRP channels [61]. INAC is one member of the PKC family of serine/threonine protein kinases [62,63], which is activated by Ca^{2+} and DAG [64]. Using phosphorylation assays in isolated signaling complexes, TRP was identified to undergo Ca^{2+} -dependent phosphorylation modification [61]. CaM is likely to be another regulator in the deactivation of TRP and TRPL channels. *In vitro* studies showed that both TRP and TRPL bind to CaM through their C-terminals [5,65,66]. *In vivo* studies demonstrated that the CaM binding sites of TRPL are critical for rapid termination and *cam* mutant shows slow termination defects [67]. Additional, Ca^{2+} also plays a critical role in the deactivation of light response [68, 69]. In the whole-cell recordings of photoreceptor cells, termination speed depends on the Ca^{2+} [69–71].

Based on these studies, several negative feedback regulation loops exist in the deactivation of visual response. Ca^{2+} /CaM/INAC/TRP pathway serves as a rapid negative feedback regulation and Ca^{2+} /CaM/dCAMTA/FBXL4 pathway acts as a slow negative feedback regulation. Other deactivation mechanisms need to be further investigated.

Table 1 Transduction mutants and phenotype

Gene	Molecular function	Alleles	Mutagen	ERG phenotype	References
<i>arr1</i>	Minor arrestin, metarhodopsin binding	<i>arr1</i> ¹	Insertion in the 2 nd intron		[72]
		<i>arr1</i> ²	C ¹⁵⁴ →S		[72]
		<i>arr1</i> ³	V ⁵² →D	PDA defect	[58]
		<i>arr2</i> ⁵	Y ²⁰ →stop	Slow termination; ERG amplitude reduced	[73, 74]
<i>arr2</i>	Major arrestin	<i>arr2</i> ³	V ⁵² →D	Slow termination;	[38]
<i>calx</i>	Na ⁺ /Ca ²⁺ exchanger	<i>Calx</i> ^A	T822→I	no PDA, inactivation of light response	[75]
<i>cam</i>	Calmodulin,	<i>cam</i> ³⁵²	Deletion the promoter	Slow termination	[76]
<i>camta</i>	Cam binding transcription activator	<i>camta</i> ^{tes-1}	Q398→stop	Slow termination	[36]
		<i>camta</i> ^{tes-2}	S577→R;L1420→stop	Slow termination	[36]
<i>cdsA</i>	CDP diglyceride synthetase	<i>CdsA</i> ¹	P-element activity	Reduction in light sensitivity;	[77]
<i>Calnexin</i>	rhodopsin chaperone			No PDA	[78]
<i>cry</i>	Cryptochrome	<i>cry</i> ^b	D410→N	very low light sensitivity	[79]
<i>Gα49B</i>	α subunit of G protein	<i>G α 49B</i> ¹	Deletion of 154-156 aa	~1000-fold reduction in sensitivity to light; slow termination; increased activation latency	[50, 80]
		<i>Gαq</i> ¹			
<i>Gβ76C</i>	β subunit of G protein	<i>Gβe</i> ²	G ²⁸⁸ →E	Reduced light sensitivity; slow termination	[81]
		<i>Gβe</i> ¹	C ²⁹³ →Y	Reduced light sensitivity; slow activation; slow termination	[82]
<i>G γ 30A</i>	γ subunit of G protein	<i>Gγ30A</i> ^{C69}	C69→G	Low light sensitivity	[83]
<i>Gprk1</i>	Rhodopsin kinase	<i>Gprk1</i> ^{hs,PL}	Constructive mutant	Small amplitude	[84]
		<i>inaC</i> ¹	An Hsp70 promoter drives expression of Gprk1.		
<i>inaC</i>	Protein kinase C (PKC)	<i>inaC</i> ²	V ²⁰¹ →D	Slow termination; defective PDA	[2]
		<i>inaC</i> ²⁰⁹	W ⁹³ →stop	Slow termination; slowly decaying receptor potential during the light pulse in ERGs;	[85]
		<i>inaC</i> ⁵	W ¹³⁹ →stop	Slow termination	[85]
		<i>InaD</i> ¹	M ⁴⁴² →K	Defective PDA; decay following the peak response in ERG	[86]
<i>inaD</i>	Scaffold protein; 5 PDZ domains	<i>InaD</i> ^{P215}			
		<i>inaD</i> ^{T1}	R ²⁷⁰ →stop	ERGs: responding only at the highest light intensities smaller amplitude;	[87, 88]
		<i>inaD</i> ^{T2}	G ⁶⁰⁵ →E	ERGs: latency, activation and deactivation are all significantly slower; smaller amplitude	[88, 89]

(To be continued on the next page)

(Continued)

Gene	Molecular function	Alleles	Mutagen	ERG phenotype	References
<i>inaF</i>	Putative regulator of TRP	<i>inaF</i> ^{P106x}	Delete the ORF and 3'-UTR and 5'-UTR	reduction in light response and longer response latency in intracellular recordings	[90]
		<i>laza</i> ^{GE}	P-element activity	Small amplitude	[41]
<i>laza</i>	PA phosphatase	<i>laza</i> ¹	Delete <i>laza</i> (-317 to +1381)	Small amplitude; Slow termination	[41]
		<i>laza</i> ²	Delete <i>laza</i> (-309 to +913)	Small amplitude; Slow termination	[41]
		<i>ninaA</i> ¹			
<i>ninaA</i>	Cyclophilin	<i>ninaA</i> ^{P228}	W ²⁰⁸ →stop	At restrictive temperature, defective PDA	[91]
		<i>ninaA</i> ²			
		<i>ninaA</i> ^{P269}	Q ⁸⁷ →stop	Defective PDA	[92]
<i>ninaB</i>	Retinal isomerase	<i>ninaB</i> ¹			
		<i>ninaB</i> ^{P315}	E ²⁸⁰ →K;M ⁴⁷¹ →L;E ⁴⁷⁷ →A	no PDA	[92]
		<i>ninaC</i> ⁵			
<i>ninaC</i>	Protein kinase	<i>ninaC</i> ^{P235}		slow termination; defective off-transient	[93, 94]
		<i>ninaC</i> ^{Δ174}	T-G transversion in the second base of the tenth intron	slow termination; defective off-transient	[95]
<i>ninaD</i>	Scavenger receptor	<i>ninaD</i> ²		No PDA	[92, 96]
		<i>ninaD</i> ^{P246}			
		<i>ninaE</i> ⁷	G ¹²⁸ →R	No PDA; reduction in sensitivity to light; no on-transient;	[97]
		<i>ninaE</i> ^{P332}			
<i>ninaE</i>	Major rhodopsin	<i>ninaE</i> ⁸	T ²⁸³ →M;W ²⁸⁹ →R;C ²⁹⁷ →S	No PDA; very small ERG response; no on/off transient;	[96]
		<i>ninaE</i> ^{P334}			
		<i>ninaE</i> ¹⁷	1.6 kb deletion of the 5' flanking and 5' coding regions	No PDA; very small ERG response; no on/off-transient	[98, 99]
		<i>ninaE</i> ¹¹⁷			
<i>ninaG</i>	Oxidoreductase	<i>ninaG</i> ^{P330}	Q71→stop	No PDA	[100]
		<i>norpA</i> ³⁶	deletion (2710–2737); substitution of 24 amino acid residues, followed by a premature termination codon	No ERG	[101]
		<i>norpA</i> ^{P24}			
		<i>norpA</i> ^{P29}			
<i>norpA</i>	Phospholipase C (PLC)	<i>norpA</i> ⁷		Very small response	[30,102]
		<i>norpA</i> ^{EE5}			
		<i>norpA</i> ³¹	S ⁵⁵¹ →Y	At restrictive temperature, no photoreceptor potential.	[103]
		<i>norpA</i> ^{H52}			
		<i>ort</i> ¹	Delete of intron 2 (110bp) and 459 nucleotides of exons 2 and 3	No PDA, light coincident response is reduced in amplitude, transient components missing.	[97,104]
<i>ort</i>	Histamine-gated Cl ⁻ channel	<i>ort</i> ⁴		Slow termination kinetics after a long pulse; a faster repolarization after a short pulse;	[105]
<i>pinta</i>	Retinoid binding	<i>pinta</i> ¹	F90→stop	No PDA	[96]
<i>pis</i>	CDP-diacylglycerol-inositol 1 3-phosphatidyltransferase	<i>pis</i> ¹	replacement of a part of the Pis gene with w	Inactivation during light response	[106]
		<i>Pld</i> ^{mult.3'}	Lack the promoter, exon 1 and most of exon 2 and contain a premature STOP codon within exon 5.	Low (~1.9%) light sensitivity	[102]
<i>Pld</i>	Phospholipase D (PLD)	<i>Pld</i> ^{mult.5'}	lacks most of exon 9, K1002→STOP	Low (~1.9%) light sensitivity	[102]
<i>Pph13</i>	Homeobox	<i>Pph13</i> ^{hazy}	W ⁵⁸ →stop	Absence of on- and off-transients	[107]
		<i>rdgA</i> ¹	Q ¹¹⁵³ →stop	Only minimal ERGs in extremely newly emerged flies	[108]
<i>rdgA</i>	DAG kinase	<i>rdgA</i> ^{BS12}			
		<i>rdgA</i> ²	G ⁸⁶⁹ →D	ERG of 7 day old flies shows reduced receptor potentials	[30]
		<i>rdgA</i> ^{KO14}			
		<i>rdgB</i> ²		Very small ERG response of 1-day-old flies	[109]
<i>rdgB</i>	PI transfer protein	<i>rdgB</i> ⁹			
		<i>rdgB</i> ^{K5222}	Q ¹⁴⁷ →stop	Small ERG response without on/off-transient;	[30]
<i>rdgC</i>	Rhodopsin phosphatase	<i>rdgC</i> ³⁰⁶		slow termination	[74, 84]
<i>repo</i>	RNA polymerase II transcription factor	<i>repo</i> ¹	P-element activity	Homozygotes exhibit an age-dependent reverse polarity ERG recording.	[110]
			Delete the translation initiation codon and the following two methionine codons	reduced PDA	[111]
<i>rtp</i>	Retinophilin	<i>rtp</i> ¹			
<i>santa-maria</i>	Scavenger receptor	<i>santa-maria</i> ¹	G217→D	No PDA	[112]
<i>stmA</i>	Possible lipase in PIP2 pathway	<i>stmA</i> ¹	G487→D	At 37°C, the ERG is completely eliminated in these flies.	[113]
		<i>rbo</i> ^{ts}			
<i>t</i>	β-Alanyl-histamine hydrolase	<i>t</i> ¹	R ¹²⁷ →P	ERG abnormal	[114]
		<i>t</i> ⁸		ERG abnormal	[115]
		<i>trp</i> ²			
		<i>trp</i> ³⁰¹		Transient potential; reduction in light response	[116, 117]
<i>trp</i>	Ca ²⁺ channel	<i>trp</i> ⁹		Transient potential, reduction in light response and longer response latency	[116, 118]
		<i>trp</i> ³⁴³			
		<i>trp</i> ^{P365}	Pro ⁵⁰⁰ →Thr;His ⁵³¹ →Asn Phe ⁵⁵⁰ →Ile;Ser ⁸⁶⁷ →Phe	Very small ERG response, but not transient	[119]
<i>trpl</i>	Ca ²⁺ channel	<i>trpl</i> ³⁰²	Q ³⁰² →stop	The receptor potential is smaller.	[116]

We apologize to those people, whose work have contributed greatly to our knowledge but were not sufficiently reviewed or were not cited owing to space limitations. We thank Yi Li for figure preparation, and people in the Han laboratory for proof-reading. This work was supported by National Natural Science Foundation of China (Grant Nos. 30970663 and 31070683) and New-Century Training Program Foundation for the Talents by the State Education Commission and Youth Foundation of Southeast University to J. H.

- 1 Pak W L, Grossfield J, White N V. Nonphototactic mutants in a study of vision of *Drosophila*. *Nature*, 1969, 222: 351–354
- 2 Ranganathan R, Harris G L, Stevens C F, et al. A *Drosophila* mutant defective in extracellular calcium-dependent photoreceptor deactivation and rapid desensitization. *Nature*, 1991, 354: 230–232
- 3 Bloomquist B T, Shortridge R D, Schneuwly S, et al. Isolation of a putative phospholipase C gene of *Drosophila*, *norpA*, and its role in phototransduction. *Cell*, 1988, 54: 723–733
- 4 Montell C, Rubin G M. Molecular characterization of the *Drosophila* *trp* locus: a putative integral membrane protein required for phototransduction. *Neuron*, 1989, 2: 1313–1323
- 5 Phillips A M, Bull A, Kelly L E. Identification of a *Drosophila* gene encoding a calmodulin-binding protein with homology to the *trp* phototransduction gene. *Neuron*, 1992, 8: 631–642
- 6 Hardie R C, Minke B. The *trp* gene is essential for a light-activated Ca^{2+} channel in *Drosophila* photoreceptors. *Neuron*, 1992, 8: 643–651
- 7 Niemeyer B A, Suzuki E, Scott K, et al. The *Drosophila* light-activated conductance is composed of the two channels TRP and TRPL. *Cell*, 1996, 85: 651–659
- 8 Fu Y, Yau K W. Phototransduction in mouse rods and cones. *Pflügers Arch*, 2007, 454: 805–819
- 9 Berson D M. Phototransduction in ganglion-cell photoreceptors. *Pflügers Arch*, 2007, 454: 849–855
- 10 Qiu X, Kumbalasiri T, Carlson S M, et al. Induction of photosensitivity by heterologous expression of melanopsin. *Nature*, 2005, 433: 745–749
- 11 Provencio I, Jiang G, De Grip W J, et al. Melanopsin: An opsin in melanophores, brain, and eye. *Proc Natl Acad Sci USA*, 1998, 95: 340–345
- 12 Provencio I, Rodriguez I R, Jiang G, et al. A novel human opsin in the inner retina. *J Neurosci*, 2000, 20: 600–605
- 13 Ruby N F, Brennan T J, Xie X, et al. Role of melanopsin in circadian responses to light. *Science*, 2002, 298: 2211–2213
- 14 Panda S, Sato T K, Castrucci A M, et al. Melanopsin (Opn4) requirement for normal light-induced circadian phase shifting. *Science*, 2002, 298: 2213–2216
- 15 Hattar S, Liao H W, Takao M, et al. Melanopsin-containing retinal ganglion cells: architecture, projections, and intrinsic photosensitivity. *Science*, 2002, 295: 1065–1070
- 16 Hannibal J, Hindersson P, Knudsen S M, et al. The photopigment melanopsin is exclusively present in pituitary adenylate cyclase-activating polypeptide-containing retinal ganglion cells of the retinohypothalamic tract. *J Neurosci*, 2002, 22: RC191.
- 17 Panda S, Nayak S K, Campo B, et al. Illumination of the melanopsin signaling pathway. *Science*, 2005, 307: 600–604
- 18 Isoldi M C, Rollag M D, Castrucci A M, et al. Rhabdomic phototransduction initiated by the vertebrate photopigment melanopsin. *Proc Natl Acad Sci USA*, 2005, 102: 1217–1221
- 19 Ready D F, Hanson T E, Benzer S. Development of the *Drosophila* retina, a neurocrystalline lattice. *Dev Biol*, 1976, 53: 217–240
- 20 O'Tousa J E, Baehr W, Martin R L, et al. The *Drosophila* *ninaE* gene encodes an opsin. *Cell*, 1985, 40: 839–850
- 21 Zuker C S, Cowman A F, Rubin G M. Isolation and structure of a rhodopsin gene from *D. melanogaster*. *Cell*, 1985, 40: 851–858
- 22 Fryxell K J, Meyerowitz E M. An opsin gene that is expressed only in the R7 photoreceptor cell of *Drosophila*. *EMBO J*, 1987, 6: 443–451
- 23 Zuker C S, Montell C, Jones K, et al. A rhodopsin gene expressed in photoreceptor cell R7 of the *Drosophila* eye: homologies with other signal-transducing molecules. *J Neurosci*, 1987, 7: 1550–1557
- 24 Montell C, Jones K, Zuker C, et al. A second opsin gene expressed in the ultraviolet-sensitive R7 photoreceptor cells of *Drosophila melanogaster*. *J Neurosci*, 1987, 7: 1558–1566
- 25 Fortini M E, Rubin G M. Analysis of cis-acting requirements of the Rh3 and Rh4 genes reveals a bipartite organization to rhodopsin promoters in *Drosophila melanogaster*. *Genes Dev*, 1990, 4: 444–463
- 26 Ting C Y, Lee C H. Visual circuit development in *Drosophila*. *Curr Opin Neurobiol*, 2007, 17: 65–72
- 27 Choe K M, Clandinin T R. Thinking about visual behavior; learning about photoreceptor function. *Curr Top Dev Biol*, 2005, 69: 187–213
- 28 Hotta Y, Benzer S. Genetic dissection of the *Drosophila* nervous system by means of mosaics. *Proc Natl Acad Sci USA*, 1970, 67: 1156–1163
- 29 Pak W L, Grossfield J, Arnold K S. Mutants of the visual pathway of *Drosophila melanogaster*. *Nature*, 1970, 227: 518–520
- 30 Harris W A, Stark W S. Hereditary retinal degeneration in *Drosophila melanogaster*. A mutant defect associated with the phototransduction process. *J Gen Physiol*, 1977, 69: 261–291
- 31 Steele F, O'Tousa J E. Rhodopsin activation causes retinal degeneration in *Drosophila* *rdgC* mutant. *Neuron*, 1990, 4: 883–890
- 32 Hotta Y, Benzer S. Abnormal electroretinograms in visual mutants of *Drosophila*. *Nature*, 1969, 222: 354–356
- 33 Cosens D J, Manning A. Abnormal electroretinogram from a *Drosophila* mutant. *Nature*, 1969, 224: 285–287
- 34 Hengstenberg R, Gotz K G. Effect of facet-separating pigments on the perception of light and contrast in eye mutants of *Drosophila*. *Kybernetik*, 1967, 3: 276–285
- 35 Pak W L, Leung H T. Genetic approaches to visual transduction in *Drosophila melanogaster*. *Recept Chann*, 2003, 9: 149–167
- 36 Han J, Gong P, Reddig K, Mitra M, Guo P, Li H S, et al. The fly CAMTA transcription factor potentiates deactivation of rhodopsin, a G protein-coupled light receptor. *Cell*, 2006, 127: 847–858
- 37 Dolph P, Nair A, Raghu P. Electroretinogram recordings of *Drosophila*. *Cold Spring Harb Protoc*, 2011: pdb.prot5549. doi: 10.1101/pdb.prot5549..
- 38 Dolph P J, Ranganathan R, Colley N J, et al. Arrestin function in inactivation of G protein-coupled receptor rhodopsin *in vivo*. *Science*, 1993, 260: 1910–1916
- 39 Xu H, Lee S J, Suzuki E, et al. A lysosomal tetraspanin associated with retinal degeneration identified via a genome-wide screen. *Embo J*, 2004, 23: 811–822
- 40 Garcia-Murillas I, Pettitt T, Macdonald E, et al. *lazar0* encodes a lipid phosphate phosphohydrolase that regulates phosphatidylinositol turnover during *Drosophila* phototransduction. *Neuron*, 2006, 49: 533–546
- 41 Kwon Y, Montell C. Dependence on the Lazaro phosphatidic acid phosphatase for the maximum light response. *Curr Biol*, 2006, 16: 723–729
- 42 Goldsmith T H, Marks B C, Bernard G D. Separation and identification of geometric isomers of 3-hydroxyretinoids and occurrence in the eyes of insects. *Vision Res*, 1986, 26: 1763–1769
- 43 Kiselev A, Subramaniam S. Activation and regeneration of rhodopsin in the insect visual cycle. *Science*, 1994, 266: 1369–1373
- 44 Emeis D, Kühn H, Reichert J, et al. Complex formation between metarhodopsin II and GTP-binding protein in bovine photoreceptor membranes leads to a shift of the photoproduct equilibrium. *FEBS Lett*, 1982, 143: 29–34
- 45 Kibelbek J, Mitchell D C, Beach J M, et al. Functional equivalence of metarhodopsin II and the Gt-activating form of photolyzed bovine rhodopsin. *Biochemistry*, 1991, 30: 6761–6768
- 46 Palczewski K, Saari J C. Activation and inactivation steps in the visual transduction pathway. *Curr Opin Neurobiol*, 1997, 7: 500–504
- 47 Yau K W, Baylor D A. Cyclic GMP-activated conductance of retinal photoreceptor cells. *Annu Rev Neurosci*, 1989, 12: 289–327
- 48 Devary O, Heichal O, Blumenfeld A, et al. Coupling of photoexcited rhodopsin to inositol phospholipid hydrolysis in fly photoreceptors.

- Proc Natl Acad Sci USA, 1987, 84: 6939–6943
- 49 Lee Y J, Shah S, Suzuki E, et al. The *Drosophila* *dgg* gene encodes a G alpha protein that mediates phototransduction. *Neuron*, 1994, 13: 1143–1157
 - 50 Scott K, Becker A, Sun Y, et al. Gq alpha protein function *in vivo*: genetic dissection of its role in photoreceptor cell physiology. *Neuron*, 1995, 15: 919–927
 - 51 Han J, Reddig K, Li H S. Prolonged G(q) activity triggers fly rhodopsin endocytosis and degradation, and reduces photoreceptor sensitivity. *Embo J*, 2007, 26: 4966–4973
 - 52 Satoh A K, Ready D F. Arrestin1 mediates light-dependent rhodopsin endocytosis and cell survival. *Curr Biol*, 2005, 15: 1722–1733
 - 53 Ranganathan R, Stevens C F. Arrestin binding determines the rate of inactivation of the G protein-coupled receptor rhodopsin *in vivo*. *Cell*, 1995, 81: 841–848
 - 54 Toyoshima S, Matsumoto N, Wang P, et al. Purification and partial amino acid sequences of phosphoinositide-specific phospholipase C of *Drosophila* eye. *J Biol Chem*, 1990, 265: 14842–14848
 - 55 Smith D P, Shieh B H, Zuker C S. Isolation and structure of an arrestin gene from *Drosophila*. *Proc Natl Acad Sci USA*, 1990, 87: 1003–1007
 - 56 Hyde D R, Mecklenburg K L, Pollock J A, et al. Twenty *Drosophila* visual system cDNA clones: one is a homolog of human arrestin. *Proc Natl Acad Sci USA*, 1990, 87: 1008–1012
 - 57 Scott K, Zuker C. Lights out: deactivation of the phototransduction cascade. *Trends Biochem Sci*, 1997, 22: 350–354
 - 58 Vinós J, Jalink K, Hardy R W, et al. A G protein-coupled receptor phosphatase required for rhodopsin function. *Science*, 1997, 277: 687–690
 - 59 Montell C. PLC fills a GAP in G-protein-coupled signalling. *Nat Cell Biol*, 2000, 2: E82–83
 - 60 Cook B, Bar-Yaacov M, Cohen Ben-Ami H, et al. Phospholipase C and termination of G-protein-mediated signalling *in vivo*. *Nat Cell Biol*, 2000, 2: 296–301
 - 61 Huber A, Sander P, Böhner M, et al. The TRP Ca^{2+} channel assembled in a signaling complex by the PDZ domain protein INAD is phosphorylated through the interaction with protein kinase C (ePKC). *FEBS Lett*, 1998, 425: 317–322
 - 62 Smith D P, Ranganathan R, Hardy R W, et al. Photoreceptor deactivation and retinal degeneration mediated by a photoreceptor-specific protein kinase C. *Science*, 1991, 254: 1478–1484
 - 63 Schaeffer E, Smith D, Mardon G, et al. Isolation and characterization of two new *Drosophila* protein kinase C genes, including one specifically expressed in photoreceptor cells. *Cell*, 1989, 57: 403–412
 - 64 Huber A, Sander P, Paulsen R. Phosphorylation of the *InaD* gene product, a photoreceptor membrane protein required for recovery of visual excitation. *J Biol Chem*, 1996, 271: 11710–11717
 - 65 Chevesich J, Kreuz A J, Montell C. Requirement for the PDZ domain protein, INAD, for localization of the trp store-operated channel to a signaling complex. *Neuron*, 1997, 18: 95–105
 - 66 Warr C G, Kelly L E. Identification and characterization of two distinct calmodulin-binding sites in the Trpl ion-channel protein of *Drosophila melanogaster*. *Biochem J*, 1996, 314 (Pt 2): 497–503
 - 67 Scott K, Sun Y, Beckingham K, et al. Calmodulin regulation of *Drosophila* light-activated channels and receptor function mediates termination of the light response *in vivo*. *Cell*, 1997, 91: 375–383
 - 68 Ranganathan R, Harris G L, Stevens C F, et al. A *Drosophila* mutant defective in extracellular calcium-dependent photoreceptor deactivation and rapid desensitization. *Nature*, 1991, 354: 230–232
 - 69 Hardie R C, Minke B. Calcium-dependent inactivation of light-sensitive channels in *Drosophila* photoreceptors. *J Gen Physiol*, 1994, 103: 409–427
 - 70 Hardie R C. whole-cell recordings of the light induced current in dissociated *Drosophila* photoreceptors: Evidence for feedback by calcium permeating the light-sensitive channels. *Proc Royal Soc London. Ser B: Biol Sci*, 1991, 245: 203–210
 - 71 Wang T, Xu H, Oberwinkler J, et al. Light activation, adaptation, and cell survival functions of the Na^{+}/Ca^{2+} exchanger CalX. *Neuron*, 2005, 45: 367–378
 - 72 Merrill, C.E., et al., Visual arrestins in olfactory pathways of *Drosophila* and the malaria vector mosquito *Anopheles gambiae*. *Proc Natl Acad Sci USA*, 2002, 99: 1633–1638
 - 73 Lee S J, Xu H, Kang L W, et al. Light adaptation through phosphoinositide-regulated translocation of *Drosophila* visual arrestin. *Neuron*, 2003, 39: 121–132
 - 74 Lee S J, Montell C. Suppression of constant-light-induced blindness but not retinal degeneration by inhibition of the rhodopsin degradation pathway. *Curr Biol*, 2004, 14: 2076–2085
 - 75 Wang T, Xu H, Oberwinkler J, et al. Light activation, adaptation, and cell survival functions of the Na^{+}/Ca^{2+} exchanger CalX. *Neuron*, 2005, 45(3): p. 367–78
 - 76 Scott K, Sun Y, Beckingham K, et al. Calmodulin regulation of *Drosophila* light-activated channels and receptor function mediates termination of the light response *in vivo*. *Cell*, 1997, 91: 375–383
 - 77 Wu L, Niemeyer B, Colley N, et al. Regulation of PLC-mediated signalling *in vivo* by CDP-diacylglycerol synthase. *Nature*, 1995, 373: 216–222
 - 78 Rosenbaum E E, Hardie R C, Colley N J. Calnexin is essential for rhodopsin maturation, Ca^{2+} regulation, and photoreceptor cell survival. *Neuron*, 2006, 49: 229–241
 - 79 Yuan Q, Lin F, Zheng X, et al. Serotonin modulates circadian entrainment in *Drosophila*. *Neuron*, 2005, 47: 115–127
 - 80 Hardie R C, Martin F, Cochrane G W, et al. Molecular basis of amplification in *Drosophila* phototransduction: roles for G protein, phospholipase C, and diacylglycerol kinase. *Neuron*, 2002, 36: 689–701
 - 81 Dolph P J, Man-Son-Hing H, Yarfitz S, et al. An eye-specific G beta subunit essential for termination of the phototransduction cascade. *Nature*, 1994, 370: 59–61
 - 82 Elia N, Frechter S, Gedi Y, et al. Excess of Gbetas over Gqalphae *in vivo* prevents dark, spontaneous activity of *Drosophila* photoreceptors. *J Cell Biol*, 2005, 171:517–526
 - 83 Schillo S, Belusic G, Hartmann K, et al. Targeted mutagenesis of the farnesylation site of *Drosophila* *Ggammae* disrupts membrane association of the G protein betagamma complex and affects the light sensitivity of the visual system. *J Biol Chem*, 2004, 279: 36309–36316
 - 84 Lee S J, Xu H, Montell C. Rhodopsin kinase activity modulates the amplitude of the visual response in *Drosophila*. *Proc Natl Acad Sci USA*, 2004, 101: 11874–11879
 - 85 Smith D P, Ranganathan R, Hardy R W, et al. Photoreceptor deactivation and retinal degeneration mediated by a photoreceptor-specific protein kinase C. *Science*, 1991, 254: 1478–1484
 - 86 Shieh B H, Niemeyer B. A novel protein encoded by the *InaD* gene regulates recovery of visual transduction in *Drosophila*. *Neuron*, 1995, 14: 201–210
 - 87 Scott K, Zuker C S. Assembly of the *Drosophila* phototransduction cascade into a signalling complex shapes elementary responses. *Nature*, 1998, 395: 805–808
 - 88 Tsunoda S, Sierralta J, Sun Y, et al. A multivalent PDZ-domain protein assembles signalling complexes in a G-protein-coupled cascade. *Nature*, 1997, 388: 243–249
 - 89 Mishra P, Socolich M, Wall M A, et al. Dynamic scaffolding in a G protein-coupled signaling system. *Cell*, 2007, 131: 80–92
 - 90 Li C, Geng C, Leung H T, et al. INAF, a protein required for transient receptor potential Ca(2+) channel function. *Proc Natl Acad Sci USA*, 1999, 96: 13474–13479
 - 91 Stamnes M A, Shieh B H, Chuman L, et al. The cyclophilin homolog ninaA is a tissue-specific integral membrane protein required for the proper synthesis of a subset of *Drosophila* rhodopsins. *Cell*, 1991, 65: 219–227
 - 92 Gu G, Yang J, Mitchell K A, et al. *Drosophila* ninaB and ninaD act outside of retina to produce rhodopsin chromophore. *J Biol Chem*, 2004, 279: 18608–18613
 - 93 Wes P D, Xu X Z, Li H S, et al. Termination of phototransduction requires binding of the NINAC myosin III and the PDZ protein INAD. *Nat Neurosci*, 1999, 2: 447–453
 - 94 Porter J A, Montell C. Distinct roles of the *Drosophila* ninaC kinase

- and myosin domains revealed by systematic mutagenesis. *J Cell Biol*, 1993, 122: 601–612
- 95 Hofstee C A, Henderson S, Hardie R C, et al. Differential effects of ninaC proteins (p132 and p174) on light-activated currents and pupil mechanism in *Drosophila* photoreceptors. *Vis Neurosci*, 1996, 13: 897–906
 - 96 Wang T, Montell C. Rhodopsin formation in *Drosophila* is dependent on the PINTA retinoid-binding protein. *J Neurosci*, 2005, 25: 5187–5194
 - 97 Scavarda N J, O'Tousa J, Pak W L. *Drosophila* locus with gene-dosage effects on rhodopsin. *Proc Natl Acad Sci USA*, 1983, 80: 4441–4445
 - 98 Iakhine R, Chorna-Ornan I, Zars T, et al. Novel dominant rhodopsin mutation triggers two mechanisms of retinal degeneration and photoreceptor desensitization. *J Neurosci*, 2004, 24: 2516–2526
 - 99 Bentreop J, Schwab K, Pak W L, et al. Site-directed mutagenesis of highly conserved amino acids in the first cytoplasmic loop of *Drosophila* Rh1 opsin blocks rhodopsin synthesis in the nascent state. *EMBO J*, 1997, 16: 1600–1609
 - 100 Sarfare S, Ahmad S T, Joyce M V, et al. The *Drosophila* ninaG oxidoreductase acts in visual pigment chromophore production. *J Biol Chem*, 2005, 280: 11895–11901
 - 101 Kim S, Chen D M, Zavarella K, et al. Substitution of a non-retinal phospholipase C in *Drosophila* phototransduction. *Insect Mol Biol*, 2003, 12: 147–153
 - 102 LaLonde M M, Janssens H, Rosenbaum E, et al. Regulation of phototransduction responsiveness and retinal degeneration by a phospholipase D-generated signaling lipid. *J Cell Biol*, 2005, 169: 471–479
 - 103 Matsumoto H, Kurien B T, Takagi Y, et al. Phosrestin I undergoes the earliest light-induced phosphorylation by a calcium/calmodulin-dependent protein kinase in *Drosophila* photoreceptors. *Neuron*, 1994, 12: 997–1010
 - 104 O'Tousa J E, Leonard D S, Pak W L. Morphological defects in oraJK84 photoreceptors caused by mutation in R1-6 opsin gene of *Drosophila*. *J Neurogenet*, 1989, 6: 41–52
 - 105 Rajaram S, Scott R L, Nash H A. Retrograde signaling from the brain to the retina modulates the termination of the light response in *Drosophila*. *Proc Natl Acad Sci USA*, 2005, 102: 17840–17845
 - 106 Wang T, Montell C. A phosphoinositide synthase required for a sustained light response. *J Neurosci*, 2006, 26: 12816–12825
 - 107 Zelhof A C, Koundakjian E, Scully A L, et al. Mutation of the photoreceptor specific homeodomain gene Pph13 results in defects in phototransduction and rhabdome morphogenesis. *Development*, 2003, 130: 4383–4392
 - 108 Johnson M A, Frayer K L, Stark W S. Characteristics of RDGA: Mutants with retinal degeneration in *Drosophila*. *J Insect Physiol*, 1982, 28: 233–242
 - 109 Milligan S C, Alb J G Jr, Elagina R B, et al. The phosphatidylinositol transfer protein domain of *Drosophila* retinal degeneration B protein is essential for photoreceptor cell survival and recovery from light stimulation. *J Cell Biol*, 1997, 139: 351–363
 - 110 Xiong W C, Okano H, Patel N H, et al. *repo* encodes a glial-specific homeo domain protein required in the *Drosophila* nervous system. *Genes Dev*, 1994, 8: 981–994
 - 111 Venkatachalam K, Wasserman D, Wang X, et al. Dependence on a retinophilin/myosin complex for stability of PKC and INAD and termination of phototransduction. *J Neurosci*, 2010, 30: 11337–11345
 - 112 Wang T, Jiao Y, Montell C. Dissection of the pathway required for generation of vitamin A and for *Drosophila* phototransduction. *J Cell Biol*, 2007, 177: 305–316
 - 113 Huang F D, Matthies H J, Speese S D, et al. Rolling blackout, a newly identified PIP2-DAG pathway lipase required for *Drosophila* phototransduction. *Nat Neurosci*, 2004, 7: 1070–1078
 - 114 Benzer S. Behavioral mutants of *Drosophila* isolated by countercurrent distribution. *Proc Natl Acad Sci USA*, 1967, 58: 1112–1119
 - 115 Lindsley D L, Zimm G G. The Genome of *Drosophila melanogaster*. viii ed. 1992. 1133
 - 116 Leung H T, Geng C, Pak W L. Phenotypes of trpl mutants and interactions between the transient receptor potential (TRP) and TRP-like channels in *Drosophila*. *J Neurosci*, 2000, 20: 6797–6803
 - 117 Popescu D C, Ham A J, Shieh B H. Scaffolding protein INAD regulates deactivation of vision by promoting phosphorylation of transient receptor potential by eye protein kinase C in *Drosophila*. *J Neurosci*, 2006, 26: 8570–8577
 - 118 Leung H T, Tseng-Crank J, Kim E, et al. DAG lipase activity is necessary for TRP channel regulation in *Drosophila* photoreceptors. *Neuron*, 2008, 58: 884–896
 - 119 Yoon J, Ben-Ami H C, Hong Y S, et al. Novel mechanism of massive photoreceptor degeneration caused by mutations in the *trp* gene of *Drosophila*. *J Neurosci*, 2000, 20: 649–659

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.