Adaptive potential of human biceps femoris muscle demonstrated by histochemical, immunohistochemical and mechanomyographical methods

Raja Dahmane · Srdjan Djordjevič · Vika Smerdu

Published online: 23 January 2007
© International Federation for Medical and Biological Engineering 2007

Erratum to: Med Bio Eng Comput 44(11):999–1006 DOI 10.1007/s11517-006-0114-5

Unfortunately, an error occurred in the Acknowledgement and also in Fig. 2. The correct version of the first sentence of Acknowledgement is given below.

This study was supported by grants from the Slovenian Olympic Committee and the TMG-BMC d.d.o.

The correct Fig. 2 is given below:

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s11517-006-0114-5>

R. Dahmane (✉) · V. Smerdu
Institute of Anatomy, Medical Faculty,
University of Ljubljana, Korytkova 2, Ljubljana, Slovenia
e-mail: raja.dahmane@guest.arnes.si

R. Dahmane
University College for Health Studies,
University of Ljubljana, Ljubljana, Slovenia

S. Djordjevič
Institute of Kinesiology, University of Primorska,
Koper, Slovenia

S. Djordjevič
Department of Biology, University of Ljubljana,
Ljubljana, Slovenia

Fig. 2 Fibre types (1, 2a, 2x and 2c) in human biceps femoris muscle demonstrated according to the reaction for mATPase at different preincubations (**a** 4.3, **b** 4.6, **c** 9.4) and immunohistochemical staining with monoclonal antibodies specific to MyHC isoforms: **d** BA-D5 (MyHC-1), **e** SC-71 (MyHC-2a), **f** BF-35 (specific to MyHC-2a if the sections are pre-treated in microwave oven. Bar is 100 μ