Erratum to: Kurtosis and skewness assessments of solid lung nodule density histograms: differentiating malignant from benign nodules on CT

Ayano Kamiya · Sadayuki Murayama ·
Hisashi Kamiya · Tsuneo Yamashiro ·
Yasuji Oshiro · Nobuyuki Tanaka

Published online: 28 February 2014
© Japan Radiological Society 2014

Erratum to: Jpn J Radiol (2014) 32(1):14–21
DOI 10.1007/s11604-013-0264-y

Unfortunately, errors appeared in Table 2 of the above-mentioned article. In the column headed “Kurtosis, Mean $\pm$ SD”, in the entries for “All nodules” the values for “Malignant ($n = 72$)” and “Benign ($n = 21$)” are reversed. They should be “ 1.10 ± 1.77 ” for malignant nodules, and “ 0.42 ± 1.75 ” for benign nodules, respectively.

The online version of the original article can be found under
doi:[10.1007/s11604-013-0264-y](https://doi.org/10.1007/s11604-013-0264-y).

A. Kamiya · S. Murayama · H. Kamiya · T. Yamashiro (✉)
Department of Radiology, Graduate School of Medical Science,
University of the Ryukyus, 207 Uehara, Nishihara,
Okinawa 903-0215, Japan
e-mail: clatsune@yahoo.co.jp

A. Kamiya
e-mail: ayano1242004@yahoo.co.jp

S. Murayama
e-mail: sadayuki@med.u-ryukyu.ac.jp

H. Kamiya
e-mail: e954123@mail.goo.ne.jp

Y. Oshiro
Department of Radiology, Okinawa National Hospital, National
Hospital Organization, 3-20-14 Ganeko, Ginowan,
Okinawa 901-2214, Japan
e-mail: oshiro4211@yahoo.co.jp

N. Tanaka
Department of Radiology, Yamaguchi University Graduate
School of Medicine, 1-1-1 Minamikogushi, Ube,
Yamaguchi 755-8505, Japan
e-mail: nbtanaka@yamaguchi-u.ac.jp