ERRATUM

Erratum to: Impact Total Psoas Volume on Short- and Long-Term Outcomes in Patients Undergoing Curative Resection for Pancreatic Adenocarcinoma: a New Tool to Assess Sarcopenia

Neda Amini¹ · Gaya Spolverato¹ · Rohan Gupta¹ · Georgios A. Margonis¹ ·
Yuhree Kim¹ · Doris Wagner¹ · Neda Rezaee¹ · Matthew J. Weiss¹ ·
Christopher L. Wolfgang¹ · Martin M. Makary¹ · Ihab R. Kamel² ·
Timothy M. Pawlik¹

Received: 29 February 2016 / Accepted: 4 March 2016 / Published online: 16 March 2016
© 2016 The Society for Surgery of the Alimentary Tract

Erratum to: J Gastrointest Surg
DOI 10.1007/s11605-015-2835-y

There is an error in Table 2. The correct table is below.

Table 2 Comparison of hospital stay and mortality of sarcopenia vs. non-sarcopenia patients

	According to TPA measurement			According to TPV measurement		
	Sarcopenia (n = 192)	No sarcopenia (n = 571)	P value	Sarcopenia (n = 152)	No sarcopenia (n = 611)	P value
Length of hospital stay, day, median (IQR)	9 (7–15)	8 (7–13)	0.05	10 (7–15.5)	8 (7–13)	0.002
Any complication, n (%)	95 (49.5)	274 (48.0)	0.72	91 (59.9)	278 (45.5)	0.002
Major complication, n (%)	38 (19.8)	88 (15.4)	0.16	34 (22.4)	92 (15.1)	0.03
Postoperative Mortality						
30 days	1 (0.5)	3 (0.5)	1.0	1 (0.7)	3 (0.5)	1.0
90 days	7 (3.7)	10 (1.8)	0.12	6 (4.0)	11 (1.8)	0.11
1 year	56 (29.2)	85 (14.9)	<0.001	45 (29.6)	96 (15.7)	<0.001
3 year	121 (63.0)	187 (32.8)	<0.001	82 (54.0)	226 (37.0)	<0.001
Survival (month), median	18.0	28.4	<0.001	17.0	26.7	<0.001

The online version of the original article can be found at <http://dx.doi.org/10.1007/s11605-015-2835-y>.

✉ Timothy M. Pawlik
tpawlik1@jhmi.edu

¹ Department of Surgery, The Johns Hopkins University School of Medicine, 600 N. Wolfe Street, Blalock 688, Baltimore, MD 21287, USA

² Department of Radiology, The Johns Hopkins University School of Medicine, Baltimore, MD, USA