4. Bearpark H, Elliot L, Grunstein R et al (1995) Snoring and sleep apnea: a population study in Australian men. *Am J Respir Crit Care Med* 15:1459–1465
5. Bliwise DL, Nekich JC, Dement WC (1991) Relative validity of sleep-reported snoring as a symptom of sleep apnea in a sleep clinic population. *Chest* 99:600–608
6. Duran J, Esnaola S, Rubio R, Iztueta A (2001) Obstructive sleep apnea-hypopnea and related clinical features in population-based sample of subjects aged 30–70 yr. *Am J Respir Crit Care Med* 163:685–689
7. Gami AS, Pressman G, Caples SM et al (2004) Association of atrial fibrillation and obstructive sleep apnea. *Circulation* 110:364–367
8. Gus M, Goncalves SC, Martinez D et al (2008) Risk of obstructive sleep apnea by Berlin Questionnaire, but not daytime sleepiness, is associated with resistant hypertension: a case-control study. *Am J Hypertens* 21:832–835
9. Harris M, Glozier N, Ratnavadivel R, Grunstein RR (2009) Obstructive sleep apnea and depression. *Sleep Med Rev* 13:437–444
10. Hiestand DM, Britz P, Goldman M, Phillips B (2006) Prevalence of symptoms and risk of sleep apnea in the US population. Results from the national sleep foundation sleep in america 2005 poll. *Chest* 130:780–786
11. Ip MS, Lam B, Launder IJ et al (2001) A community study of sleep-disordered breathing in middle-aged Chinese men in Hong Kong. *Chest* 119:62–69
12. Ip MS, Lam B, Tang LC et al (2004) A community study of sleep-disordered breathing in middle-aged Chinese women in Hong Kong: prevalence and gender differences. *Chest* 125:127–134
13. Johns MW (1991) A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep* 14(6):540–545
14. Khassawneh B, Ghazzawi M, Khader Y et al (2009) Symptoms and risk of obstructive sleep apnea in primary care patients in Jordan. *Sleep Breath* 13(3):227–232
15. Kim JK, In KH, Kim JH et al (2004) Prevalence of sleep-disordered breathing in middle-aged Korean men and women. *Am J Respir Crit Care Med* 170:1108–1113
16. National Statistical Service of the Republic of Armenia (n. d.) The results of census of Republic of Armenia of 2001 (Figures of the Republic of Armenia). Available at: <http://docs.armstat.am/census/eng.php>. Accessed 2004
17. Netzer NC, Hoegel JJ, Loube D et al (2003) Prevalence of symptoms and risk of sleep apnea in primary care. *Chest* 124:1406–1414
18. Netzer NC, Stooth RA, Netzer CM et al (1999) Using the Berlin Questionnaire to identify patients at risk for the sleep apnea syndrome. *Ann Intern Med* 131:485–491
19. (n. d.) Obesity: preventing and managing the global epidemic. Report of a WHO consultation on obesity. June 3–5, 1997. WHO/NUT/NCD/98.1, i-xv, 1–276, 1998. Geneva, Switzerland: World Health Organization. Available at: <http://diseases.who.int/uhtbin/cgiisirs/Wed+Nov+21+23:21:20+MET+2001/0/49>. Accessed 18 April 2002
20. Punjabi N, Shahar E, Redline S et al (2004) Sleep-disordered breathing, glucose intolerance, and insulin resistance. The Sleep Heart Health Study. *Am J Epidemiol* 160:521–530
21. Reddy EV, Kadiravan T, Mishra HK et al (2009) Prevalence and risk factors of obstructive sleep apnea among middle-aged urban Indians: a community-based study. *Sleep Med* 10:913–918
22. Shahar E, Whitney CW, Redline S et al (2001) Sleep-disordered breathing and cardiovascular disease. Cross-sectional results of the Sleep Heart Health Study. *Am J Respir Crit Care Med* 163:910–916
23. Sforza E, Chouchou F, Pichot V et al (2011) Is the Berlin questionnaire a useful tool to diagnose obstructive sleep apnea in the elderly? *Sleep Med* 12:142–146
24. Sharma SK, Vasudev C, Sinha S et al (2006) Validation of the modified Berlin questionnaire to identify patients at risk for the obstructive sleep apnoea syndrome. *Indian J Med Res* 124:281–290
25. Tregear S, Reston J, Schoelles K, Phillips B (2009) Obstructive sleep apnea and risk of motor vehicle crash: systemic review and meta-analysis. *J Clin Sleep Med* 5(6): 573–581
26. Tufik S, Santos-Silva R, Bittencourt LRA (2010) Obstructive sleep apnea syndrome in the Sao Paulo epidemiologic sleep study. *Sleep Med* 11:441–446
27. Young T, Palta M, Dempsey J et al (1993) The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med* 328:1230–1235
28. Young T, Shahar E, Nieto FJ et al (2002) Predictors of sleep-disordered breathing in community-dwelling adults. The Sleep Heart Health Study. *Arch Intern Med* 162:893–900

Somnologie 2012 · 16:235
DOI 10.1007/s11818-012-0585-6
© Springer-Verlag 2012

M. Karimi · J. Hedner · L. Grote

Center for Sleep and Vigilance Disorders,
Institute of Medicine, Sahlgrenska Academy,
University of Gothenburg, Gothenburg

Erratum to: Is a polysomnographic recording prior to MSLT worth the effort?

Somnologie (2011)

15:239–242

<http://dx.doi.org/10.1007/s0011818-011-0534-9>

In the original publication, the article was wrongly categorized as review article. It is an original article.

Corresponding address

M. Karimi

Center for Sleep and Vigilance Disorders,
Institute of Medicine, Sahlgrenska Academy,
University of Gothenburg
Medicinaregatan 8B, 405 30 Gothenburg
Sweden
mahssa.karimi@lungall.gu.se