Traces of pseudo-differential operators on compact and Hausdorff groups

Shahla Molahajloo · Mohammad Pirhayati

Received: 2 March 2013 / Revised: 27 May 2013 / Accepted: 4 June 2013 /

Published online: 25 June 2013

© The Author(s) 2013. This article is published with open access at Springerlink.com

Abstract We give a characterization of and a trace formula for trace class pseudo-differential operators on compact Hausdorff groups.

Keywords Pseudo-differential operators · Compact groups

Mathematics Subject Classification (2000) Primary 22A10 · 43A77 · 47G30;
Secondary 22C05

1 Introduction

Pseudo-differential operators, first developed by Kohn and Nirenberg [8] in 1965 and then is used by Hörmander [7] and others for problems in partial differential equations. Weyl transforms which are a class of pseudo-differential operators have applications in Quantization due to Hermann Weyl [18]. In [21] Weyl transforms on compact Lie groups are introduced and the heat kernels of the Laplacian on the compact Lie group is obtained. In this paper, we look at pseudo-differential operators on compact and Hausdorff groups with L^2 symbols. The L^p , $1 \leq p \leq \infty$ conditions on the symbols allowing singularities are ideal for a broad spectrum of disciplines ranging from functional analysis to operator algebras to quantization. An analogue of the results in this paper is studied for the compact Lie group $\mathbb{S}^{n-1}$, i.e., the unit sphere

S. Molahajloo (✉)
Department of Mathematics and Statistics,
Queen's University, University Avenue,
Kingston, ON K7L3N6, Canada
e-mail: smollaha@gmail.com

M. Pirhayati
Institut für Mathematik, Universität Potsdam,
Am Neuen Palais 10, 14469 Potsdam, Germany

with center at the origin in [5]. Pseudo-differential operators on the unit sphere are studied extensively in [1–11, 13, 14, 16, 17, 22].

The aim of this paper is to give a characterization of trace class pseudo-differential operators on compact and Hausdorff groups. We give a formula for the trace of pseudo-differential operators in the trace class. The main technique is to obtain a formula for the symbol of the product of two pseudo-differential operators on a compact and Hausdorff group.

In Sect. 2, we give a brief recall of Hilbert-Schmidt and trace class operators. We define pseudo-differential operators on compact and Hausdorff groups by using the space of all unitary and irreducible representations in Sect. 3. A product formula for pseudo-differential operators is given. Then we give a characterization of trace class pseudo-differential operators.

2 Hilbert-Schmidt and trace class operators

Let $\mathcal{H}$ be a complex and separable Hilbert space in which the inner product and norm are denoted by $(\cdot, \cdot)_{\mathcal{H}}$ and $\|\cdot\|_{\mathcal{H}}$. Let $A : \mathcal{H} \rightarrow \mathcal{H}$ be a compact operator. Then the absolute value of A denoted by $|A|$ is defined by

$$|A| = (A^*A)^{1/2}.$$

The operator $|A|$ is compact and positive. So, by the spectral theorem, there exists an orthonormal basis $\{\varphi_j\}_{j=1}^{\infty}$ of eigenvectors of $|A|$ with the corresponding eigenvalues $\{s_j\}_{j=1}^{\infty}$ of real numbers. The operator A is said to be in the Hilbert-Schmidt class S_2 if

$$\sum_{j=1}^{\infty} s_j^2 < \infty,$$

and we define its Hilbert-Schmidt norm by

$$\|A\|_{HS} = \left(\sum_{j=1}^{\infty} s_j^2 \right)^{1/2}.$$

The operator A is said to be in the trace class S_1 if

$$\sum_{j=1}^{\infty} s_j < \infty,$$

and we define its trace by

$$tr(A) = \sum_{j=1}^{\infty} s_j.$$

We have the following theorem which shows that the Hilbert-Schmidt norm and trace is independent of the choice of orthonormal basis for $\mathcal{H}$, for details see [12] by Reed and Simon.

Theorem 2.1 *Let $A : \mathcal{H} \rightarrow \mathcal{H}$ be a bounded operator. Then*

- $A \in S_2$ if and only if there exists an orthonormal basis $\{\varphi_j\}_{j=1}^{\infty}$ for $\mathcal{H}$ such that

$$\|A\|_{HS} = \left(\sum_{j=1}^{\infty} \|A\varphi_j\|_{\mathcal{H}}^2 \right)^{1/2} < \infty.$$

- $A \in S_1$ if and only if there exists an orthonormal basis $\{\varphi_j\}_{j=1}^{\infty}$ for $\mathcal{H}$ such that

$$\text{tr}(A) = \sum_{j=1}^{\infty} (A\varphi_j, \varphi_j)_{\mathcal{H}} < \infty.$$

The following theorem will be useful in the next section.

Theorem 2.2 *Let $A : \mathcal{H} \rightarrow \mathcal{H}$ be a bounded operator. Then $A \in S_1$ if and only if $A = BC$, where B and C are in S_2 .*

3 Pseudo-differential operators on compact and Hausdorff groups

Let G be a locally compact and Hausdorff group. Let $\mathcal{H}$ be a separable and complex Hilbert space. We denote the group of all unitary operators by $U(\mathcal{H})$. A group homomorphism $\pi : G \rightarrow U(\mathcal{H})$ is said to be a unitary representation of G on $\mathcal{H}$ if it is strongly continuous, i.e., the mapping

$$g \mapsto \pi(g)h$$

is a continuous mapping for all $h \in \mathcal{H}$. The dimension of $\mathcal{H}$ is known as the degree or the dimension of the representation π .

A closed subspace M of $\mathcal{H}$ is said to be invariant with respect to the unitary representation $\pi : G \rightarrow U(\mathcal{H})$ if

$$\pi(g)M \subseteq M, \quad \forall g \in G.$$

A unitary representation $\pi : G \rightarrow U(\mathcal{H})$ is called irreducible if it has only the trivial invariant subspaces, i.e., $\{0\}$ and $\mathcal{H}$. The following theorem is crucial in defining pseudo-differential operators on compact Hausdorff groups. For its proof see [6, 20].

Theorem 3.1 *Let G be a compact and Hausdorff group. Then any irreducible and unitary representation of G on a complex and separable Hilbert space is finite dimensional.*

Let $\mathbb{G}$ be a compact and Hausdorff group with the left (and right) Haar measure is denoted by μ . Let $\hat{\mathbb{G}}$ be the set of all irreducible and unitary representations of $\mathbb{G}$.

Let $\pi \in \hat{\mathbb{G}}$. Then π is finite dimensional. Let a_π be the degree of π and let $\mathcal{H}_\pi$ be its representation space. We denote the inner product and the norm in $\mathcal{H}_\pi$ by $(\cdot, \cdot)_{\mathcal{H}_\pi}$ and $\|\cdot\|_{\mathcal{H}_\pi}$ respectively. Let $\{\psi_1, \psi_2, \dots, \psi_{a_\pi}\}$ be an orthonormal basis for $\mathcal{H}_\pi$. Then for all $j, k = 1, 2, \dots$, we define π_{jk} by

$$\pi_{jk}(g) = \sqrt{a_\pi} (\pi(g)\psi_k, \psi_j)_{\mathcal{H}_\pi}$$

Theorem 3.2 (Peter-Weyl Theorem) $\{\pi_{jk} : \pi \in \hat{\mathbb{G}}, j, k = 1, 2, \dots, a_\pi\}$ forms an orthonormal basis for $L^2(\mathbb{G})$.

By Peter-Weyl theorem, every $f \in L^2(\mathbb{G})$ can be expressed as

$$f = \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} (f, \pi_{jk})_{L^2(\mathbb{G})} \pi_{jk}.$$

have the following Plancherel theorem.

Theorem 3.3 For all $f \in L^2(\mathbb{G})$,

$$\|f\|_{L^2(\mathbb{G})} = \left\{ \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} |(f, \pi_{jk})_{L^2(\mathbb{G})}|^2 \right\}^{1/2}.$$

Let σ be a measurable function on $\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N}$. For all measurable functions f on $\mathbb{G}$, we define $T_\sigma f$ on $\mathbb{G}$ formally by

$$(T_\sigma f)(g) = \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sigma(g, \pi, j, k) (f, \pi_{jk})_{L^2(\mathbb{G})} \pi_{jk}(g), \quad g \in \mathbb{G}.$$

T_σ is called the pseudo-differential operator on $\mathbb{G}$ corresponding to the symbol σ . For pseudo-differential operators on Euclidean spaces see for example [15, 19].

Let $L^2(\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N})$ be the space all measurable functions σ on $\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N}$ for which

$$\|\sigma\|_{L^2(\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N})} = \left\{ \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \int_{\mathbb{G}} |\sigma(g, \pi, j, k) \pi_{jk}(g)|^2 d\mu(g) \right\}^{1/2} < \infty.$$

We have the following result on the L^2 -boundedness of pseudo-differential operators on $\mathbb{G}$.

Theorem 3.4 Let $\sigma \in L^2(\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N})$. Then $T_\sigma : L^2(\mathbb{G}) \rightarrow L^2(\mathbb{G})$ is a bounded operator and

$$\|T_\sigma\|_* \leq \|\sigma\|_{L^2(\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N})},$$

where $\|\cdot\|_*$ is the norm in the C^* -algebra of all bounded linear operators from $L^2(\mathbb{G})$ into $L^2(\mathbb{G})$.

Proof Let $f \in L^2(\mathbb{G})$. By Minkowski's inequality and the Schwarz inequality, we have

$$\begin{aligned}
 & \|T_\sigma f\|_{L^2(\mathbb{G})} \\
 &= \left(\int_{\mathbb{G}} |(T_\sigma f)(g)|^2 d\mu(g) \right)^{1/2} \\
 &= \left(\int_{\mathbb{G}} \left| \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sigma(g, \pi, j, k) (f, \pi_{jk})_{L^2(\mathbb{G})} \pi_{jk}(g) \right|^2 d\mu(g) \right)^{1/2} \\
 &\leq \sum_{\pi \in \hat{\mathbb{G}}} \left(\int_{\mathbb{G}} \left| \sum_{j,k=1}^{a_\pi} \sigma(g, \pi, j, k) (f, \pi_{jk})_{L^2(\mathbb{G})} \pi_{jk}(g) \right|^2 d\mu(g) \right)^{1/2} \\
 &\leq \sum_{\pi \in \hat{\mathbb{G}}} \left(\int_{\mathbb{G}} \left(\sum_{j,k=1}^{a_\pi} |(f, \pi_{jk})_{L^2(\mathbb{G})}|^2 \right) \left(\sum_{j,k=1}^{a_\pi} |\sigma(g, \pi, j, k) \pi_{jk}(g)|^2 \right) d\mu(g) \right)^{1/2} \\
 &= \sum_{\pi \in \hat{\mathbb{G}}} \left(\sum_{j,k=1}^{a_\pi} |(f, \pi_{jk})_{L^2(\mathbb{G})}|^2 \right)^{1/2} \left(\int_{\mathbb{G}} \sum_{j,k=1}^{a_\pi} |\sigma(g, \pi, j, k) \pi_{jk}(g)|^2 d\mu(g) \right)^{1/2} \\
 &\leq \left(\sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} |(f, \pi_{jk})_{L^2(\mathbb{G})}|^2 \right)^{1/2} \left(\sum_{\pi \in \hat{\mathbb{G}}} \int_{\mathbb{G}} \sum_{j,k=1}^{a_\pi} |\sigma(g, \pi, j, k) \pi_{jk}(g)|^2 d\mu(g) \right)^{1/2} \\
 &= \|f\|_{L^2(\mathbb{G})} \|\sigma\|_{L^2(\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N})}.
 \end{aligned}$$

□

Now we are ready to give a necessary and sufficient condition on σ to guarantee Hilbert-Schmidt properties of T_σ .

Theorem 3.5 *Let σ be a measurable function on $\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N}$. Then T_σ is a Hilbert-Schmidt operator if and only if $\sigma \in L^2(\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N})$ and*

$$\|T_\sigma\|_{HS} = \|\sigma\|_{L^2(\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N})}.$$

Proof For $\tilde{\pi} \in \hat{\mathbb{G}}$ and $j_0, k_0 = 1, 2, \dots, a_{\tilde{\pi}}$, we have

$$\begin{aligned}
 (T_\sigma \tilde{\pi}_{j_0 k_0})(g) &= \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=0}^{a_\pi} \sigma(g, \pi, j, k) (\tilde{\pi}_{j_0 k_0}, \pi_{jk})_{L^2(\mathbb{G})} \pi_{jk}(g) \\
 &= \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=0}^{a_\pi} \sigma(g, \pi, j, k) (\tilde{\pi}_{j_0 k_0}, \pi_{jk})_{L^2(\mathbb{G})} \pi_{jk}(g) \\
 &= \sigma(g, \tilde{\pi}, k_0, j_0) \tilde{\pi}_{j_0 k_0}(g), \quad g \in \mathbb{G}.
 \end{aligned}$$

Hence

$$\begin{aligned}
 & \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \|T_\sigma \pi_{jk}\|_{L^2(\mathbb{G})}^2 \\
 &= \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \left(\int_{\mathbb{G}} |T_\sigma \pi_{jk}(g)|^2 d\mu(g) \right) \\
 &= \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \int_{\mathbb{G}} |\sigma(g, \pi, j, k) \pi_{jk}(g)|^2 d\mu(g) \\
 &= \|\sigma\|_{L^2(\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N})}^2.
 \end{aligned}$$

and the proof is complete. $\square$

Let σ and τ measurable functions on $\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N}$. Then we define $\sigma \circledast \tau$ by

$$\begin{aligned}
 & \sigma \circledast \tau(g, \xi, l, m) \\
 &= \left(\int_{\mathbb{G}} \tau(w, \xi, l, m) \xi_{lm}(w) \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sigma(g, \pi, j, k) \overline{\pi_{jk}(w)} \pi_{jk}(g) d\mu(w) \right) (\xi_{lm}(g))^{-1}
 \end{aligned} \tag{3.1}$$

for all $g \in \mathbb{G}$, $\xi \in \hat{\mathbb{G}}$ and $l, m = 1, \dots, a_\xi$. The following theorem shows that the composition of two pseudo-differential operators on $\mathbb{G}$ is again a pseudo-differential operator.

Theorem 3.6 *Let σ and τ be measurable functions on $\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N}$. Then*

$$T_\sigma T_\tau = T_\lambda$$

where $\lambda = \sigma \circledast \tau$.

Proof Let f be in $L^2(\mathbb{G})$. Then for all $g \in \mathbb{G}$

$$\begin{aligned}
 & (T_\sigma T_\tau f)(g) \\
 &= \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sigma(g, \pi, j, k) (T_\tau f, \pi_{jk})_{L^2(\mathbb{G})} \pi_{jk}(g) \\
 &= \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sigma(g, \pi, j, k) \left(\int_{\mathbb{G}} (T_\tau f)(w) \overline{\pi_{jk}(w)} d\mu(w) \right) \pi_{jk}(g) \\
 &= \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sigma(g, \pi, j, k) \left(\int_{\mathbb{G}} \sum_{\xi \in \hat{\mathbb{G}}} \sum_{l,m=1}^{a_\xi} \tau(w, \xi, l, m) (f, \xi_{lm})_{L^2(\mathbb{G})} \xi_{lm}(w) \overline{\pi_{jk}(w)} d\mu(w) \right) \pi_{jk}(g)
 \end{aligned}$$

$$\begin{aligned}
&= \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sigma(g, \pi, j, k) \left(\sum_{\xi \in \hat{\mathbb{G}}} \sum_{l,m=1}^{a_\xi} (f, \xi_{lm})_{L^2(\mathbb{G})} \int_{\mathbb{G}} \tau(w, \xi, l, m) \xi_{lm}(w) \overline{\pi_{jk}(w)} d\mu(w) \right) \pi_{jk}(g) \\
&= \sum_{\xi \in \hat{\mathbb{G}}} \sum_{l,m=1}^{a_\xi} \left(\int_{\mathbb{G}} \tau(w, \xi, l, m) \xi_{lm}(w) \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sigma(g, \pi, j, k) \overline{\pi_{jk}(w)} \pi_{jk}(g) d\mu(w) \right) \\
&\quad \times (\xi_{lm}(g))^{-1} (f, \xi_{lm})_{L^2(\mathbb{G})} \xi_{lm}(g) \\
&= \sum_{\xi \in \hat{\mathbb{G}}} \sum_{l,m=1}^{a_\xi} \lambda(g, \xi, l, m) (f, \xi_{lm})_{L^2(\mathbb{G})} \xi_{lm}(g), \tag{3.2}
\end{aligned}$$

where $\lambda = \sigma \circledast \tau$. Thus,

$$T_\lambda = T_\sigma T_\tau$$

□

Now we are ready to give a characterization of trace class pseudo-differential operators on $\mathbb{G}$.

Theorem 3.7 *Let λ be a measurable function in $\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N}$. Then $T_\lambda \in S_1$ if and only if there exist symbols σ and τ in $L^2(\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N})$ such that*

$$\lambda = \sigma \circledast \tau,$$

and

$$\begin{aligned}
tr(T_\lambda) &= \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \int_{\mathbb{G}} \lambda(g, \pi, j, k) |\pi_{jk}(g)|^2 d\mu(g) \\
&= \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sum_{\xi \in \hat{\mathbb{G}}} \sum_{l,m=1}^{a_\xi} (\tau(\cdot, \xi, l, m) \xi_{lm}, \pi_{jk})_{L^2(\mathbb{G})} (\sigma(\cdot, \pi, j, k) \pi_{jk}, \xi_{lm})_{L^2(\mathbb{G})}.
\end{aligned}$$

Proof The first part of the theorem follows from Theorem 2.2, Theorem 3.5 and Theorem 3.6. Now we check the absolute convergence of the series

$$\sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sum_{\xi \in \hat{\mathbb{G}}} \sum_{l,m=1}^{a_\xi} (\tau(\cdot, \xi, l, m) \xi_{lm}, \pi_{jk})_{L^2(\mathbb{G})} (\sigma(\cdot, \pi, j, k) \pi_{jk}, \xi_{lm})_{L^2(\mathbb{G})}$$

By the Schwarz inequality and Plancherel's theorem,

$$\begin{aligned}
&\sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sum_{\xi \in \hat{\mathbb{G}}} \sum_{l,m=1}^{a_\xi} \left| (\tau(\cdot, \xi, l, m) \xi_{lm}, \pi_{jk})_{L^2(\mathbb{G})} \right| \left| (\sigma(\cdot, \pi, j, k) \pi_{jk}, \xi_{lm})_{L^2(\mathbb{G})} \right| \\
&\leq \left\{ \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sum_{\xi \in \hat{\mathbb{G}}} \sum_{l,m=1}^{a_\xi} \left| (\tau(\cdot, \xi, l, m) \xi_{lm}, \pi_{jk})_{L^2(\mathbb{G})} \right|^2 \right\}^{1/2}
\end{aligned}$$

$$\begin{aligned}
& \left\{ \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sum_{\xi \in \hat{\mathbb{G}}} \sum_{l,m=1}^{a_\xi} \left| (\sigma(\cdot, \pi, j, k) \pi_{jk}, \xi_{lm})_{L^2(\mathbb{G})} \right|^2 \right\}^{1/2} \\
&= \left\{ \sum_{\xi \in \hat{\mathbb{G}}} \sum_{l,m=1}^{a_\xi} \int_{\mathbb{G}} |\tau(g, \xi, l, m) \xi_{lm}(g)|^2 d\mu(g) \right\}^{1/2} \\
& \left\{ \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \int_{\mathbb{G}} |\sigma(g, \pi, j, k) \pi_{jk}(g)|^2 d\mu(g) \right\}^{1/2} \\
&= \|\sigma\|_{L^2(\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N})} \|\tau\|_{L^2(\mathbb{G} \times \hat{\mathbb{G}} \times \mathbb{N} \times \mathbb{N})} < \infty.
\end{aligned}$$

Since $\bigcup_{\pi \in \hat{\mathbb{G}}} \{\pi_{jk} : j, k = 1, 2, \dots, a_\pi\}$ forms an orthonormal basis for $L^2(\mathbb{G})$, it follows that

$$\begin{aligned}
tr(T_\lambda) &= \sum_{\xi \in \hat{\mathbb{G}}} \sum_{l,m=1}^{a_\xi} (T_\lambda \xi_{lm}, \xi_{lm})_{L^2(\mathbb{G})} \\
&= \sum_{\xi \in \hat{\mathbb{G}}} \sum_{l,m=1}^{a_\xi} \int_{\mathbb{G}} \lambda(g, \xi, l, m) \xi_{lm}(g) \overline{\xi_{lm}(g)} d\mu(g) \\
&= \sum_{\xi \in \hat{\mathbb{G}}} \sum_{l,m=1}^{a_\xi} \int_{\mathbb{G}} \left(\int_{\mathbb{G}} \tau(w, \xi, l, m) \xi_{lm}(w) \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sigma(g, \pi, j, k) \overline{\pi_{jk}(w)} \pi_{jk}(g) d\mu(w) \right) \\
&\quad \times \overline{\xi_{lm}(g)} d\mu(g) \\
&= \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sum_{\xi \in \hat{\mathbb{G}}} \sum_{l,m=1}^{a_\xi} \int_{\mathbb{G}} \tau(w, \xi, l, m) \xi_{lm}(w) \overline{\pi_{jk}(w)} d\mu(w) \int_{\mathbb{G}} \sigma(g, \pi, j, k) \pi_{jk}(g) \\
&\quad \times \overline{\xi_{lm}(g)} d\mu(g) \\
&= \sum_{\pi \in \hat{\mathbb{G}}} \sum_{j,k=1}^{a_\pi} \sum_{\xi \in \hat{\mathbb{G}}} \sum_{l,m=1}^{a_\xi} (\tau(\cdot, \xi, l, m) \xi_{lm}, \pi_{jk})_{L^2(\mathbb{G})} (\sigma(\cdot, \pi, j, k) \pi_{jk}, \xi_{lm})_{L^2(\mathbb{G})}.
\end{aligned}$$

□

We can now look at the special case when $\mathbb{G} = \mathbb{S}^1$.

Example 3.8 Let $\mathbb{S}^1$ be the unit circle centered at the origin. For all $n \in \mathbb{Z}$, we define $\pi_n : \mathbb{S}^1 \rightarrow U(1)$ by

$$\pi_n(\theta) = (2\pi)^{-1/2} e^{in\theta}, \quad \theta \in [-\pi, \pi].$$

Then

$$\widehat{\mathbb{S}^1} = \{\pi_n : n \in \mathbb{Z}\},$$

and $\widehat{\mathbb{S}} \simeq \mathbb{Z}$. Hence for any symbol σ on $\mathbb{S}^1 \times \mathbb{Z}$ we can define pseudo-differential operator T_σ on $L^2(\mathbb{S}^1)$ by

$$(T_\sigma f)(\theta) = \sum_{n \in \mathbb{Z}} \sigma(\theta, n) (f, \pi_n)_{L^2(\mathbb{S}^1)} \pi_n(\theta), \quad f \in L^2(\mathbb{S}^1), \quad \theta \in [-\pi, \pi].$$

Hence,

$$(T_\sigma f)(\theta) = (2\pi)^{-1} \sum_{n \in \mathbb{Z}} \int_{-\pi}^{\pi} e^{in(\theta-\varphi)} \sigma(n, \theta) f(\varphi) d\varphi.$$

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

1. Agranovich, M.S.: Elliptic pseudodifferential operators on a closed curve, (Russian). Trudy Moskov. Mat. Obshch. **47**(246), 22–67 (1984)
2. Agranovich, M.S.: Spectral properties of elliptic pseudodifferential operators on a closed curve, (Russian). Funktsional. Anal. I Prilozhen **13**, 54–56 (1979)
3. Amosov, B.A.: On the theory of pseudodifferential operators on the circle, (Russian). Uspekhi Mat. Nauk. **43**, 169–170 (1988)
4. Amosov, B.A.: Translation in Russian. Math. Surv. **43**, 197–198 (1988)
5. Chen, Z., Wong, M.W.: Traces of pseudo-differential operators on $\widehat{\mathbb{S}}^{n-1}$. J. Pseudo-Differ. Oper. Appl. **4**(1), 13–24 (2013)
6. Folland, G.B.: A course in abstract Harmonic analysis. CRC Press (1995)
7. Hörmander, L.: The analysis of linear partial differential operators III. Springer-Verlag, Berlin (1985)
8. Kohn, J.J., Nirenberg, L.: An algebra of pseudo-differential operators. Comm. Pure Appl. Math. **18**, 269–305 (1965)
9. Molahajloo, S.: A characterization of compact pseudo-differential operators on $\mathbb{S}^1$, in pseudo-differential operators: analysis, applications and computations 213. Birkhäuser, Basel (2011)
10. Molahajloo, S., Wong, M.W.: Ellipticity, fredholmness and spectral invariance of pseudo-differential operators on $\mathbb{S}^1$. J. Pseudo-Differ. Oper. Appl. **1**(2), 183–205 (2010)
11. Pirhayati, M.: Spectral theory of pseudo-differential operators on $\mathbb{S}^1$, in pseudo-differential operators: analysis, applications and computations. Birkhäuser, Basel (2011)
12. Reed, M., Simon, B.: Functional analysis. Academic Press, New York (1980)
13. Ruzhansky, M., Turunen, V.: On the Fourier analysis of operators on the torus. In: Toft, J., Wong, M.W., Zhu, H. (eds.) Modern Trends in Pseudo-Differential Operators, pp. 87–105. Birkhäuser (2007)
14. Saranen, J., Vainikko, G.: Periodic integral and pseudodifferential equations with numerical approximation. Springer-Verlag, Berlin (2002)
15. Stein, E.M., Weiss, G.: Introduction to Fourier analysis on Euclidean spaces. Princeton University Press, Princeton (1971)
16. Weiss, N.J.: Multipliers on compact Lie groups. Proc. Nat. Acad. Sci. U.S.A. **68**, 930–931 (1971)
17. Weiss, N.J.: L^p estimates for bi-invariant operators on compact Lie groups. Am. J. Math **94**, 103–118 (1972)
18. Weyl, H.: The theory of groups and quantum mechanics. Dover (1950)
19. Wong, M.W.: An introduction to pseudo-differential operators, 2nd edn. World Scientific, Canada (1999)
20. Wong, M.W.: Wavelet transforms and localization operators. Birkhäuser, Basel (2002)
21. Wong, M.W.: Weyl transforms, heat kernels. Green functions and Riemann zeta functions on compact lie groups, 172nd edn, pp. 67–85. Operator Theory: Advances and Applications Birkhäuser Verlag, Basel (2006)
22. Wong, M.W.: Discrete Fourier analysis. Birkhäuser, Basel (2011)