ERRATUM

Erratum to: Solid-State Quantitative ^1H and ^{31}P MRI of Cortical Bone in Humans

Alan C. Seifert¹ • Felix W. Wehrli¹

Published online: 24 May 2016
© Springer Science+Business Media New York 2016

Erratum to: Curr Osteoporos Rep (2016) 14(3):77–86
DOI 10.1007/s11914-016-0307-2

The original version of this article unfortunately contained a mistake. While the figure captions for Figures 2 and 3 were placed correctly, the accompanying images were swapped. The corrected figures are given below.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s11914-016-0307-2>.

✉ Felix W. Wehrli
wehrli@mail.med.upenn.edu

¹ Laboratory for Structural, Physiologic, and Functional Imaging,
Department of Radiology, University of Pennsylvania Medical
Center, MRI Education Center, 1st Floor Founders, 3400 Spruce St.,
Philadelphia, PA 19104, USA

Fig. 2 Parametric maps of ^{31}P (a) and bound water ^1H (b) concentrations (mol/L) in human tibial cortical bone acquired ex vivo at 3 and 7 T, respectively (reproduced from [35••]). ^{31}P and bound water ^1H were correlated ($R^2 = 0.59$) in these data from non-osteomalacic donors. (c, d) In vivo images of cortical bone ^{31}P in the human wrist (reproduced from [53])

Fig. 3 Parametric maps of the suppression ratio (SR) obtained with and without long- T_2 suppression in six subjects (a), along with the corresponding histograms (b). As SR values increase, histograms become increasingly asymmetric with long tails toward high SR values, commensurate with the notion of elevated T_2 of the water in larger pores (reproduced from [12•]). Maps of bound (c) and pore (d) water concentrations, measured by adiabatic inversion recovery and double adiabatic full passage UTE MRI, respectively (reproduced from [34•])