Letter to the Editor

New Equations for Predicting Postoperative Risk in Patients with Hip Fracture

Ian Moppett DM, MRCP, FRCA

Published online: 13 April 2010
© The Association of Bone and Joint Surgeons® 2010

To the Editor:

I read with interest the paper by Hirose et al. [2] describing the use of E-PASS to predict postoperative risk based on preoperative factors in patients with hip fractures. The authors are to be congratulated on attempting to validate their risk prediction in a separate cohort of patients. However, I have some concerns regarding the data as presented.

First, the subjects chosen were a nonrepresentative population. Patients with no complications were excluded from the study groups, yet this group represents a substantial proportion of patients with hip fractures. As the definition of lack of complications is a retrospective finding, it is not clear how the score then can be used preoperatively.

Second, in addition to the comments made by Zhou and Fan [6], I question the appropriateness of the analysis performed. For a score to be assessed as reliable, there should be some statistical assessment. Hirose et al. report observed-to-estimated ratios, but make no statistical assessment. The Hosmer-Lemeshow test is an appropriate test in these circumstances [3].

Third, I question the rationale for the comparisons made in the paper. POSSUM previously was shown to be not particularly useful in this population [5]. The surgical stress

score has a very narrow range in this population, and therefore is unlikely to have discriminatory ability. As Hirose et al. stated, several scoring systems have been developed. Maxwell et al. [4] reported on a large series of patients, validating a hip fracture score, which is based on recognized risk factors for postoperative mortality, and comparing it with the ASA and Donati scores [1]. Thus, it is unclear why E-PASS would be a better technique than those that have gone before.

References

1. Donati A, Ruzzi M, Adrario E, Pelaia P, Coluzzi F, Gabbanelli V, Pietropaoli P. A new and feasible model for predicting operative risk. *Br J Anaesth*. 2004;93:393–399.
2. Hirose J, Ide J, Irie H, Kikukawa K, Mizuta H. New equations for predicting postoperative risk in patients with hip fracture. *Clin Orthop Relat Res*. 2009;467:3327–3333.
3. Lemeshow S, Hosmer DW Jr. A review of goodness of fit statistics for use in the development of logistic regression models. *Am J Epidemiol*. 1982;115:92–106.
4. Maxwell MJ, Moran CG, Moppett IK. Development and validation of a preoperative scoring system to predict 30 day mortality in patients undergoing hip fracture surgery. *Br J Anaesth*. 2008;101: 511–517.
5. Ramanathan TS, Moppett IK, Wenn R, Moran CG. POSSUM scoring for patients with fractured neck of femur. *Br J Anaesth*. 2005;94:430–433.
6. Zhou Z, Fan S. Letter to the editor: New equations for predicting postoperative risk in patients with hip fracture. *Clin Orthop Relat Res*. 2010 Feb 6. (Epub ahead of print)

(Re: Hirose J, Ide J, Irie H, Kikukawa K, Mizuta H. New equations for predicting postoperative risk in patients with hip fracture. *Clin Orthop Relat Res*. 2009;467:3327–3333.)

I. Moppett (✉)
University of Nottingham, Anaesthesia and Intensive Care,
Queen's Medical Centre Campus, Nottingham University
Hospitals NHS Trust, Derby Road,
Nottingham, Notts NG7 2UH, UK
e-mail: iain.moppett@nottingham.ac.uk