Erratum to: Infirmary and Injury Complexity are Risk Factors for Surgical-site Infection after Operative Fracture Care

Abdo Bachoura MD, Thierry G. Guitton MSc,
R. Malcolm Smith MD, Mark S. Vrahas MD,
David Zurakowski PhD, David Ring MD, PhD

Published online: 29 September 2011
© The Association of Bone and Joint Surgeons® 2011

Erratum to: Clin Orthop Relat Res (2011) 469:2621–2630 DOI 10.1007/s11999-010-1737-2

Due to a formatting error, in Table 8, the data for the second study by Minnema et al. were shifted to the right, thereby showing the data under the incorrect column headings. The table in correct format is shown below.

The online version of the original article can be found under doi:
[10.1007/s11999-010-1737-2](https://doi.org/10.1007/s11999-010-1737-2).

A. Bachoura, T. G. Guitton, D. Ring
Orthopaedic Hand and Upper Extremity Service, Massachusetts
General Hospital, Harvard Medical School, Boston, MA, USA

R. M. Smith, M. S. Vrahas
Department of Orthopaedics, Massachusetts General Hospital,
Harvard Medical School, Boston, MA, USA

D. Zurakowski
Departments of Anesthesia and Surgery, Children's Hospital
Boston, Harvard Medical School, Boston, MA, USA

D. Ring (✉)
Massachusetts General Hospital, Yawkey Center, Suite 2100,
55 Fruit Street, Boston, MA 02114, USA
e-mail: dring@partners.org

Table 8. Comparison of incidence and independent predictors of SSI in the orthopaedic literature

Study	Subspecialty	SSI overall incidence	Independent predictors of SSI (odds ratio, 95% CI)				CHF	Wound drain	Site	Preoperative length of stay
			Wound class	Prophylactic antibiotics	Number of surgeries	Diabetes				
Ridgeway et al. [37]	Hip arthroplasty	3.1%	No	No	Not assessed	Not assessed	Not assessed	Not assessed	Not assessed	No
Minnema et al. [22]	Knee arthroplasty	Not assessed	Not assessed	No	Not assessed	Not assessed	Not assessed	Closed suction drain 7.0(2.1–25)	Not assessed	Not assessed
Pulido et al. [34]	Joint arthroplasty	0.7%	Not assessed	Not assessed	Not assessed	No	No	Not assessed	Not assessed	Not assessed
Olsen et al. [26]	Spine	2.0%	Not assessed	Suboptimal timing 3.4(1.5–7.9)	Not assessed	3.5(1.2–10)	Not assessed	No	Cervical spine 0.3(0.1–0.6)	Not assessed
Pull ter Gunne & Cohen [35]	Spine	4.2%	Not assessed	Not assessed	Not assessed	2.2(1.2–4.0)	Not assessed	Not assessed	Not assessed	Not assessed
Pull ter Gunne et al. [36]	Spine deformity	5.5%	Not assessed	Not assessed	No	No	Not assessed	Not assessed	Not assessed	Not assessed
Blam et al. [3]	Spine trauma	9.4%	Not assessed	Not assessed	Not assessed	Not assessed	Not assessed	Not assessed	Not assessed	(In ICU) No
Simchen et al. [39]	General	4.8%	Not assessed	No	> 2 operation 9.6(8.2–11.0)	Not assessed	Not assessed	Open drain 4.6(3.8–6.5)	No	Not assessed
De Boer et al. [7]	General	1.1%	Not assessed	Not assessed	> 1 operation 2.5(1.9–3.0)	Not assessed	Not assessed	Not assessed	Not assessed	> 4 days 3.3(2.5–4.0)
		2.2%	Dirty & contaminated 4.1(3.7–4.5)	1.5(1.2–1.9)	> 1 operation 2.4(1.8–2.9)	Not assessed	Not assessed	Not assessed	Not assessed	> 4 days 2.4(2.0–2.9)
Maksimovic et al. [20]	General	22.7%	Dirty & contaminated 12.1(5.6–26.3)	Not assessed	Not assessed	No	Not assessed	No	Not assessed	Not assessed
Thu et al. [42]	Trauma	12.5%	Dirty 8.7(4.6–16.4)	No	Not assessed	Not assessed	Not assessed	No	Not assessed	No
Herruzo-Cabrera et al. [14]	Trauma	3.9%	Dirty/clean 10.5(5.1–21.9)	Incorrect/correct 1.5(1–2.2)	Not assessed	Not assessed	Not assessed	Not assessed	Not assessed	> 4/4 days 1.6(1.1–2.4)
Current study	Trauma	4.2%	No	Not assessed	> 1 operation 3.4(2.0–6.0)	2.1(1.2–3.8)	2.8(1.3–6.5)	2.3(1.3–3.8)	Elbow 2.2(1.1–4.7) tibial shaft/plateau 2.3(1.3–4.)	No

SSI = surgical-site infection; CHF = congestive heart failure.