Erratum to: Catalytic Properties of Two *Rhizopus oryzae* 99-880 Glucoamylase Enzymes Without Starch Binding Domains Expressed in *Pichia pastoris*

Jeffrey A. Mertens · Jay D. Braker · Douglas B. Jordan

Published online: 13 July 2010

© Springer Science+Business Media, LLC 2010

Erratum to: Appl Biochem Biotechnol
DOI 10.1007/s12010-010-8994-0

In the original publication, Fig. 3a and b are the same image. The corrected figure is reproduced here.

Fig. 3 Temperature stabilities and optima of AmyC and AmyD. Temperature stabilities of AmyC at pH 5.5 and AmyD at pH 6.0 were determined by pre-incubating the enzymes for 30 min at 20 °C to 80 °C followed by activity assays (1% soluble starch) at 40 °C. Temperature optima of AmyC and AmyD were determined by performing activity assays (1% soluble starch) at temperatures ranging from 20 °C to 80 °C. For temperature optima, 100% activity corresponds to rate of 94.9 s⁻¹ and 49.7 s⁻¹ for AmyC and AmyD, respectively

The online version of the original article can be found at <http://dx.doi.org/10.1007/s12010-010-8994-0>.

J. A. Mertens (✉) · J. D. Braker · D. B. Jordan
Bioenergy Research Unit, National Center for Agricultural Utilization Research,
USDA Agricultural Research Service, 1815 N. University St., Peoria, IL 61604, USA
e-mail: jeffrey.mertens@ars.usda.gov