Letter regarding Guddat et al. “Fatal spontaneous subdural bleeding due to neonatal giant cell hepatitis: a rare differential diagnosis of shaken baby syndrome”

Neil E. I. Langlois

Published online: 24 September 2011
© Springer Science+Business Media, LLC 2011

Dear Sir,

I read with interest the report by Guddat and colleagues regarding a case of fatal spontaneous subdural haematoma attributed to neonatal giant cell hepatitis [1]. As it has been noted that intraocular findings may vary depending on the cause [2, 3] I would be grateful if the authors could describe the position of the hemorrhages within the layers of the retina and their distribution around the globe, with a comment if retinoschisis or any perimacular folds were present. This may be helpful for future comparison between traumatic and non-traumatic causes of retinal changes.

References

1. Guddat SS, Ehlich E, Martin H, Tsokos M. Fatal spontaneous subdural bleeding due to neonatal giant cell hepatitis: a rare differential diagnosis of shaken baby syndrome. *Forensic Sci Med Pathol*. 2011;7:294–7.
2. Levin AV. Retinal hemorrhages: advances in understanding. *Pediatr Clin N Am*. 2009;56:333–44.
3. Bhardwaj G, Chowdhury V, Jacobs MB, Moran KT, Martin FJ, Coroneo MT. A systematic review of the diagnostic accuracy of ocular signs in pediatric abusive head trauma. *Ophthalmology*. 2010;117:983–92.

N. E. I. Langlois (✉)
Forensic Science South Australia & University of Adelaide,
21 Divett Place, Adelaide, SA 5000, Australia
e-mail: Neil.Langlois@sa.gov.au