ERRATUM

Erratum to: The Clinical Utility of TIMP3 Expression in ACTH-Secreting Pituitary Tumor

Bowen Sun¹ · Xiaohai Liu¹ · Yakun Yang¹ · Congxin Dai¹ · Ying Li² · Yonghui Jiao¹ · Zhenqing Wei¹ · Yong Yao¹ · Ming Feng¹ · Xinjie Bao¹ · Kan Deng¹ · Ning Wang² · Renzhi Wang¹

Published online: 19 February 2016
© Springer Science+Business Media New York 2016

Erratum to: J Mol Neurosci (2016) 58:137–144
DOI: 10.1007/s12031-015-0698-z

The original version of this article unfortunately contained mistakes in Figs. 3 and 4. The correct versions are presented below:

Fig. 3 **a** TIMP3 protein levels in ACTH-secreting pituitary adenomas and adjacent non-tumorous pituitary tissues assessed by Western blotting. Graph shows the TIMP3 GAPDH relative ratio in 10 tumors samples and

10 ANPT. $p < 0.001$. **b** Panel representative western blots of 5 tumors and 4 ANPT samples. *ACTH* adrenocorticotrophic hormone, *ANPT* non-tumorous pituitary tissue

The online version of the original article can be found at <http://dx.doi.org/10.1007/s12031-015-0698-z>.

✉ Renzhi Wang
wrzpunch@163.com

¹ Department of Neurosurgery, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences, Peking Union Medical College, Beijing 100730, China

² Department of Neurosurgery, the First Affiliated Hospital, Harbin Medical University, Harbin 150001, China

Fig. 4 **a** TIMP3 mRNA expression in invasive and noninvasive PAs. TIMP3 mRNA levels were significantly higher in the noninvasive PAs than in the invasive PAs ($p < .05$). **b** TIMP3 protein levels in different subtype pituitary adenomas and adjacent non-tumorous pituitary tissues

assessed by Western blotting, and compared invasive PAs with noninvasive PAs. TIMP3 tissue inhibitor of metalloproteinase 3, NF nonfunctional adenoma, PRL prolactin, GH growth hormone, ANPT non-tumorous pituitary tissue