ERRATUM

Erratum to: Third-line treatment of colorectal liver metastases using DEBIRI chemoembolization

Germano Scevola¹ · Giorgio Loreni¹ · Marco Rastelli¹ · Stefano Sposato¹ · Sara Ramponi² · Vittorio Miele³

Published online: 6 April 2017
© Springer Science+Business Media New York 2017

Erratum to: Med Oncol (2017) 34:37
DOI 10.1007/s12032-017-0890-9

The original version of the article unfortunately contained a mistake. The author of this article has not included the figures for this article. This is now corrected through this erratum, and the figures for this article are given (Figs. 1, 2 and 3).

The online version of the original article can be found under doi:[10.1007/s12032-017-0890-9](https://doi.org/10.1007/s12032-017-0890-9).

✉ Germano Scevola
germano.scevola@gmail.com

¹ Department of Interventional Radiology, S. Pertini Hospital, Rome, Italy

² Department of Medical Oncology, S. Pertini Hospital, Rome, Italy

³ Department of Radiology, Careggi University Hospital, Florence, Italy

Fig. 1 Case N°1. **a** CT scan, single focal liver metastasis from colorectal cancer in the VIII segment, **b** selective embolization of the VIII segment, **c** final check with selective catheterization of the hepatic artery at the end of treatment DEBIRI: embolization of

branches of VIII segment with stopping the flow, and **d** CT scan performed at 26-month follow-up demonstrates significant reduction in lesion

Fig. 2 Case N°2. **a** CT scan, single focal liver metastasis from colorectal cancer in the IV segment, **b** selective embolization and DEBIRI treatment of the segment IV, and **c** CT scan after 24 months demonstrates a hypodense necrotic metastasis in IV segment

Fig. 3 Case N°3. **a** CT scan, recurrent single focal liver metastasis from colorectal cancer in the VIII segment, previously treated with RF thermoablation, **b** selective embolization and DEBIRI treatment of the segment VIII, and **c** CT scan after 32 months demonstrates a hypodense necrotic metastasis in VIII segment