$$\int_{t_1}^{t_2} \frac{dA}{dt} \cdot dt = \frac{1}{2}l(t_2 - t_1).$$

The radius vector joining the planet and the Sun sweeps out equal areas in equal intervals of time, which is Kepler's second law.

In the case of an ellipse, the area covered in one orbit is

$$A = \pi a \cdot b,$$

where a is the semi-major axis and b is the semi-minor axis. In terms of the eccentricity ε , $b = a\sqrt{1 - \varepsilon^2}$. The rate at which the radius vector sweeps out the orbit is $dA/dt = l/2$, a constant. So the period of an orbit is $P = A/\dot{A} = (2\pi a^2\sqrt{1 - \varepsilon^2})/l$. Therefore

$$\left(\frac{P}{2\pi}\right)^2 = \frac{a^4(1 - \varepsilon^2)}{l^2} = \frac{a^4(1 - \varepsilon^2)}{pGM} = \frac{a^4(1 - \varepsilon^2)}{GM \cdot a(1 - \varepsilon^2)} = \frac{a^3}{GM}.$$

$$P^2 \propto a^3$$

This is Kepler's third law.

A detailed derivation of the laws and properties of ellipses can be found in Wikipedia, the free encyclopedia.

Address for Correspondence: Renuka Ravindran, Atria Institute of Technology, Bangalore 560 024, India.
Email: renrav@math.iisc.ernet.in

Errata

Resonance, Vol 14, No. 11, November 2009

The Solar Physics Observatory at Kodaikanal and John Evershed (pp. 1032–1039):

Due to an inadvertence on the part of the author, the article did not carry the credits to go with the photographs. The photographs appearing in the article were obtained from the IIA Archives. The error is regretted.

What is the Unit of Natural Selection (pp. 1047–1059): We regret that the author's name Ambika Kamath has been misspelt as Ambika Karanth.