Erratum to: Evaluation of primary prostate cancer using ^{11}C -methionine-PET/CT and ^{18}F -FDG-PET/CT

Masato Shiiba · Keiichi Ishihara · Go Kimura · Tomoyuki Kuwako ·
Hisashi Yoshihara · Hidetaka Sato · Yukihiro Kondo ·
Shin-ichi Tsuchiya · Shin-ichiro Kumita

Published online: 14 February 2012
© The Japanese Society of Nuclear Medicine 2012

Erratum to: Ann Nucl Med
DOI 10.1007/s12149-011-0551-6

The fifth coauthor's name was incorrectly given as Naohisa Yoshihara. It should be Hisashi Yoshihara. In the affiliation line, it should be shown as H. Yoshihara, not N. Yoshihara.

Also, the title and the contents of Table 2 were incorrect. They should be given as shown below.

Table 2 Correlation coefficients between PSA level and SUV_{max}

	All specimens	Excluding metastases
MET	0.132 ($P = 0.07$)	0.578 ($P = 0.00$)
FDGe	0.103 ($P = 0.17$)	0.430 ($P = 0.03$)
FDGd	0.100 ($P = 0.18$)	0.461 ($P = 0.03$)

PSA Prostate Specific Antigen, MET methionine, FDGe FDG early phase, FDGd FDG delayed phase

The online version of the original article can be found under doi:
[10.1007/s12149-011-0551-6](https://doi.org/10.1007/s12149-011-0551-6).

M. Shiiba (✉) · K. Ishihara · T. Kuwako · H. Yoshihara ·
H. Sato · S. Kumita
Department of Radiology, Nippon Medical School,
1-1-5, Sendagi, Bunkyo-ku, Tokyo 113-0022, Japan
e-mail: m-shiiba@nms.ac.jp

G. Kimura · Y. Kondo
Department of Urology, Nippon Medical School, Tokyo, Japan

S. Tsuchiya
Department of Diagnostic Pathology, Nippon Medical School
Main Hospital, Tokyo, Japan