Erratum to: An effective salvage treatment using ifosfamide, etoposide, cytarabine, dexamethasone, and rituximab (R-IVAD) for patients with relapsed or refractory aggressive B-cell lymphoma

Katsuhiro Miura · Kazuhiro Takei · Sumiko Kobayashi · Satomi Kiso · Yukio Hirabayashi · Atsuko Hojo · Hitomi Kodaira · Mai Yagi · Daisuke Kurita · Yujin Kobayashi · Toshitake Tanaka · Noriyoshi Iriyama · Yoshihiro Hatta · Yoshimasa Kura · Tetsuo Yamazaki · Umihiko Sawada · Jin Takeuchi

Published online: 14 July 2011
© The Japanese Society of Hematology 2011

**Erratum to: Int J Hematol (2011) 94:90–96
DOI 10.1007/s12185-011-0884-x**

An error appeared in the original version of this article, unfortunately. On the second page, in Sect. 2.2, “Treatment,” on line 10 it is stated that dexamethasone was administered on days 1–3; however, the administration period should be correctly stated as days 3–5.

The online version of the original article can be found under doi:[10.1007/s12185-011-0884-x](https://doi.org/10.1007/s12185-011-0884-x).

K. Miura (✉) · K. Takei · S. Kobayashi · S. Kiso · Y. Hirabayashi · A. Hojo · H. Kodaira · M. Yagi · D. Kurita · Y. Kobayashi · T. Tanaka · N. Iriyama · Y. Hatta · Y. Kura · T. Yamazaki · U. Sawada · J. Takeuchi
Division of Medicine, Department of Hematology and Rheumatology, Nihon University School of Medicine, 30-1 Oyaguchikamicho, Itabashi Ward, Tokyo, Japan
e-mail: javis@med.nihon-u.ac.jp