Correction to: Influence of Gd-EOB-DTPA on proton density fat fraction using the six-echo Dixon method in 3 Tesla magnetic resonance imaging

Tatsuya Hayashi¹ · Kei Fukuzawa² · Hiroshi Kondo³ · Hiroshi Onodera⁴ · Shuji Toyotaka⁴ · Rie Tojo⁴ · Shimpei Yano⁴ · Masakatsu Tano² · Tosiaki Miyati⁵ · Jun'ichi Kotoku¹ · Takahide Okamoto¹ · Keiko Toyoda³ · Hiroshi Oba³

Published online: 14 March 2018

© Japanese Society of Radiological Technology and Japan Society of Medical Physics 2018

Correction to: Radiol Phys Technol (2017) 10:483–488
<https://doi.org/10.1007/s12194-017-0420-7>

A diameter of glass bottles in phantoms in the above article (2 cm) was incorrect. The correct diameter is 4.5 cm.

The original article can be found online at <https://doi.org/10.1007/s12194-017-0420-7>.

✉ Tatsuya Hayashi
t-hayashi@med.teikyo-u.ac.jp

¹ Graduate School of Medical Technology, Teikyo University, 2-11-1 Kaga, Itabashi-ku, Tokyo 173-8605, Japan

² Department of Radiological Technology, Toranomon Hospital, Tokyo, Japan

³ Department of Radiology, Teikyo University School of Medicine, Tokyo, Japan

⁴ Central of Radiology, Teikyo University Hospital, Tokyo, Japan

⁵ Faculty of Health Sciences, Institute of Medical, Pharmaceutical and Health Sciences, Kanazawa University, Ishikawa, Japan