Erratum to: Taxonomic dissolution of *Sarcostemma* (Apocynaceae: Asclepiadoideae)

Ulrich Meve¹ & Sigrid Liede-Schumann¹

Summary. Due to a production error, an out-of-date (uncorrected) pdf of this paper was uploaded onto Springerlink, and used for the print version of *Kew Bulletin* 67 (4): 751 – 758 ([10.1007/s12225-012-9384-2](https://doi.org/10.1007/s12225-012-9384-2)). Consequently the authority for *Cynanchum viminalis* (L.) Bassi and its subspecies was incorrect. These names are now listed again below with the correct authority.

Cynanchum viminalis (L.) Bassi (1768: 17) subsp. **viminalis**

NOTES. The protologue of *Cynanchum* (*Euphorbia*) *viminalis* (Linnaeus 1753) refers to an illustration of a plant from Egypt (Fefel Taval) as the type and states explicitly ‘Habitat in Africae maritimis’.

Cynanchum viminalis (L.) Bassi subsp. **australe** (R. Br.) Meve & Liede, **comb. nov.**

<http://www.ipni.org/urn:lsid:ipni.org:names:77120428-1>

Sarcostemma australe R. Br., *Prodr.*: 463 (Brown 1810). Type: Australia, South Australia, Bird Island, Petral Bay, Isle St. Francis, 3 Feb. 1802, *R. Brown* (Iter Australiense 2872) (lectotype BM!, designated by Forster (1992)).

Sarcostemma viminalis subsp. *australe* (R. Br.) P. I. Forst., *Austral. Syst. Bot.* 5: 64 (Forster 1992).

Cynanchum viminalis (L.) Bassi subsp. **brunonianum** (Wight & Arn.) Meve & Liede, **comb. nov.**

<http://www.ipni.org/urn:lsid:ipni.org:names:77120429-1>

Sarcostemma brunonianum Wight & Arn., in Wight, *Contr. Bot. India*: 59 (1834). Type: India, [‘Peninsula Ind. Orientalis’], Cody Cally, village of Autcherroccum, *Herb. Wright* 1557 (lectotype G-DC! (No. G00301010), designated here; isolectotypes E, G-DC! (No. G00136527)).

Sarcostemma viminalis (L.) R. Br. subsp. *brunonianum* (Wight & Arn.) P. I. Forst. (Forster 1992: 63).

Cynanchum viminalis (L.) Bassi subsp. **orangeanum** (Liede & Meve) Liede & Meve, **comb. nov.**

<http://www.ipni.org/urn:lsid:ipni.org:names:77120430-1>

Sarcostemma viminalis subsp. *orangeanum* Liede & Meve, *Bot. J. Linn. Soc.* 112: 7 (1993). Type: South Africa, Northern Cape, 11.4 km E Kuruman, *Liede & Meve* 579 (holotype K!; isotype MSUN!).

Cynanchum viminalis (L.) Bassi subsp. **stocksii** (Hook. f.) Meve & Liede, **comb. nov.**

<http://www.ipni.org/urn:lsid:ipni.org:names:77120431-1>

Sarcostemma stocksii Hook. f., *Fl. Brit. India* 4: 27 (Hooker 1883). Type: Pakistan, Sind, *Stocks* 509 (holotype K!). *Sarcostemma viminalis* subsp. *stocksii* (Hook. f.) Ali (1983: 31).

Cynanchum viminalis (L.) Bassi subsp. **thunbergii** (G. Don) Liede & Meve, **comb. nov.**

<http://www.ipni.org/urn:lsid:ipni.org:names:77120432-1>

Sarcostemma thunbergii G. Don, *Gen. Hist.* 4: 156 (1838). Type: South Africa, Cape province, Worcester, Karoo Botanic Garden, *Bayer* 68 (neotype NBG!, designated by Liede & Meve (1993)).

Sarcostemma viminalis subsp. *thunbergii* (G. Don) Liede & Meve (1993: 10).

Cynanchum viminalis (L.) Bassi subsp. **welwitschii** (Hiern) Meve & Liede, **comb. et stat. nov.**

The online version of the original article can be found at <http://dx.doi.org/10.1007/s12225-012-9384-2>

Published online 26 March 2013

¹ Department of Plant Systematics, University of Bayreuth, 95440 Bayreuth, Germany. e-mail: ulrich.meve@uni-bayreuth.de

<http://www.ipni.org/urn:lsid:ipni.org:names:77120433-1>

Sarcostemma wekwitschii Hiern, *Cat. Afr. Pl.* 1: 689 (1898).
Type: Angola, Huilla, base of Morro de Lopollo,
Wekwitsch 4261 (holotype BM!; isotypes BR!, C).

References

- Ali, S. I. (1983). Asclepiadaceae, No. 150. In: E. Nasir & S. I. Ali (eds), *Flora of Pakistan*, pp. 1 – 65. Shamim Printing Press, Karachi.
- Bassi, F. (1768). Novae plantarum species. *Bonomiensi Sci. Inst. Acad. Comment.* 6 (Opusc.): 13 – 20.
- Brown, R. (1810). Asclepiadeae. In: *Prodromus Florae Novae Hollandiae*, pp. 458 – 464. J. Johnson & Co., London.
- Don, G. (1838). Asclepiadeae. In: *General History of the Dichlamydeous Plants* 4, pp. 106 – 166. J. G. & F. Rivington, London.
- Forster, P. I. (1992). A taxonomic revision of *Sarcostemma* R. Br. Subgenus *Sarcostemma* (Asclepiadaceae: Asclepiadeae) in Australia. *Austral. Syst. Bot.* 5: 53 – 70.
- Hiern, W. P. (1898). Asclepiadaceae. In: *Catalogue of the African Plants collected by Dr. Friedrich Wekwitsch, dicotyledons, part 3*, pp. 676 – 698. Printed by order of the Trustees, London.
- Hooker, J. D. (1883). Asclepiadaceae to Amaranthaceae. *The Flora of British India* 4: 1 – 78. L. Reeve & Co., London (Repr. Singh, Dehra Dun).
- Liede, S. & Meve, U. (1993). Towards an understanding of the *Sarcostemma viminalis* (Asclepiadaceae) complex. *Bot. J. Linn. Soc.* 112: 1 – 15.
- Linnaeus, C. (1753). *Euphorbia viminalis*. *Species Plantarum*. P. 452. L. Salvii, Holmiae (Stockholm).
- Wight, R. (1834). Asclepiadae Indicae. In R. Wight (ed.), *Contributions to the Botany of India*, pp. 29 – 67. Parbury, Allen & Co., London.