Erratum to: Before it had a name: Diagnostic characteristics, geographic distribution, and the conservation of *Cupressus tonkinensis* (Cupressaceae)

DAMON P. LITTLE¹, PHILIP THOMAS², HIÊP T. NGUYỄN^{3,4}, AND LỘC K. PHAN^{3,4,5}

¹ Lewis B. and Dorothy Cullman Program for Molecular Systematics, The New York Botanical Garden, Bronx, NY 10458-5126, USA; e-mail: dlittle@nybg.org

² Royal Botanic Garden, Edinburgh, EH3 5LR Scotland, UK

³ Institute of Ecology and Biological Resources, Vietnam Academy of Natural Science and Technology, 18 Hoang Quoc Viet, Cau Giay, Ha Noi, Việt Nam

⁴ Center for Plant Conservation, Vietnam Union of Science and Technology Associations (VUSTA), No. 25/32 Lane 191, L c Long Quan Rd, Nghia Do, Cau Giay, Ha Noi, Việt Nam

⁵ Hanoi University of Science, 334 Nguyen Trai, Thanh Xuan, Ha Noi, Việt Nam

In this article, published in Volume 63 Number 2, pp. 171–196, Fig. 10 was inadvertently omitted. It is reproduced here.

FIG. 10. Branch segments on the plane of the paper are drawn as solid-fill lines, segments pointing out of the plane of the paper away from the viewer are drawn as dashed lines, and segments pointing out of the plane of the paper towards the viewer are drawn as unfilled lines. The antepenultimate segment is labeled. The penultimate segments are labeled at their apex with “p”. The remaining unlabeled segments are all ultimate segments. Penultimate segments A and D bear ultimate segments on only one plane (character 35) whereas segments B and C bear segments on two planes. The frequency of ultimate segment branches on the primary plane (character 12) for segment A=100%, B=60% (6 out of 10), C=62% (5 out of 8), and D=100%. The frequency of apical pointing ultimate segments (character 13) for segment A=50% (5 out of 10), B=50% (2 out of 4), C=60% (3 out of 5), and D=66% (2 out of 3).