Foreword

Wm. Wayt Thomas^{1,2}

¹ The New York Botanical Garden, New York, USA

² Author for correspondence; e-mail: wthomas@nybg.org

Published online: 24 January 2009

© The New York Botanical Garden 2009

The International Botanical Congress, held in Vienna, Austria in July of 2005, included two symposia on the family Cyperaceae. The cosmopolitan Cyperaceae are the world's seventh largest plant family, with about 105 genera and perhaps 5,500 species. The largest genus, *Carex* (ca. 2,000 spp.), is remarkable for its exceptional diversity, varied ecology, unusual cytology and diverse phytogeography. Recently, through molecular studies, many conflicting hypotheses about phylogenetic relationships in Cyperaceae have been resolved, as have relationships within some of the major clades and larger genera. These exciting advances have helped solve some of the most difficult problems in Cyperaceae, including assessing morphological and anatomical convergence due to the highly reduced morphology and difficulties in interpreting homology in the evolution of spikelets and flowers.

The first symposium, "Sedges have edges—Phylogenetic relationships and evolution," organized by Anton Reznicek, comprised seven talks on the evolution and phylogenetic relationships of the whole family or non-caricoid groups in the family. The second symposium, "Evolution in the genus *Carex* and tribe Cariceae: New ideas from new phylogenies," organized by Matthias Hendrichs, Julian R. Starr, and Bruce A. Ford, comprised six talks and focused on the huge genus *Carex*. These 13 talks have resulted in eight papers presented in this issue of Botanical Review. In editing this volume, I was assisted by David Simpson, Anton Reznicek and Julian Starr.