Ying Huang
Department of Respiratory Medicine
Children's Hospital of Chongqing Medical University
No.136, Zhongshan 2nd Road, Yuzhong District
Chongqing 400014, China
Email: huangying62@126.com

References

- Basaran S, Guler-Uysal F, Ergen N, Seydaoglu G, Bingol-Karakoc G, Ufuk Altintas D. Effects of physical exercise on quality of life, exercise capacity and pulmonary function in Children with asthma. *J Rehabil Med* 2006;38:130-135.
- Carlsen KH, Anderson SD, Bjermer L, Bonini S, Brusasco V, Canonica W, et al. Treatment of exercise-induced asthma, respiratory and allergic disorders in sports and the relationship to doping: Part II of the report from the Joint Task Force of European Respiratory Society(ERS) and European Academy of Allergy and Clinical Immunology (EAACI) in cooperation with GA(2)LEN. *Allergy* 2008;63:492-505.
- Lucas SR, Platts-Mills TA. Physical activity and exercise in asthma: relevance to etiology and treatment. *J Allergy Clin Immunol* 2005;115:928-934.
- Cheng BL, Huang Y, Shu C, Lou XL, Fu Z, Zhao J. A cross-sectional survey of participation of asthmatic children in physical activity. *World J Pediatr* 2010;6:238-243.

doi: 10.1007/s12519-011-0272-x

Controlled insufflation of air for diagnosis and treatment of acute intestinal intussusception

To the Editor

I have read with great interest the article by Kong et al^[1] where the authors refer to a successful pneumatic reduction of acute intestinal intussusception in 17/23 cases (74% [95%CI 56-92]). The article makes no mention of whether the machine of air/barium enema utilized for the intussusception's reduction is connected to the patient by means of an occlusive catheter.

The purpose of this letter is to provide a little historical contribution regarding the first description of insufflation of air for diagnosis and treatment of intestinal intussusception.

A new method termed "controlled insufflation" of air for diagnosis and treatment of acute intestinal intussusception was developed by Fiorito et al in 1953,^[2] and the authors described an original apparatus called a baroscope, connected to the patient's anus by means of an occlusive catheter (Fig.), which under fluoroscopic control allowed a successful reduction of intestinal intussusception in 81/86 cases (94% [95%CI 91-99]).^[3] In contrast to Kong et al's patients, Fiorito et al's patients were all between 3 and 18 months of age.

According to the authors, the "controlled insufflation"

(Reproduced with permission from the American Academy of Pediatrics)
Fig. The baroscope (schematic). A: mercury manometer with intraluminal wires at the bottom and at 60 and 130 mm; B: relay that operates a valve; C: interruptor to select highest pressures of 60 or 130 mmHg; D: valve for releasing gas pressure.

had the following advantages. A) In diagnosis: 1) Simplicity, without the troubles and interruptions during fluoroscopic examination commonly seen in the barium-enema technique. 2) Fast diffusion of the air so that it advances promptly to the intussusception. 3) Clear-cut visualization of intussusception in all cases as it is seen surrounded by gas. 4) No contraindications. It can be used in the diagnosis of advanced cases with the highest pressure being 60 mmHg [6 Kpa]. 5) No interference with normal bowel physiology. 6) In all cases air passes the ileo-cecal valve. Ileo-cecal forms can therefore be diagnosed. B) In treatment: 1) The complete filling of the small bowel with air confirms the full reduction of the intussusception (Working within a maximum pressure of 130 mmHg [13 kPa]). 2) In our experience the results are much better than those obtained with the barium-enema technique.

Eduardo Cuestas
Department of Pediatrics and Neonatology
Hospital Privado
Córdoba, Argentina
Email: ecuestas@hospitalprivadosa.com.ar

References

- Kong FT, Liu WY, Tang YM, Zhong L, Wang XJ, Yang G, et al. Intussusception in infants younger than 3 months: a single center experience. *World J Pediatr* 2010;6:55-59.
- Fiorito ES, Recalde Cuestas LA. The "controled insufflation" (baro-fluoroscopically), method of choice for the diagnosis and treatment of intussusception in infants. *Rev Soc Pediatr Litoral* 1953;18:3.
- Fiorito ES, Recalde Cuestas LA. Diagnosis and treatment of acute intestinal intussusception with controlled insufflation of air. *Pediatrics* 1959;24:241-244.

doi: 10.1007/s12519-011-0273-9