Erratum to: Effect of different storage conditions on the mycotoxin contamination of *Fusarium culmorum*-infected and non-infected wheat straw

Dirk Rohweder · Hana Valenta · Sarah Sondermann ·
Margit Schollenberger · Winfried Drochner ·
Guenter Pahlow · Susanne Döll · Sven Dänicke

Published online: 30 March 2011
© Society for Mycotoxin Research and Springer 2011

Erratum to: Mycotoxin Research
DOI 10.1007/s12550-011-0087-6

The publisher regrets that Table 1 and Table 2 were displayed incorrectly in the published article. The correct tables are shown below.

The following error appeared in Table 1:

The marks for alteration differs significant from zero were placed incorrect.

The following error appeared in Table 2:

In line “3-Ac-DON”, column “CON 86”, “CON 82” and “CON 78” the superscript “a” must be placed behind 0.4, 0.6 and 0.4, respectively.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s12550-011-0087-6>.

D. Rohweder · H. Valenta · S. Döll (✉) · S. Dänicke
Institute of Animal Nutrition, Friedrich-Loeffler-Institute (FLI)
Federal Research, Institute for Animal Health,
Bundesallee 50,
38116 Braunschweig, Germany
e-mail: susanne.doell@fli.bund.de
URL: www.fli.bund.de

S. Sondermann · M. Schollenberger · W. Drochner
Institute of Animal Nutrition, University of Hohenheim,
Stuttgart, Germany

G. Pahlow
Institute for Crop and Soil Science, Julius-Kühn-Institute, Federal
Research Centre for Cultivated Plants,
Braunschweig, Germany

Table 1 FSE initial dry matter and mycotoxin concentrations of wheat straw and the identically cultivated *Fusarium culmorum*-inoculated variant, and the alteration per day during 32 weeks of storage in a barn and outdoors ($\mu\text{g/kg}$, estimated intercepts and alterations per day of the mixed model, $n=3$ at each of the six sampling times)

Straw storage	CON			FUS		
	Initial	Barn	Outdoor	Initial	Barn	Outdoor
Dry matter (%)	81.8	0.02 ^{a*}	−0.28 ^{b*}	81.2	<0.01 ^a	−0.19 ^{b*}
DON	526 ^A	−0.4 ^a	<0.1 ^a	28 713 ^B	−17.8 ^a	−69.2 ^{b*}
15-Ac-DON	41 ^A	−0.1	−0.2	69 ^B	−0.3	−0.4
3-Ac-DON	37 ^A	−0.1 ^a	<0.1 ^a	3 371 ^B	0.6 ^a	−16.3 ^{b*}
ZEA	87 ^A	0.4 ^{a*}	2.9 ^{b*}	585 ^B	0.1 ^a	6.2 ^{c*}

^{AB} Different superscripts describe significant effect in mycotoxin concentration of inoculation, ^{abc} Different superscripts describe significant differences in alteration between the variants, * Alteration differs significantly from zero

Table 2 Initial mycotoxin concentrations of wheat straw and its identical cultivated, with *Fusarium culmorum* inoculated variant and The alteration per day during 16 weeks of storage at a target dry matter (DM) of 86%, 82% and 78% of the laboratory scale experiment ($\mu\text{g/kg}$, estimated intercepts and alterations per day of the mixed model, $n=1$ at each of the five sampling times)

Straw storage DM (%)	CON				FUS			
	Initial	86	82	78	Initial	86	82	78
Dry matter (%)		0.02 ^{b*}	0.03 ^{b*}	0.05 ^{c*}		0.02 ^{b*}	0.02 ^{ab*}	<0.01 ^a
DON	548 ^A	−0.1	0.6	−0.9	30 323 ^B	−65.7	10.5	−68.6
15-Ac-DON	46 ^A	<0.1	−0.2	−0.3	67 ^B	−0.3	<0.1	−0.4
3-Ac-DON	20 ^A	0.4 ^a	0.6 ^a	0.4 ^a	2705 ^B	−8.0 ^b	−14.6 ^b	−27.8 ^{b*}
ZEA	250 ^A	−1.2	−1.0	0.4	813 ^B	−0.2	−3.1	−0.6

^{AB} Different superscripts describe significant effect in mycotoxin concentration of inoculation, ^{abc} Different superscripts describe significant differences in alteration between the variants, * Alteration differs significantly from zero