Growth, Feeding and Ecological Roles of the Mixotrophic and Heterotrophic Dinoflagellates in Marine Planktonic Food Webs

Hae Jin Jeong^{1*}, Yeong Du Yoo¹, Jae Seong Kim², Kyeong Ah Seong³, Nam Seon Kang¹, and Tae Hoon Kim⁴

¹*School of Earth and Environmental Sciences, College of Natural Sciences, Seoul National University, Seoul 151-747, Korea*

²*Red Tide Research Center, Kunsan National University, Kunsan 573-701, Korea*

³*Saemankeum Environmental Research Center, Kunsan National University, Kunsan 573-701, Korea*

⁴*Research Institute of Oceanography, College of Natural Sciences, Seoul National University, Seoul 151-747, Korea*

Received 15 March 2010; Revised 28 April 2010; Accepted 2 May 2010

© KSO, KORDI and Springer 2010

Abstract – Planktonic mixotrophic and heterotrophic dinoflagellates are ubiquitous protists and often abundant in marine environments. Recently many phototrophic dinoflagellate species have been revealed to be mixotrophic organisms and also it is suggested that most dinoflagellates may be mixotrophic or heterotrophic protists. The mixotrophic and heterotrophic dinoflagellates are able to feed on diverse prey items including bacteria, picoeukaryotes, nanoflagellates, diatoms, other dinoflagellates, heterotrophic protists, and metazoans due to their diverse feeding mechanisms. In turn they are ingested by many kinds of predators. Thus, the roles of the dinoflagellates in marine planktonic food webs are very diverse. The present paper reviewed the kind of prey which mixotrophic and heterotrophic dinoflagellates are able to feed on, feeding mechanisms, growth and ingestion rates of dinoflagellates, grazing impact by dinoflagellate predators on natural prey populations, predators on dinoflagellates, and red tides dominated by dinoflagellates. Based on this information, we suggested a new marine planktonic food web focusing on mixotrophic and heterotrophic dinoflagellates and provided an insight on the roles of dinoflagellates in the food web.

Key words – grazing, harmful algal bloom, ingestion, predation, predator, prey, protist, red tide

Marine dinoflagellates are ubiquitous and some genera are cosmopolitan at present (Lessard 1984; Jeong 1999; Rublee et al. 2004). Also, they have been easily observed in the fossil record, reflecting their ubiquitous presence in the past (e.g. Radi et al. 2007). They often dominate the plankton

assemblages and sometimes form red tides or harmful blooms (e.g. Jeong 1999). They have very diverse morphology (size, shape etc.) and genetics which are two most critical keys for their classification (e.g. Daugbjerg et al. 2000). They have 3 different major trophic modes (i.e. autotrophic, mixotrophic, and heterotrophic) and the mixotrophic and heterotrophic dinoflagellates have diverse feeding mechanisms. Also, the mixotrophic and heterotrophic dinoflagellates are known to feed on diverse types and sizes of prey. In turn, dinoflagellates are excellent prey for mixotrophic and heterotrophic protists and metazoans. Therefore, dinoflagellates play diverse roles in marine planktonic food webs.

In the last 2 decades, many phototrophic dinoflagellates which had previously been thought to be exclusively autotrophic dinoflagellates have been revealed to be mixotrophic dinoflagellates (i.e. capable of both photosynthesis and ingesting prey) (Jacobson and Anderson 1996; Stoecker 1999; Jeong et al. 2005c, 2010; Seong et al. 2006). The kind of prey which these mixotrophic dinoflagellates are able to feed on is very diverse including bacteria, other dinoflagellates, and heterotrophic protists. In particular, recently, feeding by dinoflagellates on a single bacterial cell has been confirmed (Seong et al. 2006; Jeong et al. 2008). Several dinoflagellate genera such as *Karenia*, *Karlodinium*, *Lepidodinium* have been discovered to have fucoxanthin or chlorophyll *b* as accessory pigments, not peridinin which had been thought to be the common carotenoid accessory pigment of phototrophic dinoflagellates (Watanabe et al. 1990; Garces et al. 2006). Thus, it is believed that these

*Corresponding author. E-mail: hjjeong@snu.ac.kr

pigments inside these dinoflagellates were originated from prey plastids through feeding (Bhattacharya et al. 2003). Feeding is likely to be one of the most important driving forces for the evolution of dinoflagellates. When combining the nutrition of modern dinoflagellates with the presence of fossil dinoflagellates in geological records, dinoflagellates have been ubiquitous for a long time (> 100 MY) due to their diverse trophic modes and prey items. When their common prey became rare or disappeared in geological time scales, dinoflagellates might change their trophic modes and/or evolve their morphology and digestive enzymes to feed on the new prey items. Therefore, investigating predator-prey relationships among dinoflagellates and between dinoflagellates and diverse plankton and exploring roles of dinoflagellates in food webs may be critical steps in understanding the nature of the ecosystems at present and in the past and also evolution of dinoflagellates.

In this paper, we reviewed the kind of prey which mixotrophic and heterotrophic dinoflagellates are able to feed on, feeding mechanisms, growth and ingestion rates of dinoflagellates, grazing impact, and predators. In addition, we suggested a possible mechanism of the outbreak of red tides or harmful blooms dominated by mixotrophic dinoflagellates in offshore and/or oceanic waters where the nutrient concentrations are low. We also suggested new marine planktonic food webs focussing on dinoflagellates based on the literature and also addressed the roles of dinoflagellates in these food webs. In the section on the kind of prey and feeding mechanisms, i) whether dinoflagellates are able to feed on most types and sizes of prey, even very small or large prey, ii) if so, how dinoflagellates capture and ingest these diverse prey, and iii) in an evolutionary sense, whether morphological diversity of dinoflagellates could be related to their feeding mechanisms were explored. In the section on the growth and ingestion rates and grazing impact by dinoflagellates on diverse prey, iv) when these predators' sizes are similar, whether the maximum growth rates of autotrophic, mixotrophic, and heterotrophic dinoflagellates are also similar or not, v) whether the maximum ingestion rates of mixotrophic dinoflagellates are comparable to those of heterotrophic dinoflagellates or not, and vi) what are the important factors affecting the maximum growth and ingestion rates were investigated. In the section on predators, vii) whether maximum ingestion rates of major predators (other dinoflagellates, ciliates, and copepods) on mixotrophic dinoflagellates are similar to

those of heterotrophic dinoflagellates and viii) whether the mortality rate of dinoflagellate prey due to predation by heterotrophic protistan and metazoan predators is critical in controlling dinoflagellate red tides were addressed.

This review paper provides an insight on how dinoflagellates come to be ubiquitous and often dominate the plankton assemblages at present and in the past, how dinoflagellates have diverse trophic modes, feeding mechanisms, and diverse prey items, and to which direction dinoflagellates have evolved their morphology and/or enzymes.

1. Trophic Modes

Marine dinoflagellates have 3 major trophic modes (i.e. autotrophy, mixotrophy, and heterotrophy) (Lessard and Swift 1985; Burkholder et al. 1992; Steidinger et al. 1996; Sherr and Sherr 2002; Mason et al. 2007). Phototrophic dinoflagellates (autotrophic or mixotrophic dinoflagellates) have been thought to be one of the most important phytoplankton groups for a long time. Therefore, there have been many studies on the ecology and physiology of the phototrophic dinoflagellates assuming that they were exclusively autotrophic dinoflagellates (e.g. Smayda 1997). They have often formed huge red tides which have sometimes caused large-scale mortalities of fin-fish and shellfish and thus great losses to the aquaculture and tourist industries of many countries (ECOHAB 1995; Azanza et al. 2005). The mechanisms of the outbreak, persistence, and decline of the red tides dominated by phototrophic dinoflagellates have also been studied based on the assumption that they were exclusively autotrophic dinoflagellates. However, many phototrophic dinoflagellates have recently been revealed to be mixotrophic dinoflagellates (MTDs) and thus the study on phagotrophy of phototrophic dinoflagellates is rapidly increasing (Larsen 1988; Jacobson and Anderson 1994; Granéli et al. 1997; Jeong et al. 1997; Adolf et al. 2006; Burkholder et al. 2008). In addition, recently some newly described phototrophic dinoflagellate species have been revealed to be mixotrophic dinoflagellates (e.g. Kang et al. 2010; Yoo et al. 2010). Thus, mixotrophy by the causative phototrophic dinoflagellate species should be considered when the mechanisms of the outbreak, persistence, and decline of red tides are being explored.

Heterotrophic dinoflagellates (HTDs) have been known to have diverse feeding mechanisms and feed on diverse prey (e.g. Hansen and Calado 1999). Recently, their feeding

mechanisms and prey items have been newly discovered (e.g. Jeong et al. 2008). Thus, difference in maximum growth rates among similar-sized autotrophic, mixotrophic, and heterotrophic dinoflagellates and differences in the kind of prey, feeding mechanisms, and maximum ingestion rates between MTDs and HTDs may give a clue to understanding the evolution in the trophic modes of dinoflagellates in a geological time scale.

2. The Kind of Prey which Dinoflagellates are Able to Feed on

Mixotrophic and heterotrophic dinoflagellates usually co-occur with diverse types and sizes of plankton in marine environments. Thus, the following questions arise; (1) Are dinoflagellates able to feed on most plankton taxa? Even toxic prey species? (2) Is the kind of prey which MTDs are able to feed on different from that of HTDs? (3) Are dinoflagellates able to feed on plankton prey with a wide range of size, even very small (i.e. bacteria) or large prey

(larval fish)? If so, how do dinoflagellates capture and ingest these very small or large prey? (i.e. what are their feeding mechanisms?) (4) Is the lower or upper prey size limit affected by the predator size?.

2a. Prey taxa

Both MTDs and HTDs have been known to feed on diverse taxa including phototrophic (Strom 1991; Jeong et al. 2005b) and heterotrophic bacteria (Jeong et al. 2008), pico-eukaryotes (Lee 2006), cryptophytes (Li et al. 2000), haptophytes (Goldman et al. 1989; Berge et al. 2008), chlorophytes (Strom and Buskey 1993), prasiophytes (Berge et al. 2008), raphidophytes (Nakamura et al. 1995), diatoms (Jacobson and Anderson 1986; Menden-Deuer et al. 2005; Yoo et al. 2009), other dinoflagellates (Adolf et al. 2007; Tillmann 2004), heterotrophic nanoflagellates (Jeong et al. 2007b), and ciliates (Hansen 1991; Bockstahler and Coats 1993a, 1993b) (Table 1). However, some HTDs have been reported to be able to feed on the blood, flesh, eggs, early naupliar stages, and adult forms of metazoans, while MTDs

Table 1. The kind of prey taxa which mixotrophic (MTD) and heterotrophic dinoflagellate (HTD) are able to feed on.

(DE: Direct engulfment feeders, PAL: Pallium feeders, PED: Peduncle feeders, Numbers: feeding occurred and numbers are references, -: tested, but not found yet, Blank: not tested)

Prey / Predator	MTD	HTD	DE	PAL	PED
heterotrophic bacteria	(1)	(2, 3)	(3)	(3)	(4)
autotrophic bacteria	(5)	(6)	(6)		(4)
pico-eukaryotes	(7)				
Cryptophytes	(8)	(9, 10)	(11)	(10)	(4)
Haptophyta (=Prymnesiophyta)	(12)	(13, 10)	(13)	(10)	(12)
Chlorophytes (Dunaliella)		(13, 10)	(13)	(10)	
Prasiophytes	(12)	(10)	(14)	(15)	
Euglenophytes					
Raphidophytes	(12)	(15, 16)	(15)	(16)	(16)
Bacillariophytes	(17)	(18, 19)	(18)	(20)	(19)
Dinoflagellates	(21)	(22, 16)	(22)	(13)	(16)
Heterotrophic nanoflagellates	(23)	(24)	(24)		(24)
Ciliates	(25, 26)	(4)	(27)		(26)
Eggs of metazoans	-	(28, 29)		(28)	(29)
Naupliar stages of metazoans	-	(28, 29)		(21)	(29)
Bloods of metazoans	-	(29)			(29)
Flesh of metazoans	-	(29)			(29)

Note. (1): Seong et al. (2006), Nygaard and Tobiesen (1993), (2): Lessard and Swift (1985), Burkholder and Glasgow (1995, 1997), (3): Jeong et al. (2008), (4): Burkholder and Glasgow (1995), (5): Jeong et al. (2005b), (6): Strom (1991), Glasgow et al. (1998); (7): Lee (2006), (8): Li et al. (2000), Jakobsen et al. (2000), Stoecker et al. (1997), (9): Feinstein et al. (2002), (10): Strom and Buskey (1993), (11): Jakobson and Hansen (1997), (12): Berge et al. (2008), (13): Goldman et al. (1989), (14): Hansen et al. (1996), (15): Nakamura et al. (1992), (16): Jeong et al. (2006, 2007a), (17): Legrand et al. (1998), Bockstahler and Coats (1993b), (18): Naustvoll LJ (1998), Nakamura et al. (1995), Hansen PJ (1992), Strom and Buskey (1993), (19): Parrow et al. (2001), (20): Menden-Deuer et al. (2005), (21): Skovgaard (1996a), Hansen and Nielsen (1997), (22): Nakamura et al. (1995), (23): Jeong et al. (unpubl.data), (24): Jeong et al. (2007b), (25): Smalley et al. (1999), Bockstahler and Coats (1993a), Hansen PJ (1991), (26): Park et al. (2006), (27): Bursa AS (1961), (28): Jeong (1994b), (29): Burkholder and Glasgow (1997)

Fig. 1. The size (Equivalent Spherical Diameters, ESD, μm) of the largest prey which engulfment feeding mixotrophic (A. open circles, MTD) and heterotrophic dinoflagellates (B. closed circles, HTD) are able to feed on as a function of the predator size. The equation of the regression was $\text{ESD of the MTD predator } (\mu\text{m}) = 0.587 (\text{ESD of prey}) + 1.67$, $r^2 = 0.687$ ($p < 0.005$)

(At: *Alexandrium tamarense*, As: *Akashiwo sanguinea*, Cp: *Cochlodinium polykrioides*, Fm: *Fragilidium cf. mexicanum*, Gc: *Gymnodinium catenatum*, Gi: *Gymnodinium impudicum*, Ht: *Heterocapsa triquetra*, Lp: *Lingulodinium polyedrum*, Pd: *Prorocentrum donghaiense*, Pmc: *P. micans*, Pm: *P. minimum*, Pt: *P. triestinum*, St: *Scrippsiella trochoidea*, Oxy: *Oxyrrhis marina*, Gdo: *Gyrodinium dominans*, Gsp: *Gyrodinium spirale*, Poly: *Polykrikos kofoidii*)

were not (e.g. Jeong 1994b, 2007a). Among HTDs, the pallium (feeding veil) and peduncle (feeding tube) feeders were known to be able to feed on these animal prey items, while the engulfment feeders were not (see feeding mechanism section for details). For the engulfment feeding HTDs, eggs, early naupliar stages, and adult forms of metazoans are too large to engulf. There are some MTD species that feed on prey using a peduncle (Park et al. 2006; Berge et al. 2008; Jeong et al. 2010; Yoo et al. 2010). Therefore, it is worthwhile to discover any MTD that is able

to feed on these animal prey items.

Even though several MTDs and HTDs have co-occurred with euglenophytes during euglenophyte blooms (our unpublished data), there has been no study on feeding by MTDs and HTDs on euglenophytes yet. It is worthwhile to explore this topic to understand the dynamics of euglenophytes and roles of dinoflagellates in these dynamics. Also, there has been no study on the feeding by MTDs on chlorophytes yet. Further study on whether MTDs and HTDs are able to feed on prey not tested yet is necessary (Table 1).

2b. Toxic prey

Some MTDs and HTDs were able to feed on toxic algae; the MTDs *Lingulodinium polyedrum* and *Akashiwo sanguinea* were able to feed on the toxic dinoflagellate *Alexandrium tamarense* (Jeong et al. 2005c); also, the HTDs *Polykrikos kofoidii* fed on the toxic MTD *Gymnodinium catenatum* and detoxified the toxin produced by the prey (Jeong et al. 2003b). These dinoflagellates may have enzyme-detoxifying phytochemicals and it is worthwhile to explore the presence and nature of these enzymes. Dinoflagellates may evolve to feed on toxic algae which might evolve to have toxins for anti-predation.

2c. Size of prey and predator

The size range of plankton co-occurring with dinoflagellates are very wide, from very small bacteria to large larval fish. Many think that a single bacterium cell is too small to be eaten by dinoflagellates. Some studies reported signs of bacteria inside some MTD and HTD predators using bacteria prelabelled with tritiated-thymidine (Lessard and Swift 1985; Nygaard and Tobiesen 1993). However, these studies neither reported the feeding behavior of HTDs on bacteria nor provided transmission electron microscopy (TEM) pictures showing ingested bacterial cells or bacteria sized beads in the food vacuoles inside the protoplasm of predator cells. Therefore, for a long time, some argued that the signs of bacteria inside the dinoflagellate predators might be derived from detritus to which the bacteria are attached or labelled bacteria stuck to the predator surface. Recently, Seong et al. (2006) and Jeong et al. (2005b, 2008), using confocal microscopy, TEM, and video microscopy (see details in the feeding mechanism section), have clearly showed that MTDs and HTDs were able to feed on a single bacterium cell. Therefore, the lower prey size limits of MTDs and HTDs goes down to pico-size. In many previous studies, the lower prey size limits of MTDs and HTDs had

Table 2. Feeding occurrence by each mixotrophic dinoflagellate predator on diverse prey items which were identified in the taxonomic level listed below

(+) : A dinoflagellate predator was observed to feed on prey cells, (–): the dinoflagellate predator was observed not to feed on prey cells, No (+) or (–) means not tested, ESD: Equivalent Spherical Diameters (μm), HB: Heterotrophic bacteria, CB: Cyanobacteria, HP: Haptophyte, CR: Cryptophyte, Ac: *Amphidinium carterae*, DA: a small diatom, Ha: *Heterosigma akashiwo*, Pm: *Prorocentrum minimum*, Ht: *Heterocapsa triquetra*, Cp: *Cochlodinium polykrikoides*, Pc: *P. micans*, At: *Alexandrium tamarense*, As: *Akashiwo sanguinea*, Lp: *Lingulodinium polyedrum*, Ce: *Ceratium* spp., HD: Heterotrophic dinoflagellates, CL: Ciliates)

Predator / Prey	ESD	HB	CB	HP	CR	Ac	DA	Ha	Pm	Ht	Cp	Pc	At	As	Lp	Ce	HD	CL	Ref.
<i>Heterocapsa rotundata</i>	5.8	+	+				+												(6,11,15)
<i>Amphidinium carterae</i>	6.6						+												(15)
<i>Prorocentrum minimum</i>	12.1	+	+	+	+	+	+	+		–	–	–	–	–	–				(6,7,9,11,13,15)
<i>Paragymnodinium shiwhaense</i>	12.4			+	+	+	–	+	–	–	–	–	–	–	–				(18)
<i>Gyrodinium galatheanum</i>	12.5				+														(9)
<i>Prorocentrum triestinum</i>	12.6	+		+	+	+	+	+	+	–	–	–	–	–	–				(6,7,11,15)
<i>Karlodinium armiger</i>	13.1			+	+		+	+	+	+		+		+			+		(17)
<i>Prorocentrum donghaiense</i>	13.3	+	+	+	+	+	+	+	+	–	–	–	–	–	–				(6,7,11,15)
<i>Heterocapsa triquetra</i>	15.0	+	+	+	+	+	+	+	+		–	–	–	–	–				(6,7,11,15)
<i>Alexandrium minutum</i>	16.7		+																(6)
<i>Gymnodinium impudicum</i>	17.8	+	+	+	+	+	+	+	–	–	–	–	–	–	–				(6,7,11,15)
<i>Gymnodinium aureolum</i>	19.4	+	+	+	+	+	–	+	–	–	–	–	–	–	–				(16)
<i>Karenia brevis</i>	20.3		+																(6)
<i>Scrippsiella trochoidea</i>	22.8		+	+	+	+	+	+	+	–	–	–	–	–	–				(6,7,11,15)
<i>Cochlodinium polykrikoides</i>	25.9	+	+	+	+	+	+	+	–	–		–	–	–	–				(6,7,11,15)
<i>Prorocentrum micans</i>	26.6	+	+	+	+	+	+	+	+	+	–		–	–	–				(6,7,11,15)
<i>Alexandrium tamarense</i>	28.1	+	+	+	+	+	+	+	+	–	–	–		–	–				(6,7,11,15)
<i>Ceratium furca</i>	29.0																	+	(3)
<i>Gyrodinium pavillardii</i>	29.5																	+	(1)
<i>Akashiwo sanguinea</i>	30.8	+	+	+	+	+	+	+	+	+	–	–	+		–			+	(2,3,6,7,11,15)
<i>Gonyaulax polygramma</i>	32.5	+	+	+	+	+	+	+	+	+	–	–	–	–	–				(6,8,11,15)
<i>Alexandrium catenella</i>	32.6	+	+																(6,11)
<i>Gymnodinium catenatum</i>	33.9	+	+	+	+	+	+	+	+	–	–	–	–	–	–				(6,7,11,15)
<i>Gonyaulax spinifera</i>	35.0		+																(6)
<i>Lingulodinium polyedrum</i>	38.2	+	+	+	+	+	+	+	+	+	–	–	+	–					(6,7,11,15)
<i>Gyrodinium uncatenum</i>	39.0																	+	(3)
<i>Gyrodinium instriatum</i>	43.2																	+	(14)
<i>Fragilidium subglobosum</i>	50.0															+			(12)
<i>Fragilidium mexicanum</i>	55.6					–			+		–	+		+	+		+		(4)

Note. (1): Biecheler (1952), (2): Bockstahler and Coats (1993a), (3): Bockstahler and Coats (1993b), (4): Jeong et al. (1999a), (5): Jeong et al. (2004c), (6): Jeong et al. (2005b), (7): Jeong et al. (2005c), (8): Jeong et al. (2005d), (9): Li et al. (1996), (10): Nygaard and Tobiesen (1993), (11): Seong et al. (2006), (12): Skovgaard (1996a), (13): Stoecker et al. (1997), (14): Uchida et al. (1997), (15): Yoo et al. (2009), (16): Jeong et al. (2010), (17): Berge et al. (2008), (18): Yoo et al. (2010)

been reported to be 3–4 μm (Hansen 1992; Jakobsen and Hansen 1997; Hansen and Calado 1999). These studies tested algal prey with the assumption that dinoflagellates might feed on prey using raptorial feeding (see feeding mechanism section). However, some small HTDs fed on a single bacterium cell using feeding mechanisms (filter/interception feeding) different from those used for large algal prey (Jeong et al. 2008). Discovery of new feeding mechanisms of dinoflagellates might give a clue for understanding

feeding by dinoflagellates on a single bacterium cell.

For the engulfment feeding dinoflagellates, the size (Equivalent Spherical Diameter, ESD) of the largest prey which MTDs are able to feed on was significantly positively correlated with the ESD's of the predators ($p < 0.005$, linear regression ANOVA; Fig. 1A). These MTDs engulf large prey cells through the sulcus which is in general proportional to the body size (Jeong et al. 2005c). Thus, in general the upper size limit of prey for the engulfment feeding MTDs is

proportional to the predator size (Table 2). However, the ESD of the largest prey which the engulfment feeding HTDs are able to feed on was not significantly positively correlated with the ESD's of the predators ($p>0.1$; Fig. 1B). When the sizes are similar, the upper prey size limits of the HTDs were larger than those of MTDs and thus the ratios of predator to prey size for the HTDs were clearly lower than those of the MTDs (Fig. 1A and B). These HTDs may have more flexible sulcus than the MTDs. Also, microtubules involved in opening and closing sulcus of HTDs may be more well developed than MTDs. In addition, force involved in pulling prey cells into the protoplasm of HTDs may be stronger than those of MTDs. Further study on this topic is necessary.

The pallium feeding HTDs or peduncle feeding MTDs and HTDs are able to feed on prey larger than themselves (Gaines and Taylor 1984; Jacobson 1987; Jeong et al. 2007a; Berge et al. 2008; Yoo et al. 2010) (see next subsection for details). Therefore, their upper prey size limits are larger than those of the engulfment feeding MTDs and HTDs.

3. Feeding Mechanisms and Evolution in Morphology

Dinoflagellates are able to feed on diverse types and sizes of prey as mentioned above. Thus, several questions arise; (1) what are feeding mechanisms of dinoflagellates? Are they diverse? (2) Is there any dinoflagellate having 2 or more feeding mechanisms? (3) What makes feeding mechanisms of dinoflagellates be diverse? (4) Is there any relationship between the morphology and feeding mechanisms of dinoflagellates?

3a. Types

In general, there are 3 major feeding mechanisms of free-living protists (Boenigk and Arndt 2000; Fenchel 1987; Sleight 1989): raptorial feeding, filter/interception feeding, and diffusion feeding. In raptorial feeding, predators search, capture, ingest, and digest prey cells, while in filter/interception feeding, predators generate feeding currents and filter or intercept prey cells in the feeding current. In diffusion feeding, predators wait until prey cells arrive in the feeding part of the predators and ingest the prey cells. For a long time, many studies reported that planktonic dinoflagellates fed on prey cells using raptorial feeding because these studies focused on nano-sized or larger prey.

Fig. 2. A diagram of possible evolution in the morphology of dinoflagellates ingesting a prey cell through the sulcus. Dinoflagellates with a large displacement of the cingulum and/or extension of the sulcus (Dinoflagellate B) may have an advantage in engulfing larger prey cells over ones with smaller or no displacement of the cingulum (Dinoflagellate A)

3b. The number and size of the mouth

There are 3 major feeding mechanisms in the raptorial feeding; direct engulfment feeding, pallium feeding, and peduncle feeding. Direct engulfment feeders ingest prey cells through openings (like mouth) of the predator body. Thus, several questions arise; (1) where is a mouth? (2) Is there only one mouth in a dinoflagellate? (3) Is it located in a similar area among dinoflagellates? Most dinoflagellates are known to engulf a prey cell through the sulcus (Skovgaard 1996a; Jeong et al. 2005c). However, recently it was found that the MTDs *Gonyaulax polygramma* and *Scrippsiella* spp. engulfed a small prey through the apical horn, whereas they engulfed a larger prey through the sulcus (i.e. 2 mouths; Jeong et al. 2005c, 2005d). Surprisingly, *Prorocentrum* spp. engulfs prey cells through sutures on the sides of several parts of their bodies (i.e. multiple mouths; Jeong et al. 2005c). Sometime a *P. micans* cell engulfed 3 cyrptophyte prey cells at 3 different locations simultaneously. Therefore, the sulcus, apical horn, and suture are mouths of dinoflagellates and some dinoflagellates have 2 or more mouths. Considering the locations through which the dinoflagellates engulf, the dinoflagellates are close to amoebae that engulf prey cells through several parts of the body rather than common ciliates that engulf prey cells through the cytostome.

The size of the opening (sulcus, apical horn, suture etc.) is likely to be one of the critical factors determining predator and prey in the feeding battle among the engulfment feeding dinoflagellates (Jeong et al. 2005c). For dinoflagellates ingesting a prey cell through the sulcus, having a large displacement of the cingulum and/or extension of the sulcus may provide

an advantage in engulfing larger prey cells over dinoflagellates with smaller or no displacement of the cingulum (Fig. 2). The engulfment feeding dinoflagellates may evolve to have larger openings to engulf larger prey cells.

Dinoflagellates may develop pallium or peduncle feeding mechanisms to feed on prey larger than the predator themselves, as mentioned above. Some pallium and peduncle feeding dinoflagellates are able to feed on prey 10 times larger than themselves (e.g. Jacobson 1987). Unlike most engulfment feeding dinoflagellates, several pallium and peduncle feeding dinoflagellate cells are able to attack a prey cell simultaneously (Jeong 1994b). When the range of size and types of prey are considered, the pallium or peduncle feeding dinoflagellates may be more evolved than the direct engulfment feeding dinoflagellates.

3c. Feeding mechanism for bacterial prey

Is a dinoflagellate able to feed on bacterium cells using raptorial feeding? Dinoflagellates may have difficulty in detecting and capturing tiny bacterium cells. Jeong et al. (2008) revealed that the HTDs *Oxyrrhis marina* and *Gyrodinium* spp. fed on a single bacterium cell like filter/interception feeders; they generate feeding currents using the flagella. In *O. marina* feeding, feeding currents moved from above the predator toward the cingular depression of the predator along the flow lines (Fig. 3A). *O. marina* intercepted and then ingested a single bacterial cell in the feeding current. In *Gyrodinium* spp. feeding, feeding currents flowed from above the epicone of the predator, via the long and narrow sulcus, to below the hypocone of the predator or flowing along the sides of the body of the predator, via the cingulum and the sulcus, to below the hypocone. *Gyrodinium* spp. ingested a single living bacterium in the feeding currents when the prey arrived at the lower part of the sulcus, by interception. Displacement in the cingulum of dinoflagellates (like *Gyrodinium* spp.) makes a longer and narrower sulcus that in turn may generate a stronger feeding current due to venturi effect (Fig. 3B). Also, bacterial cells in the narrower sulcus may be more easily captured by the predator than those in the wider sulcus. The peduncle cover of *Pfiesteria piscicida* is likely to make a narrow duct and thus may also generate a strong feeding current to feed on pico-sized prey (Fig. 3C). Thus, morphological characteristics of HTDs such as cingular depression, long and narrow sulcus, and probably peduncle cover may be a result of evolution to feed on pico-sized prey. It

Fig. 3. Some morphological characteristics of heterotrophic dinoflagellates such as cingular depression (A. *Oxyrrhis marina*), long and narrow sulcus (B. *Gyrodinium* sp.), and probably the peduncle cover of *Pfiesteria piscicida* (C) may be a result of evolution to feed on pico-sized prey. Displacement in the cingulum of dinoflagellates (B) makes a longer and narrower sulcus which in turn may generate a stronger feeding current due to venturi effect. c: cingular depression, s: sulcus, pc: peduncle cover

is worthwhile to explore relationships among the morphology, feeding mechanisms, and evolution of dinoflagellates.

3d. Number of feeding mechanisms for a dinoflagellate predator

Oxyrrhis marina and *Gyrodinium* spp. exhibit 2 different feeding behaviors; they feed on pico-sized prey as described above and they engulf nano- or micro-sized prey through the cingulum depression and the sulcus. The MTD *Karlodinium armiger* also exhibit 2 different feeding mechanisms (Berge et al. 2008): it fed on small prey cells using direct engulfment, while it fed on large prey using peduncle feeding. Some dinoflagellates may have evolved to feed on more diverse prey items by utilizing different feeding behaviors depending on the type and/or size of prey. These dinoflagellates have an advantage in gaining energy compared to predators that have only one feeding behavior that limits them to being able to feed on only a certain sized prey.

Which feeding mechanism of dinoflagellates is better? The MTD *Fragilidium* cf. *mexicanum* can feed on the HTD

Table 3. Maximum growth rate of each mixotrophic dinoflagellate species when growing photosynthetically (ESD: Equivalent spherical diameter (μm), T: Temperature ($^{\circ}\text{C}$), MGR: Maximum growth rate (d^{-1}))

Species name	ESD	T	MGR	Ref.
<i>Prorocentrum minimum</i>	12.1	25	0.942	Kondo et al. (1990)
<i>Prorocentrum donghaiense</i>	13.2	20	0.821	Jeong et al. (unpubl data)
<i>Heterocapsa triquetra</i>	15.1	15	0.720	Hansen (2002)
<i>Heterocapsa circularisquama</i>	18.0	30	0.901	Yamaguchi et al. (1997)
<i>Gymnodinium aureolum</i>	19.4	20	0.120	Jeong et al. (2010)
<i>Gymnodinium impudicum</i>	22.0	30	0.580	Jeong et al. (unpubl data)
<i>Karenia brevis</i>	23.0	20	0.400	Richardson et al. (2006)
<i>Ceratium lineatum</i>	25.0	15	0.220	Hansen et al. (2007)
<i>Cochlodinium polykrikoides</i>	25.8	25	0.576	Jeong et al. (unpubl data)
<i>Prorocentrum micans</i>	26.6	20	0.531	Jeong et al. (unpubl data)
<i>Ceratium furca</i>	28.1	24	0.720	Baek et al. (2008)
<i>Alexandrium affine</i>	29.7	25	0.598	Jeong et al. (unpubl data)
<i>Alexandrium tamarense</i>	29.7	15	0.381	Yamamoto & Tarutani (1997)
<i>Akashiwo sanguinea</i>	30.8	25	0.783	Matsubara et al. (2007)
<i>Gonyaulax polygramma</i>	32.5	20	0.488	Jeong et al. (unpubl data)
<i>Gymnodinium catenatum</i>	33.9	25	0.407	Jeong et al. (unpubl data)
<i>Alexandrium catenella</i>	34.0	14	0.208	Navarro et al. (2006)
<i>Gyrodinium instriatum</i>	36.5	20	0.510	Nagasoe et al. (2006)
<i>Lingulodinium polyedrum</i>	36.6	20	0.182	Jeong et al. (2005c)
<i>Fragilidium subglobosum</i>	45.0	15	0.160	Hansen and Nielsen (1997)

Table 4. Optimal prey and maximum mixotrophic growth (MGR), ingestion (MIR), and clearance rates (MCR) of each mixotrophic dinoflagellate predator species (ESD: Equivalent spherical diameter (μm), RPP: ratio of predator to prey ESD, T: Temperature ($^{\circ}\text{C}$), LI: Light intensity ($\mu\text{E m}^{-2}\text{s}^{-1}$), MGR: Maximum growth rate (d^{-1}), MIR: Maximum ingestion rate ($\text{ng C grazer}^{-1}\text{d}^{-1}$), MCR: Maximum clearance rate ($\mu\text{l grazer}^{-1}\text{h}^{-1}$))

Predator	ESD	Optimal prey	ESD	RPP	T	LI	MGR	MIR	MCR	Ref.
<i>Gymnodinium gracilentum</i> *	9.1	<i>Teleaulax amphioxeia</i>	8.5	1.1	15	60	1.510	0.08		(1)
<i>Karlodinium veneticum</i>	11.0	<i>Storeatula major</i>	5.6	2.0	20	250	0.520	0.13	7.50	(2)
<i>Gyrodinium galatheanum</i>	11.0	<i>Storeatula major</i>	5.6	2.0	20	100	0.940	0.12		(3)
<i>Paragymnodinium shiwhaense</i>	12.4	<i>Teleaulax</i> sp.	5.6	2.2	20	20	1.097	0.38	0.7	(4)
<i>Prorocentrum donghaiense</i>	13.2	<i>Teleaulax</i> sp.	5.6	2.4	20	20	0.510	0.03	0.04	(5)
<i>Karlodinium armiger</i>	16.7	<i>Rhodomonas baltica</i>	10.7	1.6	15	180	0.650	0.97	10.80	(6)
<i>Gymnodinium aureolum</i>	19.4	<i>Teleaulax</i> sp.	5.6	3.5	20	20	0.169	0.06	0.003	(7)
<i>Cochlodinium polykrikoides</i>	25.8	<i>Teleaulax</i> sp.	5.6	4.6	20	50	0.324	0.16	0.33	(8)
<i>Prorocentrum micans</i>	26.6	<i>Teleaulax</i> sp.	5.6	4.8	20	20	0.197	0.04	0.05	(5)
<i>Ceratium furca</i>	29.0	Natural prey populations			11.5	100	0.600	3.55		(9)
<i>Gonyaulax polygramma</i>	32.5	<i>Teleaulax</i> sp.	5.6	5.8	20	50	0.278	0.18	0.18	(10)
<i>Dinophysis acuminata</i>	35.0	<i>Myrionecta rubra</i>	22.0	1.6	20	60	0.950			(11)
<i>Lingulodinium polyedrum</i>	36.6	<i>Scrippsiella trochoidea</i>	25.1	1.5	20	50	0.303	0.36	0.14	(5)
<i>Fragilidium subglobosum</i>	45.0	<i>Ceratium tripos</i>	59.5	0.8	15	45	0.500	6.27		(12)
<i>Dinophysis norvegica</i>	45.0	Natural prey populations			10	67	0.630	2.59		(9)
<i>Dinophysis acuta</i>	48.5	Natural prey populations			10	67	0.410	1.49		(9)
<i>Fragilidium cf. mexicanum</i>	54.5	<i>Lingulodinium polyedrum</i>	37.9	1.4	22	20	0.360	7.00	6.00	(13)

Note. (1): Jakobsen et al. (2000), (2): Adolf et al. (2006), (3): Li et al. (2000), (4): Yoo et al. (2010), (5): Jeong et al. (2005c), (6): Berge et al. (2008), (7): Jeong et al. (2010), (8): Jeong et al. (2004c), (9): Granéli et al. (1997), (10): Jeong et al. (2005d), (11): Park et al. (2006), (12): Hansen and Nielsen (1997), (13): Jeong et al. (1999a). *: Heterotrophic dinoflagellate conducting mixotrophy using kleptoplastids from prey cells.

Table 5. Optimal prey and maximum growth, ingestion, and clearance rates of each heterotrophic dinoflagellate predator species (ESD: Equivalent spherical diameter (μm), FM: Feeding mechanism, DE: Direct engulfment, PD: Peduncle, PL: Pallium, Tx: Taxon, RPP: Ratio of predator to prey ESD, T: Temperature ($^{\circ}\text{C}$), MGR: Maximum growth rate (d^{-1}), MIR: Maximum ingestion rate ($\text{ng C grazer}^{-1} \text{d}^{-1}$), MCR: Maximum clearance rate ($\mu\text{l grazer}^{-1} \text{h}^{-1}$), BL: Blood, CR: Cryptophyte, DA: Diatom, DN: Dinoflagellate, RA: Raphidophyte, *: prey species for supporting MGR and MIR of *Polykrikos kofoidii* were different)

Predator	ESD	FM	Optimal Prey	Tx	ESD	RPP	T	MGR	MIR	MCR	Ref.
<i>Gymnodinium</i> sp.	6.0	DE	<i>Rhodomonas salina</i>	CR	8.0	0.8	15	0.936	0.04	0.05	(1)
<i>Protoperidinium bipes</i>	7.8	PL	<i>Skeletonema costatum</i>	DA	5.9	1.3	20	1.370	2.9	1.00	(2)
<i>Protoperidinium hirobis</i>	8.7	PL	<i>Leptocylindrum danicus</i>	DA	19.7	0.4	20	1.230	0.8	0.50	(3)
<i>Luciella masanensis</i>	13.5	PD	perch bloods	BL	6.1	2.2	20	1.460	2.6	0.83	(4)
<i>Pfiesteria piscicida</i>	13.5	PD	perch bloods	BL	6.1	2.2	20	1.740	4.3	2.50	(5)
<i>Stoeckeria algicida</i>	13.9	PD	<i>Heterosigma akashiwo</i>	RA	11.5	1.2	20	1.630	0.8	3.70	(6)
<i>Oxyrrhis marina</i>	15.6	DE	<i>Heterosigma akashiwo</i>	RA	11.5	1.4	20	1.430	1.3	0.30	(7)
<i>Gyrodinium dominans</i>	20.0	DE	<i>Skeletonema costatum</i>	DN	5.4	3.7	24	0.370	-		(8)
<i>Protoperidinium vorax</i>	21.0	PL	<i>Sk. pseudocostatum</i>	DA	4.0	5.3	20	1.090	2.3	19.90	(9)
<i>Oblea rutunda</i>	21.6	PL	<i>Ditylum brightwellii</i>	DA	33.0	0.7	20	0.660	1.3	0.70	(10)
<i>Protoperidinium steinii</i>	25.8	PL	<i>Heterocapsa triquetra</i>	DN	15.0	1.7	15	0.180			(11)
<i>Protoepridinium huberi</i>	26.5	PL	<i>Ditylum brightwellii</i>	DA	26.5	1.0	20	0.720	17.8	23.00	(12)
<i>Diplopsalis lenticula</i>	31.0	PL	<i>Ditylum brightwellii</i>	DA	35.2	0.9	15	0.250	19.4	2.70	(13)
<i>Gyrodinium spirale</i>	31.8	DE	<i>Prorocentrum minimum</i>	DN	12.1	2.6	20	0.790	13.6	5.30	(14)
<i>Protoperidinium excentricum</i>	35.8	PL	<i>Ditylum brightwellii</i>	DA	35.2	1.0	12	0.330			(15)
<i>Protoperidinium pellucidum</i>	36.1	PL	<i>Ditylum brightwellii</i>	DA	26.5	1.4	20	0.700	11.5		(16)
<i>Polykrikos kofoidii</i> *	43.5	DE	<i>Gymnodinium catenatum</i>	DN	34.0	1.3	20	1.120	17.1	4.60	(17)
<i>Polykrikos kofoidii</i> *	43.5	DE	<i>Lingulodinium polyedrum</i>	DN	37.9	1.1	20	0.826	24.4	5.90	(17)
<i>Protoperidinium spiniferum</i>	45.0	PL	<i>Leptocylindrum danicus</i>	DA	19.7	2.3	20	0.300			(3)
<i>Protoperidinium conicum</i>	45.7	PL	<i>Ditylum brightwellii</i>	DA	35.2	1.3	12	1.130	17.7		(15)
<i>Protoperidinium pallidum</i>	46.6	PL	<i>Ditylum brightwellii</i>	DA	35.2	1.3	15	0.280			(11)
<i>Zygbikodinium lenticulatum</i>	50.0	PL	<i>Ditylum brightwellii</i>	DA	35.2	1.4	15	0.200			(11)
<i>Protoperidinium divergens</i>	61.0	PL	<i>Lingulodinium polyedrum</i>	DN	38.2	1.6	19	0.484	12.0	0.67	(18)
<i>Protoperidinium crassipes</i>	73.0	PL	<i>Lingulodinium polyedrum</i>	DN	36.6	2.0	19	0.308	5.3	0.47	(18)
<i>Protoperidinium depressum</i>	81.0	PL	<i>Ditylum brightwellii</i>	DA	35.2	2.3	12	0.210	12.1		(15)

Note. (1): Jakobsen and Hansen (1997), (2): Jeong et al. (2004d), (3): Jacobson and Anderson (1986), (4): Jeong et al. (2007a), (5): Jeong et al. (2006), (6): Jeong et al. (2005a), (7): Jeong et al. (2003a), (8): Nakamura et al. (1995), (9): Siano and Montresor (2005), (10): Strom and Buskey (1993), (11): Naustvoll (2000), (12): Buskey et al. (1994), (13): Naustvoll (1998), (14): Kim and Jeong (2004), (15): Menden-Deuer et al. (2005), (16): Buskey (1997), (17): Jeong et al. (2001b), (18): Jeong and Latz (1994)

Protoperidinium cf. *divergens*, while *P. cf. divergens* also feed on *F. cf. mexicanum* (Jeong et al. 1997). However, in general, *F. cf. mexicanum* predominates over *P. cf. divergens* in their competition even though their sizes are similar. Also, the engulfment feeding *Oxyrrhis marina* dominated over the peduncle feeding *Pfiesteria piscicida*, *Stoeckeria algicida*, and *Luciella masanensis* (Jeong et al. 2007c). Therefore, engulfment feeding may be a more effective feeding mechanism than pallium feeding or peduncle feeding if there are no supplementary tools such as nematocysts or trichocysts. Feeding mechanisms may be a critical factor determining predator and prey relationships among dinoflagellates and thus it is worthwhile exploring this topic further.

4. Growth and Ingestion Rates

The range of the growth and ingestion rates of dinoflagellates is very wide (Table 3, 4, 5). The growth and ingestion rates of dinoflagellates have been known to be affected by diverse factors; prey species and concentrations are primary factors affecting the rates (e.g. Jeong et al. 2007a). In this section, we examined whether the growth and ingestion rates of dinoflagellate predators are also affected by size, trophic mode, or feeding mechanisms of the predators or not.

4a. Predator size effect

When the optimal prey for each dinoflagellate predator species was provided, MGRs of dinoflagellates growing autotrophically (ATDs) and HTDs were significantly

Fig. 4. The maximum growth rate (MGR, d^{-1}) of phototrophic dinoflagellates growing exclusively photosynthetically (open triangles) and growing mixotrophically (open circles) and heterotrophic dinoflagellates (closed circles) as a function of the size (Equivalent Spherical Diameters, ESD, μm) of the predator when each predator fed on the optimal prey species (see Table 2, 3, 4). The equation of the regression was $MGR (d^{-1}) = -0.017 \times (ESD \text{ of predator}) + 9.80$, $r^2 = 0.351$ for autotrophic growth ($n = 20$, $p < 0.01$); $MGR (d^{-1}) = -0.016 \times (ESD \text{ of predator}) + 1.31$, $r^2 = 0.472$ for heterotrophic growth ($n = 25$, $p < 0.005$). *Gymnodinium gracilentum* (Gg) is a heterotrophic dinoflagellate conducting mixotrophy using kleptoplastids from prey cells and thus was treated as a MTD in this calculation. Temperature effect was not considered because optimal temperature for each dinoflagellate species is different from the other

negatively correlated with the predator size ($p < 0.01$ for ATD and $p < 0.005$ for HTD; Fig. 4). However, MIRs of MTDs and HTDs were significantly positively correlated with the predator size ($p < 0.01$ for MTD and $p < 0.05$ for HTD; Fig. 5). When single prey item was offered, feeding by dinoflagellates on some prey was affected by the predator size, while that on other prey was not; ingestion rates by MTDs and HTDs on heterotrophic bacteria (ESD $< 0.5 \mu m$) were not significantly affected by the size of the predators (Seong et al. 2006; Jeong et al. 2008), while those on the cyanobacterium *Synechococcus* sp. (ESD = $2 \mu m$) were affected (Jeong et al. 2005b). Ingestion rates by engulfment feeding MTDs on an unidentified cryptophyte were also affected by the predator size (Jeong et al. 2005c). Thus, for the filter/interception feeding dinoflagellate predator (on heterotrophic bacteria), the shape (cingulum depression, displacement of the cingulum as described above) of the dinoflagellate predators may be more important than the size, while for engulfment feeding dinoflagellate predator (on larger algal prey), the size of the predator may be an

important factor affecting the ingestion rates of the predators.

4b. Trophic mode effect

The highest MGR of the HTDs ($1.7 d^{-1}$ for *Pfiesteria piscicida*; Jeong et al. 2006) is double the highest MGR of dinoflagellates growing autotrophically ($0.94 d^{-1}$ for *Prorocentrum minimum*; Kondo et al. 1990) (Table 3, 4, 5). Also, MGRs of small HTDs are much higher than those of similar sized ATDs (Fig. 4). Energy gain of small HTDs through feeding may be higher than that of small ATDs through photosynthesis. Also, enzymes involved in photosynthesis may lower MGRs of dinoflagellates and it is worthwhile exploring this topic. The range of MIRs of each HTD was $0.04\text{--}24.4 \text{ ng C dinoflagellate}^{-1} d^{-1}$, while that of each MTD species was $0.03\text{--}7.0 \text{ ng C dinoflagellate}^{-1} d^{-1}$ (Fig. 5). Also, MIRs of HTDs were higher than those of similar sized MTDs (Fig. 5). Heterotrophic activity of HTDs (feeding and digestion) is likely to be higher than that of MTDs. Based on these results, we suggest that to increase growth and ingestion rates, some MTDs may evolve to HTDs by discarding the plastids. It is worthwhile to explore the relationships between growth and ingestion rates of dinoflagellates and evolution.

Fig. 5. The maximum ingestion rate (MIR, $ng \text{ C dinoflagellate}^{-1} d^{-1}$) of phototrophic dinoflagellates growing mixotrophically (open circles) and heterotrophic dinoflagellates (closed circles) as a function of the size (Equivalent Spherical Diameters, ESD, μm) of the predator when each predator fed on the optimal prey species (see Table 4, 5). The equation of the regression was $MIR (ng \text{ C dinoflagellate}^{-1} d^{-1}) = 0.108 \times (ESD \text{ of predator}) - 1.49$, $r^2 = 0.517$ for mixotrophs ($n = 16$, $p < 0.01$); $MIR (ng \text{ C dinoflagellate}^{-1} d^{-1}) = 0.178 \times (ESD \text{ of predator}) - 2.89$, $r^2 = 0.265$ for heterotrophs ($n = 16$, $p < 0.05$)

Fig. 6. The maximum growth (MGR, d^{-1} , A) and ingestion rate (MIR, $ng\ C\ dinoflagellate^{-1}d^{-1}$, B) of heterotrophic dinoflagellates having different feeding mechanisms as a function of the size (Equivalent Spherical Diameters, ESD, μm) of the predator when each predator fed on the optimal prey species (see Table 5). Pink triangles: Peduncle feeders (PE). Green squares: Direct engulfment feeders (EG). Blue circles: Pallium feeders (PA). The equation of the regression for PA in (A) was $MGR\ (d^{-1}) = -0.012 \times (ESD\ of\ predator) + 1.06$, $r^2 = 0.382$ ($n = 16$, $p < 0.05$); The equation of the regression for EG in (B) was $MIR\ (ng\ C\ dinoflagellate^{-1}d^{-1}) = 0.523 \times (ESD\ of\ predator) - 5.09$, $r^2 = 0.944$ for mixotrophs ($n = 5$, $p < 0.05$)

4c. Feeding mechanism effect

When the predators' ESDs were $< 21\ \mu m$, the mean of the 3 highest MGRs of the peduncle feeders (*Pfiesteria piscicida*, *Stoeckeria algicida*, and *Luciella masanensis*) was significantly higher than that of the engulfment feeding HTDs (*Oxyrrhis marina*, *Gymnodinium* sp., and *Gyrodinium dominans*) and the pallium feeding HTDs (*Protoperidinium bipes*, *P. hirobis*, and *P. vorax*) ($p < 0.05$ for both, one-tailed t-test) which were not significantly different from each other ($p > 0.1$, two-tailed t-test) (Fig. 6A). To the contrary, the mean of the 2 highest MIRs of the peduncle feeders was significantly lower than that of the engulfment feeders and

the pallium feeders ($p < 0.05$, one-tailed t-test) which were not significantly different from each other ($p > 0.1$, two-tailed t-test) (Fig. 6B). Therefore, the growth efficiency of the peduncle feeders is higher than that of the engulfment feeders or the pallium feeders; The peduncle feeding HTDs may spend less energy cost to feed, ingest, and grow compared to the engulfment feeding or the pallium feeding HTDs. Engulfing a prey cell may spend large energy in opening and closing the sulcus or the apical horn, the suture using microtubules. Also, the pallium feeding may spend large energy in towing and handling large prey cells and in form large pallium. However, the peduncle feeding HTD may just form tiny feeding tube and suck prey materials.

4d. Optimal prey size

When the MGR of each dinoflagellate species on its optimal prey was achieved, the range of the ratios of MTD predator to prey size (0.8–5.8) is similar to that of HTD predator to prey size (0.4–5.3) (Fig. 7A and B). The high MGRs (i.e. $> 0.8\ d^{-1}$ for MTDs and $> 1.2\ d^{-1}$ for HTDs) were obtained when the ratios for MTDs and HTDs were 1.1–2.4 and 0.4–2.2, respectively. The ratios of predator to prey size when the 3 highest MGRs were achieved were 0.4–1.3 for the pallium feeding HTDs, 1.3–1.4 for the engulfment feeding HTDs, and 1.2–2.2 for the peduncle feeding HTDs (Fig. 7B). The lowest ratio was achieved by the pallium feeding HTDs because they are able to feed on prey larger than themselves (e.g. Naustvoll 2000).

When the high MIRs (i.e. $> 5\ ng\ C\ dinoflagellate^{-1}d^{-1}$ for MTDs and $> 10\ ng\ C\ dinoflagellate^{-1}d^{-1}$ for HTDs) were obtained, the range of the ratios of HTDs predator to prey size (0.9–2.6) was wider than that of MTDs (0.8–1.4) (Fig. 7C–D). When the 2 highest MIRs were achieved, the ratios of predator to prey size of pallium feeding HTDs (0.9–1.0) were markedly lower than those of peduncle feeding HTDs (1.2–2.2). Therefore, the highest MIRs were achieved when the pallium feeding HTDs fed on prey similar to themselves, while the peduncle feeders feed on prey having half the size of the predators. Therefore, the optimal ratios of predator to prey size of dinoflagellates are affected by the types of the feeding mechanisms, which may cause separation of ecological niches of the dinoflagellates.

4e. Light and nutrient effects

In addition to the biological factors, light and nutrient conditions sometimes affect the ingestion rates of MTDs;

Fig. 7. The maximum growth (d^{-1}) and ingestion rates (MIR , $ng\ C\ dinoflagellate^{-1}\ d^{-1}$) of mixotrophic (A, C) and heterotrophic dinoflagellates having different feeding mechanisms (B, D) as a function of the ratio of predator relative to prey size when each predator fed on the optimal prey species (see Table 5) (Pink triangles: Peduncle feeders (PE), Green squares: Direct engulfment feeders (EG), Blue circles: Pallium feeders (PA))

the ingestion rates of the MTDs *Fragilidium subglobosum*, *Gymnodinium gracilentum*, and *Karlodinium veneficum* (previously *Gyrodinium galatheanum* and *Karlodinium micrum*) increased continuously, or increased and then were saturated, with increasing light intensity up to ca. 75–100 $\mu E\ m^{-2}\ s^{-1}$ (Skovgaard 1996b, 1998, Hansen and Nielsen 1997, Li et al. 2000, Skovgaard et al. 2000), while those of *F. cf. mexicanum* (20–200 $\mu E\ m^{-2}\ s^{-1}$, Jeong et al. 1999a) and *P. minimum* (between 6% and 100% incident light intensity; Stoecker et al. 1997) were not significantly affected by light intensity under the provided conditions. The ingestion rates of *Ceratium furca* and *K. veneficum* were also affected by nutrient conditions (Li et al. 2000; Smalley et al. 2003), while that of *F. cf. mexicanum* (Jeong et al. 1999a) was not significantly affected by nutrient conditions. Therefore, the effects of light and nutrient conditions on feeding in MTDs may depend on the species. Mixotrophy in phototrophic dinoflagellates, whose feeding is not affected by light and

nutrient conditions, may enable the dinoflagellates to have an advantage in surviving using photosynthesis when the abundance of prey is low, while mixotrophy in phototrophic dinoflagellates, whose feeding is affected by light and nutrient conditions, may enable the dinoflagellates to have an advantage in surviving, using phagotrophy when light and nutrient conditions are not favorable for photosynthesis. Therefore, light and nutrient conditions may be important driving forces on the evolution of trophic modes of MTDs.

5. Grazing Impact by Dinoflagellates on the Natural Populations of Prey

Dinoflagellate predators often have considerable grazing impact on and sometimes control the natural population of prey including heterotrophic bacteria (Seong et al. 2006; Jeong et al. 2008), cyanobacteria (Jeong et al. 2005b), nanoflagellates (Jeong et al. 2007b), diatoms (Jeong et al.

2004d; Yoo et al. 2009), other dinoflagellates (Jeong et al. 2005c), and ciliates (Smalley and Coats 2002). In general, the grazing impact by MTDs and HTDs on a population of co-occurring prey increased with increasing the abundance of the dinoflagellate predators. The ingestion rates and also grazing impact by dinoflagellates on bacteria are comparable to those by mixotrophic nanoflagellates (MNF) or HNFs because the abundance of the former predators is also comparable to that of the nanoflagellates (Seong et al. 2006; Jeong et al. 2005b, 2008). However, the abundance of dinoflagellates is usually much higher than that of ciliates or copepods. Therefore, even though the ingestion rates of dinoflagellates are lower than that of ciliates and copepods, grazing impact of the former predators on natural population of prey are usually higher than that of the latter predators. For example, grazing coefficients attributable to *Acartia* spp. on *Prorocentrum minimum* in the Korean coastal waters (0–0.001 h⁻¹) are much lower than those for *Gyrodinium*

dominans on co-occurring *P. minimum* (0–0.07 h⁻¹) or for *G. spirale* (0–0.23 h⁻¹) (e.g. Kim and Jeong 2004). Much lower abundances of *Acartia* spp. (0.01–0.40 ind. l⁻¹) compared to those of *Gyrodinium* spp. (1,600–140,000 ind. l⁻¹) may be responsible for these lower grazing coefficients, even though the ingestion rates of *Acartia* spp. on *P. minimum* (10,900 ng C grazer⁻¹d⁻¹) are much higher than that of *Gyrodinium* spp. (1–14 ng C grazer⁻¹d⁻¹). Therefore, when investigating the population dynamics of target prey, we must measure the grazing impact by dinoflagellate predators.

6. Mortality due to Predation

To explore the population dynamics of dinoflagellates, the mortality rate due to predation in addition to the growth rate should be obtained. Other MTDs (Jeong et al. 2005c), HTDs (Hansen 1991), ciliates (Jeong et al. 1999b), copepods (Berggreen et al. 1988; Jeong 1994a; Kim and Jeong 2004;

Table 6. Maximum ingestion and clearance rates of heterotrophic dinoflagellates (HTD) on mixotrophic (MTD) and HTDs (RPP: Ratio of predator to prey ESD, T: Temperature (°C), MIR: Maximum ingestion rate (ng C grazer⁻¹d⁻¹), MCR: Maximum clearance rate (μl grazer⁻¹h⁻¹), ESD: Equivalent spherical diameter (μm), *: Highest MIR and MCR of each predator species)

Predator	ESD	Prey	Taxon	ESD	RPP	T	MIR	MCR	Ref.
<i>Luciella masanensis</i>	13.5	<i>Amphidinium carterae</i>	MTD	9.7	1.4	20	0.3*	0.06*	(1)
<i>Pfiesteria piscicida</i>	13.5	<i>Heterocapsa rotundata</i>	MTD	5.8	2.3	20	0.2	0.01	(2)
<i>Pfiesteria piscicida</i>	13.5	<i>Amphidinium carterae</i>	MTD	9.7	1.4	20	1.1*	1.50*	(2)
<i>Pfiesteria piscicida</i>	13.5	<i>Akashiwo sanguinea</i>	MTD	30.8	0.4	20	0.1	0.01	(2)
<i>Pfiesteria piscicida</i>	13.5	<i>Gymnodinium catenatum</i>	MTD	33.9	0.4	20	0.2	0.01	(2)
<i>Gyrodinium dominans</i>	15.3	<i>Prorocentrum minimum</i>	MTD	12.1	1.3	20	1.2	0.90*	(3)
<i>Oxyrrhis marina</i>	15.6	<i>Amphidinium carterae</i>	MTD	9.7	1.6	20	2.8	2.40	(4)
<i>Oxyrrhis marina</i>	15.6	<i>Karlodinium veneficum</i>	MTD	9.1	1.7	20	5.5*	2.40*	(5)
<i>Oxyrrhis marina</i>	15.6	<i>Prorocentrum minimum</i>	MTD	12.1	1.3	20	2.0	0.80	(6)
<i>Gyrodinium dominans</i>	20.0	<i>Heterocapsa triquetra</i>	MTD	15.3	1.3	24	7.5*	0.19	(7)
<i>Gyrodinium spirale</i>	28.0	<i>Heterocapsa triquetra</i>	MTD	15.8	1.8	15	7.5	0.30	(8)
<i>Gyrodinium spirale</i>	31.8	<i>Prorocentrum minimum</i>	MTD	12.1	2.6	20	13.6*	5.30*	(3)
<i>Protoberidinium pellucidum</i>	36.1	<i>Prorocentrum micans</i>	MTD	32.7	1.1	20	7.7*		(9)
<i>Polykrikos kofoidii</i>	43.5	<i>Gymnodinium catenatum</i>	MTD	34.0	1.3	20	17.1	4.60	(10)
<i>Polykrikos kofoidii</i>	43.5	<i>Scrippsiella trochoidea</i>	MTD	25.1	1.7	20	16.6	1.10	(10)
<i>Polykrikos kofoidii</i>	43.5	<i>Lingulodinium polyedrum</i>	MTD	37.9	1.1	20	24.4*	5.90*	(10)
<i>Polykrikos kofoidii</i>	43.5	<i>Ceratium furca</i>	MTD	29.0	1.5	20	9.8	3.70	(10)
<i>Polykrikos kofoidii</i>	43.5	<i>Prorocentrum micans</i>	MTD	26.0	1.7	20	4.6	2.30	(10)
<i>Polykrikos kofoidii</i>	43.5	<i>Gyrodinium impudicum</i>	MTD	23.2	1.9	20	5.4	1.30	(10)
<i>Protoberidinium divergens</i>	61.0	<i>Lingulodinium polyedrum</i>	MTD	38.2	1.6	19	12.0*	0.67*	(11)
<i>Protoberidinium crassipes</i>	73.0	<i>Lingulodinium polyedrum</i>	MTD	36.6	2.0	19	5.3*	0.47*	(11)
<i>Oxyrrhis marina</i>	15.6	<i>Pfiesteria piscicida</i>	HTD	13.5	1.2	20	0.3*	0.34*	(12)
<i>Oxyrrhis marina</i>	15.6	<i>Stoeckeria algicida</i>	HTD	13.5	1.2	20	0.1	0.61	(12)
<i>Oxyrrhis marina</i>	15.6	<i>Luciella masanensis</i>	HTD	13.5	1.2	20	0.1	0.01	(12)

Note. (1): Jeong et al. (2007a), (2): Jeong et al. (2006), (3): Kim and Jeong (2004), (4): Jeong et al. (2001a), (5): Adolf et al. (2007), (6): Lee (1998), (7): Nakamura et al. (1995), (8): Hansen (1992), (9): Buskey (1997), (10): Jeong et al. (2001b), (11): Jeong and Latz (1994), (12): Jeong et al. (2007b)

Table 7. Maximum ingestion and clearance rates of ciliates on mixotrophic (MTD) and heterotrophic dinoflagellate prey (HTD) (T: Temperature (°C), MIR: Maximum ingestion rate (ng C grazer⁻¹d⁻¹), MCR: Maximum clearance rate (μl grazer⁻¹h⁻¹), ESD: Equivalent spherical diameter (μm), *: Highest MIR and MCR of each predator species)

Predator	ESD	Prey	Taxon	ESD	RPP	T	MIR	MCR	Ref.
<i>Balanion</i> sp.	32	<i>Heterocapsa triquetra</i>	MTD	15.0	2.1	15	20*	1.2*	(1)
<i>Tiarina fusus</i>	37	<i>Lingulodinium polyedrum</i>	MTD	37.9	1.0	19	23*	4.5*	(2)
<i>Tiarina fusus</i>	37	<i>Scrippsiella trochoidea</i>	MTD	25.1	1.5	19	10	0.2	(2)
<i>Tiarina fusus</i>	37	<i>Prorocentrum minimum</i>	MTD	12.8	2.9	19	2	0.2	(2)
<i>Tiarina fusus</i>	37	<i>Amphidinium carterae</i>	MTD	16.1	2.3	19	3	0.01	(2)
<i>Tiarina fusus</i>	37	<i>Heterocapsa triquetra</i>	MTD	12.8	2.9	19	3	0.0	(2)
<i>Tiarina fusus</i>	37	<i>Prorocentrum micans</i>	MTD	26.0	1.4	19	3	0.4	(2)
<i>Favella</i> sp.	72	<i>Heterocapsa triquetra</i>	MTD	15.0	4.8	15	130	11.4	(1)
<i>Favella</i> sp.	74	<i>Scrippsiella trochoidea</i>	MTD	26.5	2.8	20	237*	43.0*	(1)
<i>Strombidinopsis</i> sp.	102	<i>Cochlodinium polykrikoides</i>	MTD	23.3	4.4	19	353*	52.3	(3)
<i>Strombidinopsis</i> sp.	102	<i>Gymnodinium sanguineum</i>	MTD	36.3	2.8	19	343	85.6	(3)
<i>Strombidinopsis</i> sp.	102	<i>Prorocentrum minimum</i>	MTD	12.8	8.0	19	267	110.0*	(3)
<i>Strombidinopsis</i> sp.	102	<i>Lingulodinium polyedrum</i>	MTD	37.9	2.7	19	222	110.0	(3)
<i>Strombidinopsis</i> sp.	102	<i>Scrippsiella trochoidea</i>	MTD	25.1	4.1	19	207	41.0	(3)
<i>Strombidinopsis jeokjo</i>	94	<i>Gyrodinium dominans</i>	HTD	15.3	6.1	20	108*	14.5*	(4)
<i>Strombidinopsis jeokjo</i>	94	<i>Oxyrrhis marina</i>	HTD	15.6	6.0	20	87	13.4	(4)
<i>Strombidinopsis jeokjo</i>	94	<i>Pfiesteria piscicida</i>	HTD	13.5	6.9	20	44	15.4	(5)
<i>Strombidinopsis jeokjo</i>	94	<i>Stoeckeria algicida</i>	HTD	13.5	6.9	20	49	6.6	(5)
<i>Strombidinopsis jeokjo</i>	94	<i>Luciella masanensis</i>	HTD	13.5	6.9	20	10	5.4	(5)

Note. (1): Stoecker and Evans (1985), (2): Jeong et al. (2002), (3): Jeong et al. (1999), (4): Jeong et al. (2004b), (5): Jeong et al. (2007c)

Verity and Paffenhöfer 1996), larvae of mussels (Jeong et al. 2004a) and polychaeta (Song and Jeong unpubl. data; Watras et al. 1985), rotifers (Heinbokel et al. 1988), and larval fish (Scura and Jerde 1977) have been known to feed on MTDs. Also, HTDs were fed by MTDs (Jeong et al. 1997), other HTDs, ciliates (Jeong et al. 2004b; Stoecker et al. 2002) and copepods (Koski and Riser 2006). However, there has been no report on the feeding by larvae of mussels and polychaeta and rotifers on HTDs yet and it is worthwhile to explore this topic. The range of the ingestion rates of heterotrophic protistan and metazoan predators on MTDs and HTDs is very wide because the taxonomical characteristics, the size, and feeding mechanisms of these predators are diverse and the morphological and biochemical properties of MTD and HTD prey are also diverse.

6a. Predation rates of diverse predators

There have been many studies on ingestion rates of heterotrophic protists and metazoans on MTDs (Table 6, 7, 8). The maximum ingestion rates (MIRs) of HTD species on the optimal MTDs (0.3–24.4 ng C grazer⁻¹d⁻¹) was comparable to that of the small ciliates (< 40 μm in body length; 20–23 ng C grazer⁻¹d⁻¹), but markedly lower than that of the larva of

the mussel *Mytilus galloprovincialis* (69 ng C grazer⁻¹d⁻¹) or large ciliates (130–350 ng C grazer⁻¹d⁻¹) (Table 6, 7). MIRs of the copepod species on MTDs (0.1–73.6 μg C copepod⁻¹d⁻¹) were ca. 10–1000 times higher than those of heterotrophic protists (Table 8). The maximum clearance rates (MCRs) of HTDs, ciliates, and metazoans on MTDs showed a pattern similar to MIRs; MCRs of the HTD species on MTDs (0.01–5.9 μl grazer⁻¹h⁻¹) was comparable to that of the small ciliates (0.01–4.5 μl grazer⁻¹h⁻¹), but lower than that of the larva of *M. galloprovincialis* (11.4 μl grazer⁻¹h⁻¹), large ciliates (11.4–110.0 μl grazer⁻¹h⁻¹), or the larva of the polychaeta *Polydora* sp. (520 μl grazer⁻¹h⁻¹; Watras et al. 1985) (Table 6, 7). MCRs of copepods on MTDs (0.3–5.0 ml grazer⁻¹h⁻¹) were also ca. 10–1000 times higher than those of heterotrophic protists (Table 8). Therefore, in MIRs and MCRs of the predators on MTDs, the general sequence was copepods > large ciliates = the larvae of benthos > small ciliates = HTDs.

There have been only a few studies on feeding by other HTDs or ciliates on HTDs (Jeong et al. 2004b, 2007c). The MIR of the HTD *Oxyrrhis marina* on HTDs (0.3 ng C grazer⁻¹d⁻¹) was much lower than that of the large ciliate *Strombidinopsis jeokjo* (108 ng C grazer⁻¹d⁻¹) (Table 6, 7).

Table 8. Maximum ingestion and clearance rates of copepods on mixotrophic (MTD) and heterotrophic dinoflagellates (HTD) (T: Temperature (°C), MIR: Maximum ingestion rate ($\mu\text{g C copepod}^{-1}\text{d}^{-1}$), MCR: Maximum clearance rate ($\text{ml copepod}^{-1}\text{h}^{-1}$), ESD: Equivalent spherical diameter (μm), *: Highest MIR and MCR of each predator species)

Predator	Prey	Taxon	ESD	T	MIR	MCR	Ref.
<i>Acartia tonsa</i>	<i>Amphidinium carterae</i>	MTD	9.7	20	3.5		(1)
<i>Acartia tonsa</i>	<i>Prorocentrum minimum</i>	MTD	12.0	20	10.0	3.5	(2)
<i>Acartia tonsa</i>	<i>Gymnodinium sanguineum</i>	MTD	28.0	20	3.2	5.0	(3)
<i>Acartia tonsa</i>	<i>Prorocentrum minimum</i>	MTD	12.6	20	19.7*		(4)
<i>Acartia tonsa</i>	<i>Amphidinium carterae</i>	MTD	9.7	20	6.1		(4)
<i>Acartia tonsa</i>	<i>Karenia brevis</i>	MTD	20.4	21	3.8		(5)
<i>Calanus sinicus</i>	<i>Alexandrium tamarense</i> (toxic)	MTD	30.0	18	0.1*	4.0	(6)
<i>Calanus sinicus</i>	<i>Alexandrium tamarense</i> (non-toxic)	MTD	30.0	18	-	1.8	(6)
<i>Calanus helgolandicus</i>	<i>Prorocentrum micans</i>	MTD	30.0	15	73.6*	1.7	(7)
<i>Paracalanus crassirostris</i>	<i>Alexandrium tamarense</i> (toxic)	MTD	30.0	18		0.5	(6)
<i>Paracalanus crassirostris</i>	<i>Alexandrium tamarense</i> (non-toxic)	MTD	30.0	18		0.6	(6)
<i>Paracalanus crassirostris</i>	<i>Prorocentrum dentatum</i>	MTD	14.5	18		0.3	(6)
<i>Temora turbinata</i>	<i>Karenia brevis</i>	MTD	20.4	21	23.4*		(5)
<i>Temora turbinata</i>	<i>Prorocentrum minimum</i>	MTD	10.5	21			(5)
<i>Centropages typicus</i>	<i>Karenia brevis</i>	MTD	20.4	21	56.9*		(5)
<i>Centropages typicus</i>	<i>Prorocentrum minimum</i>	MTD	10.5	21	0.1		(5)
<i>Acartia</i> spp.**	<i>Oxyrrhis marina</i> on <i>P. minimum</i>	HTD	15.6	20	4.6*		(8)
<i>Acartia</i> spp.**	<i>Oxyrrhis marina</i> on <i>Amphidinium carterae</i>	HTD	15.6	20	1.9		(8)
<i>Acartia</i> spp.**	<i>Pfiesteria piscicida</i>	HTD	13.5	20	3.9	0.2	(9)
<i>Acartia</i> spp.**	<i>Stoeckeria algicida</i>	HTD	13.5	20	3.6	0.8	(9)
<i>Acartia</i> spp.**	<i>Luciella masanensis</i>	HTD	13.5	20	3.5	0.4	(9)
<i>Acartia tonsa</i>	<i>Oxyrrhis</i> sp.	HTD	13.9	20	5.0*		(2)
<i>Acartia tonsa</i>	<i>Gyrodinium dominans</i> (on <i>Rhodomonas salina</i>)	HTD	14.6	17-20		2.0	(3)
<i>Acartia tonsa</i>	<i>Gyrodinium dominans</i> (on <i>T. pseudonana</i>)	HTD	14.6	17-20	1.0	2.5	(3)
<i>Acartia tonsa</i>	<i>Pfiesteria piscicida</i>	HTD	10.0	20	4.8	1.4	(4)

Note. (1): Houde and Roman (1987), (2): Besiktepe and Dam (2002), (3): Broglio et al. (2003), (4): Roman et al. (2006), (5): Cohen et al. (2007), (6): Liu and Wang (2002), (7): Huskin et al. (2000), (8): Jeong et al. (2001a), (9): Jeong et al. (2007c). **: *Acartia omorii* & *A. hongii*.

The MCR of *O. marina* on HTDs ($0.014 \mu\text{l grazer}^{-1}\text{h}^{-1}$) was also much lower than that of *S. jeokjo* ($14.5 \mu\text{l grazer}^{-1}\text{h}^{-1}$) (Table 6, 7). Large ciliates may be one of the major predators on HTDs. MIRs ($0.7\text{--}5.0 \mu\text{g C copepod}^{-1}\text{d}^{-1}$) and MCRs ($2.5\text{--}32.6 \text{ml copepod}^{-1}\text{h}^{-1}$) of the copepods on HTDs were also ca. 10–1000 times higher than those of heterotrophic protists (Table 6, 7, 8). To understand the population dynamics of HTDs, much more study on the mortality rate due to predators is necessary.

Calbet et al. (2003) showed that grazing by microzooplankton (several ciliates and rotifers) on the toxic MTD *Alexandrium minutum* was much higher than that by copepods. Higher abundance of microzooplankton than copepods might cause this difference. Therefore, when exploring mortality of dinoflagellates due to predation, we must measure feeding by both microzooplankton and metazooplankton predators on dinoflagellates.

6b. MTD prey vs HTD prey

The growth and ingestion rates of the HTD *Oxyrrhis marina* on MTDs ($n=3$) were significantly higher than those on HTDs ($p<0.05$, one tailed t-test; Fig. 8A). The ingestion rates of the large ciliate *Strombidinopsis* spp. ($n=5$) on MTDs were also significantly higher than those on HTDs ($p<0.05$; Fig. 8B), even though the growth rates of *Strombidinopsis* spp. on MTDs were not significantly higher than those on HTDs ($p>0.1$). The highest MIRs of the copepods *Acartia* spp. on MTDs were also clearly greater than those on HTDs (Fig. 8C). Therefore, compared to MTDs, HTDs may have more effective anti-predation tools against these predators. The maximum swimming speeds of HTDs (*Pfiesteria piscicida*, *Stoeckeria algicida*, and *Luciella masanensis*) were much higher than those of MTDs (*Amphidinium carterae*, *Karlodinium veneticum*, and *Prorocentrum minimum*). Higher swimming speeds of

Fig. 8. The maximum ingestion rate (MIR, ng C grazer⁻¹d⁻¹) of the heterotrophic dinoflagellate *Oxyrrhis marina* (A), the ciliate *Strombidinopsis* spp. (B), and the copepods *Acartia* spp. (C) as a function of the size (Equivalent Spherical Diameters, ESD, μm) when the predator fed on mixotrophic (open circles) and heterotrophic dinoflagellate prey (closed circles) (see Table 6, 7, 8). Pm: *Prorocentrum minimum* prey

the HTDs may escape from being captured by predators compared to MTDs. Also, these HTDs may attack the predators, while MTDs have been not known to attack the predators yet (Jeong et al. 2007c).

6c. Toxic dinoflagellate prey

Toxins of dinoflagellate prey may kill potential predators, be not eaten, or support the positive growth of predators; i) some *Alexandrium* strains caused loss of motility and cell lysis of the HTDs *Oblea rotunda* and *Oxyrrhis marina* by extracellular substances (Tillmann and John 2002). ii) a toxic strain of *A. tamarensis* inhibited feeding by *Favella* spp. (Hansen 1995) and also the toxic naked dinoflagellate *Gyrodinium aureolum* suppresses the growth of *F. ehrenbergii* (Hansen 1989). Also, *Strombidinopsis* sp. and *Tiarina fusus* did not grow on the toxic dinoflagellate *Amphidinium carterae* (Jeong et al. 1999b, 2002). Some copepods have been known to reject some toxic MTDs (Huntley et al. 1983, 1986; Turner and Anderson 1983; Uye and Takamatsu 1990). However, whether or not copepods are able to ingest toxic MTDs is variable (reviewed by Turner and Tester 1997; Colin and Dam 2003). iii) However, a non-toxic strain of *Alexandrium tamarensis*, the toxic dinoflagellates *A. catenella* and *Dinophysis acuminata*, were excellent prey for *Favella* spp. (Stoecker et al. 1981; Maneiro et al. 2000; Kamiyama and Arima 2001). Also, the HTD *Polykrikos kofoidii* exhibited a maximum growth rate on the toxic dinoflagellate *Gymnodinium catenatum* among diverse dinoflagellate prey. In addition, *O. marina* grew well on *Amphidinium carterae* (Jeong et al. 2001a). Therefore, it is concluded that feeding by heterotrophic protistan and metazoan predators on toxic dinoflagellates was affected by

the prey species and strains of a toxic dinoflagellate and also the predator species.

In the ecological aspects, the presence of toxic dinoflagellates may determine the dominant heterotrophic protistan and metazoan predators. The predators which have strong bodies against cell lysis caused by phytotoxins of the toxic dinoflagellates and/or are able to feed on the toxic dinoflagellates (and detoxify the phytotoxins) may dominate zooplankton assemblages. If a predator detoxifies the phytotoxins, the predator plays the role of a sink of phytotoxins in marine food webs. Rapid division of a heterotrophic protistan predator may accelerate the detoxification by attenuating the phytotoxins inside a predator cell. Some copepods are known to transfer phytotoxins sometimes from toxic MTDs to higher level organisms such as fish, seabirds, and marine mammals (e.g. Tester et al. 2000). The copepods may tolerate the toxins during this transfer. However, a heterotrophic protistan predator may not play a role of a transfer of phytotoxins in marine food webs because it has weak tolerance for toxins, cannot detoxify, or cannot survive for a long time without eating prey.

7. Red Tides or Harmful Blooms Dominated by Mixotrophic Dinoflagellates

Many MTDs and some HTDs have been known to cause red tides or harmful blooms (e.g. Jeong 1995). There have been many studies on these red tides since the 19th century. However, for a long time, there have been some unsolved issues in red tides dominated by MTDs; (1) a mechanism of the outbreak and/or the persistence of red tides dominated by some MTDs such as *Karenia brevis*, *Prorocentrum*

Fig. 9. A diagram on a possible mechanism of the outbreak and/or persistence of red tides dominated by mixotrophic dinoflagellates in offshore and/or oceanic waters where inorganic nutrient concentrations are low. Mixotrophic dinoflagellates grow and form a red tide patch by feeding on some cyanobacteria which can conduct nitrogen fixation and on heterotrophic bacteria which usually have high phosphorus contents

donghaiense, and *P. minimum* in offshore and/or oceanic waters where the nutrient concentrations are low have been a mystery to the scientists. (2) Also, a red tide dominated by one MTD species has often been followed by a successive red tide dominated by another MTD species in a short period (a few days). When considering the photogrowth rate of the MTD species, the rapid succession of another MTD species in serial red tides has been difficult to explain. However, the discovery of mixotrophy in phototrophic dinoflagellates may give some clue on these unsolved issues.

By combining the results of Jeong et al. (2005b) and Seong et al. (2006), I propose a possible mechanism of the outbreak and/or the persistence of offshore or oceanic red tides (Fig. 9); many MTDs such as *K. brevis*, *P. donghaiense*, and *P. minimum* are able to feed on *Synechococcus* sp. and heterotrophic bacteria. Therefore, if the MTDs feed on some cyanobacteria which can conduct nitrogen fixation

(e.g. Mitsui et al. 1986) and heterotrophic bacteria which usually have high P : N ratios (e.g. Tezuka 1990), the MTDs are able to obtain nitrogen and phosphorus simultaneously for their growth in offshore or oceanic waters. It is worthwhile to test this hypothesis in offshore and oceanic waters.

Jeong et al. (2005c) suggested that feeding by larger MTDs on smaller MTDs may be a driving force for the succession of dominant species during serial red tides based on their field data; in Masan Bay, Korea in 2004, a bloom dominated by a mixture of *Amphidinium* sp. and the raphidophyte *Heterosigma akashiwo* was followed by one dominated by a mixture of *Prorocentrum minimum* and *P. triestinum*, then by *Cochlodinium polykrikoides*, and then by *P. micans* in series. *P. minimum* and *P. triestinum* which were able to feed on *A. carterae* and *H. akashiwo* were ingested in turn by *C. polykrikoides* and *P. micans* (Jeong et al. 2004c, 2005c). Under these situations, nutrients may not play such an important role in the succession of dominant species during serial red tides; direct transfer of the materials and energy of the organisms that form a red tide into MTDs by predation may not need the uptake and release of nitrogen and phosphorus by the organisms. This may cause uncoupling between nutrient concentrations and MTD abundance in natural environments.

The discovery of mixotrophy in many phototrophic dinoflagellates may change our views of the mechanisms of the outbreak, persistence, and decline of red tides dominated by dinoflagellates, and also of the driving forces for succession in serial red tides. Therefore, to understand the process of red tides, we should consider mixotrophy in dominant dinoflagellates.

8. Marine Planktonic Food Webs and Roles of Dinoflagellates

Establishing the food webs in the marine planktonic community is one of the most important steps in understanding the cycling of materials and flows of energy and predicting further responses of marine ecosystems to diverse-scale-environmental changes. For the last four decades, several new ecological concepts in marine planktonic food webs have been established (Azam et al. 1983). In particular, dinoflagellates have been one of the triggers for discovering new pathways and in turn establishing new ecological concepts in marine ecosystems (Bockstahler and Coats 1993a; Jeong 1994b; Jeong et al. 2005b, 2008; Seong et al. 2006).

Fig. 10. A diagram of marine planktonic food webs focusing on dinoflagellates. Line (1): Feeding by mixotrophic dinoflagellates on bacteria (e.g. Jeong et al. 2005b; Seong et al. 2006). Line (2): Feeding by mixotrophic dinoflagellates on phytoplankton (e.g. Stoecker et al. 1997). Line (3): Feeding by mixotrophic dinoflagellates on other mixotrophic dinoflagellates and mixotrophic nanoflagellates (e.g. Jeong et al. 2005c). Line (4): Feeding by mixotrophic dinoflagellates on heterotrophic protists (e.g. Bockstahler and Coats 1993a). Line (5): Feeding by heterotrophic protists on mixotrophic dinoflagellates (e.g. Jeong and Latz 1994). Line (6): Feeding by metazooplankton on mixotrophic dinoflagellates (e.g. Turner and Tester 1997). Line (7): Feeding by heterotrophic dinoflagellates on bacteria (e.g. Jeong et al. 2008). Line (8): Feeding by heterotrophic dinoflagellates on phytoplankton (Hansen 1992; Jeong et al. 2004d). Line (9): Predator-prey relationships among heterotrophic dinoflagellates (Jeong et al. unpubl. data). Line (10): Feeding by heterotrophic dinoflagellates on heterotrophic protists (Jeong et al. 2007b). Line (11): Feeding by heterotrophic dinoflagellates on metazooplankton including eggs and naupliar stages (Jeong 1994b). Line (12): Feeding by heterotrophic protists on heterotrophic dinoflagellates (Jeong et al. 2004b). Line (13): Feeding by metazooplankton on heterotrophic dinoflagellates (Gifford and Dagg 1991). Are there any exclusively autotrophic nanoflagellate (ANF) or dinoflagellate (ATD)?

(HT bacteria: Heterotrophic bacteria, MTD: Mixotrophic dinoflagellates, HTD: Heterotrophic dinoflagellates, MNF: Mixotrophic nanoflagellate, HNF: Heterotrophic nanoflagellate, PicoEuk: Picoeukaryote)

For the last decade, several models, including mixotrophic protists in marine microbial food webs, have been established (Stoecker 1998; Stickney et al. 2000; Jost et al. 2004; Hammer and Pitchford 2005). However, these models may need to be modified in response to recent discoveries

related to dinoflagellates. For example, the model established by Stickney et al. (2000) needs to include the data on the feeding by large MTDs on bacteria (Nygaard and Tobiesen 1993; Jeong et al. 2005b; Seong et al. 2006). Also, most models mentioned above have not included diverse predator-

Fig. 11. The prey and predators of *Prorocentrum minimum* (A) and *P. micans* (B) so far reported. The blue-colored arrows (routes) emerging from the variety of prey all point to *P. minimum* or *P. micans*, while the red-colored arrows (routes) emerging from *P. minimum* or *P. micans* lead to the diverse predator species. *P. minimum* or *P. micans* box looks like a hub airport and thus we suggest calling each diagram “the *Prorocentrum minimum* hub” or “the *Prorocentrum micans* hub” (BL: Bivalve larvae, CB: Cyanobacteria, CL: Ciliates, CO: Copepods, CP: Cryptophytes, DA: Diatoms, HD: Heterotrophic dinoflagellates, HNF: Heterotrophic nanoflagellates, HP: Haptophytes, HT bacteria: Heterotrophic bacteria, MD: Mixotrophic dinoflagellates, RP: Raphidophytes, PL: Polychaeta larvae, PC: Picoeukaryotes)

prey relationships among MTDs at the species level and allocated only one or few boxes for mixotrophic organisms. The box should be split into many boxes because many MTDs coexist in water parcels, there are many predator-prey relationships among these MTDs, and the ecological niche of a mixotrophic dinoflagellate is different from that of other MTDs (Jeong et al. 2005c). Also, Hammer and Pitchford (2005) established a model to address the effect of mixotrophy in plankton bloom dynamics, mainly focussing on Model III mixotrophic organisms which are called photosynthetic “protozoa” by Stoecker (1998). They assumed that the portion of mixotrophs involving phytoplankton or zooplankton would be small. However, now many have been revealed to be MTDs and also all phototrophic dinoflagellates are suggested to be MTDs. Including large portions of MTDs into Model III mixotrophs may affect their results; mixotrophy may be a large perturbation to the predator-prey system in their model, not a small perturbation.

Based on literature and our recent findings, we established a diagram describing marine planktonic food webs focusing on dinoflagellates (Fig. 10). One of the most interesting things is the fact that the ecological roles of MTDs in marine plankton food webs are quite different from those of other phytoplankton such as diatoms and phototrophic nano- and microflagellates. Also, two diagrams could be drawn to describe predator-prey relationships between each of two different MTD species (e.g. *Prorocentrum minimum* and *P. micans*) and other plankton (Fig. 11). These two diagrams (so called, *Prorocentrum minimum* hub and *P. micans* hub) show that the prey and/or predators of one MTD species are quite different from those of the other MTD species. Therefore, each MTD may have a different ecological niche from that of other MTD species. These different ecological niches and thus they may give an explanation for the phytoplankton co-existence by Hutchinson (1961) and help us to understand the roles of each MTD in marine ecosystems at the species level and also to establish effective ecological models.

As described above, MTDs and HTDs interact with very diverse marine organisms and thus they play diverse roles in marine plankton food webs as follows; (1) in the community level, MTDs and HTDs are effective predators on diverse prey (taxa, size etc.) from bacteria (maybe virus) to metazoans due to their diverse feeding mechanisms. Also, in the species level, a dinoflagellate species is an effective predator on diverse prey items with a wide size

range because it has 2 or more different feeding mechanisms depending on the type and/or size of prey. (2) Dinoflagellates are excellent prey for diverse heterotrophic protistan and metazoan predators. Thus, they sometimes play a role of a link between bacteria and some metazoan predators which are unable to ingest bacteria directly. (3) They are strong competitors with red-tide algae, HNFs, and ciliates for picoplankton and with ciliates and metazoans for nanoplankton and larger prey. (4) They sometimes control the populations of diverse prey and play the role of a driving force for rapid succession of dominant species in serial red tides. Also, grazing impact of dinoflagellates on a prey population is often higher or comparable to that of ciliates or copepods due to higher abundance. (5) Some dinoflagellates are able to detoxify phytotoxin. They may play a role of sinking phytotoxins from marine food webs. (6) Dinoflagellates make food webs long and complicated, but they make limited materials to be utilized in maximum. In particular, many predator-prey relationships among dinoflagellates make food webs more complex but support co-existence of many dinoflagellate species with separation of ecological niches at the level of species.

9. Further Studies

To understand the roles of MTDs and HTDs in marine plankton food webs, it is worth exploring the following questions in future researches: (1) Are there exclusively autotrophic dinoflagellates? So far, approximately 70 phototrophic dinoflagellates have been revealed to be mixotrophic (e.g. Jacobson and Anderson 1996; Stoecker 1999; Jeong et al. 2005c, 2010; Seong et al. 2006). Considering the fact that all the MTDs tested in our recent studies were able to feed on bacteria, and also that the endosymbiosis theory describing the origin of plastids is a widely accepted theory (Bhattacharya and Medlin 1995; Hackett et al. 2004; McFadden 2001; Yoon et al. 2002), other untested dinoflagellates may also be mixotrophic or heterotrophic. It is worthwhile testing the ability of phagotrophy in all phototrophic dinoflagellates not yet tested; (2) Plastids of phototrophic prey organisms belonging to diverse classes such as cyanophyta, cryptophyta, haptophyta, bacillariophyta, dinophyta, can be transferred into MTDs by predation (e.g. Jeong et al. 2005c). Thus, there is a chance that MTDs can contain diverse plastids originating from diverse classes. Feeding by MTDs on diverse

phototrophic organisms having different plastids supports the hypothesis of tertiary endosymbiosis (ancestor of dinoflagellates engulfing a eukaryotic cell with a secondary plastid) and multiple origins of dinoflagellates (Bhattacharya et al. 2003). However, considering the ability of all MTDs tested in our study to feed on cyanobacterium *Synechococcus* (Jeong et al. 2005b), there may be some dinoflagellates containing plastids originating from cyanophyta. To explore evolution in algae, it is worthwhile exploring the relationships between the feeding occurrences, types of plastids, DNA sequences, and gene expression of MTDs and their prey. (3) There may be some predator-prey relationships between MTDs and other marine organisms so far not observed. For example, MTDs may feed on small heterotrophic nanoflagellates, or vice versa. However, this has been not tested yet. The discovery of unknown food webs is critical and serves our understanding of ecology and evolution in phototrophic organisms and increases the validity of ecological models and predictions. (4) DNA difference in dinoflagellates may cause difference in morphology and enzymes involving in feeding and digesting prey. Therefore, it is worthwhile exploring the relationships among phylogeny tree based on DNA sequence, morphology, feeding enzymes, and morphology of dinoflagellates.

Acknowledgements

We thank Jae Yeon Park, Seong Taek Kim, Jae Yoon Song, Jong Hyeok Kim, Soo Kyeom Kim for technical support. This paper was funded by grants from National Research Foundation (2009-0058298) and Ecological Disturbance Research Program, Korea Institute of Marine Science & Technology Promotion/Ministry of Land, Transportation and Marine Affairs (KMLTM) award to HJ Jeong.

References

- Adolf JE, Stoecker DK, Harding LW Jr (2006) The balance of autotrophy and heterotrophy during mixotrophic growth of *Karlodinium micrum* (Dinophyceae). *J Plankton Res* **28**: 737-751
- Adolf JE, Krupatkin D, Bachvaroff T, Place AR (2007) Karlotoxin mediates grazing by *Oxyrrhis marina* on strains of *Karlodinium veneficum*. *Harmful Algae* **6**:400-412
- Azam F, Fenchel T, Field JG, Gray JS, Meyer-Reil L, Thingstad F (1983) The ecological role of water-column microbes in the sea. *Mar Ecol Prog Ser* **10**:257-263
- Azanza RV, Fukuyo Y, Yap LG, Takayama H (2005) *Prorocentrum minimum* bloom and its possible link to a massive fish kill in Bolinao, Pangasinan, Northern Philippines. *Harmful Algae* **4**:519-524
- Baek SH, Shimode J, Kikuchi T (2008) Growth of dinoflagellates, *Ceratium furca* and *Ceratium fusus* in Sagami Bay, Japan: the role of temperature, light intensity and photoperiod. *Harmful algae* **7**:163-173
- Berge T, Hansen PJ, Moestrup Ø (2008) Feeding mechanism, prey specificity and growth in light and dark of the plastidic dinoflagellate *Karlodinium armiger*. *Aquat Microb Ecol* **50**: 279-288
- Berggreen B, Hansen B, Kiørboe T (1988) Food size spectra, ingestion and growth of the copepod *Acartia tonsa* during development: implications for determination of copepod production. *Mar Biol* **99**:341-352
- Besiktepe S, Dam HG (2002) Coupling of ingestion and defecation as a function of diet in the calanoid copepod *Acartia tonsa*. *Mar Ecol Prog Ser* **229**:151-164
- Bhattacharya D, Medlin L (1995) The phylogeny of plastids: a review based on comparisons of small-subunit ribosomal RNA coding regions. *J Phycol* **31**:489-498
- Bhattacharya D, Yoon HS, Hackett JD (2003) Photosynthetic eukaryotic unite: endosymbiosis connects the dots. *BioEssays* **26**:50-60
- Biecheler B (1952) Recherches sur les Peridiniens. *Bull Biol Fr Belg Suppl* **36**:1-149
- Bockstahler KR, Coats DW (1993a) Grazing of the mixotrophic dinoflagellate *Gymnodinium sanguineum* on ciliate population of Chesapeake Bay. *Mar Biol* **116**:447-487
- Bockstahler KR, Coats DW (1993b) Spatial and temporal aspects of mixotrophy in Chesapeake Bay dinoflagellates. *J Eukaryot Microbiol* **40**:49-60
- Boenigk J, Arndt H (2000) Comparative studies on the feeding behavior of two heterotrophic nanoflagellates: the filter-feeding choanoflagellate *Monosiga ovata* and the raptorial-feeding kinetoplastid *Rhynchomonas nasuta*. *Aquat Microb Ecol* **22**:243-249
- Broglio E, Jonasdottir SH, Calbet A, Jakobsen HH, Saiz E (2003) Effect of heterotrophic versus autotrophic food on feeding and reproduction of the calanoid copepod *Acartia tonsa*: relationship with prey fatty acid composition. *Aquat Microb Ecol* **31**:267-278
- Burkholder JAM, Glasgow HB Jr (1995) Interactions of a toxic estuarine dinoflagellate with microbial predators and prey. *Arch Protistenkd* **145**:177-188
- Burkholder JAM, Glasgow HB Jr (1997) Trophic controls on stage transformations of a toxic ambush-predator dinoflagellate. *J Eukaryot Microbiol* **44**:200-205
- Burkholder JAM, Glibert PM, Skelton HM (2008) Mixotrophy, a major mode of nutrition for harmful algal species in eutrophic

- waters. *Harmful Algae* **8**:77-93
- Burkholder JAM, Noga EJ, Hobbs CW, Glasgow HB Jr, Smith SA (1992) New "phantom" dinoflagellate is the causative agent of major estuarine fish kills. *Nature* **358**:407-410
- Bursa AS (1961) The annual oceanographic cycle at Igloolik in the Canadian Arctic II. The Phytoplakton. *J Fish Res Bd Can* **18**:563-615
- Buskey EJ (1997) Behavioral components of feeding selectivity of the heterotrophic dinoflagellate *Protoperdinium pellucidum*. *Mar Ecol Prog Ser* **153**:77-89
- Buskey EJ, Coulter CJ, Brown SL (1994) Feeding, growth and bioluminescence of the heterotrophic dinoflagellate *Protoperdinium huberi*. *Mar Biol* **121**:373-380
- Calbet A, Vaque D, Felipe J, Vila M, Sala MM, Alcaraz M, Estrda M (2003) Relative grazing impact of microzooplankton and mesozooplankton on a bloom of the toxic dinoflagellate *Alexandrium minutum*. *Mar Ecol Prog Ser* **259**:303-309
- Cohen JH, Tester PA, Forward RB Jr (2007) Sublethal effects of the toxic dinoflagellate *Karenia brevis* on marine copepod behavior. *J Plankton Res* **29**:301-315
- Colin SP, Dam HG (2003) Effects of the toxic dinoflagellate *Alexandrium fundyense* on the copepod *Acartia hudsonica*: a test of the mechanisms that reduce ingestion rates. *Mar Ecol Prog Ser* **248**:55-65
- Daugbjerg N, Hansen G, Larsen J, Moestrup Ø (2000) Phylogeny of some major genera of dinoflagellates based on ultrastructure and partial LSU rDNA sequence data, including the erection of 3 new genera of unarmoured dinoflagellates. *Phycologia* **39**:302-317
- ECOHAB (1995) The ecology and oceanography of harmful algal blooms: a national research agenda. Woods Hole Oceanographic Institute, Woods Hole, MA, 66 p
- Feinstein TN, Traslavina R, Sun M, Lin S (2002) Effects of light on photosynthesis, grazing, and population dynamics of the heterotrophic dinoflagellate *Pfiesteria piscicida* (Dinophyceae). *J Phycol* **38**:659-669
- Fenchel T (1987) Ecology of protozoa: the biology of free living phagotrophic protists. Springer-Verlag, New York, 197 p
- Gaines G, Taylor FJR (1984) Extracellular digestion in marine dinoflagellates. *J Plankton Res* **6**:1057-1061
- Garces E, Fernandez M, Penna A, Van Lenning K, Gutierrez A, Camp J, Zapata M (2006) Characterization of MW Mediterranean *Karlodinium* spp. (Dinophyceae) strains using morphological, molecular, chemical, and physiological methodologies. *J Phycol* **42**: 1096-1112
- Gifford DJ, Dagg MJ (1991) The microzooplankton-mesozooplankton link: consumption of planktonic protozoa by the calanoid copepods *Acartia tonsa* Dana and *Neocalanus plumchrus* Murukawa. *Mar Microb Food Webs* **5**:161-177
- Glasgow HB, Lewitus AJ, Burkholder JAM (1998) Feeding behavior of the ichthyotoxic estuarine dinoflagellate, *Pfiesteria piscicida*, on amino acids, algal prey, and fish vs. *Mammalian erythrocytes*. In: Reguera B, Blanco J, Fernandez ML, Wyatt T (eds) *Harmful microalgae*. Intergovernmental Oceanographic Commission of UNESCO, Paris, pp 394-397
- Goldman JC, Dennett MR, Gordin H (1989) Dynamics of herbivorous grazing by the heterotrophic dinoflagellate *Oxyrrhis marina*. *J Plankton Res* **11**:391-407
- Granéli E, Anderson DM, Carlsson P, Maestrini SY (1997) Light and dark carbon uptake by *Dinophysis* species in comparison to other photosynthetic and heterotrophic dinoflagellates. *Aquat Microb Ecol* **13**:177-186
- Hackett JD, Anderson DM, Erdner DL, Bhattacharya D (2004) Dinoflagellates: a remarkable evolutionary experiment. *Am J Bot* **91**:1523-1534
- Hammer A, Pitchford J (2005) The role of mixotrophy in plankton bloom dynamics, and the consequences for system productivity. *J Mar Sci* **62**:833-840
- Hansen FC, Witte HJ, Passarge J (1996) Grazing in the heterotrophic dinoflagellate *Oxyrrhis marina*: size selectivity and preference for calcified *Emiliania huxleyi* cells. *Aquat Microb Ecol* **10**:307-313
- Hansen PJ (1989) The red tide dinoflagellate *Alexandrium tamarense*: effects on behaviour and growth of a tintinnid ciliate. *Mar Ecol Prog Ser* **53**:105-116
- Hansen PJ (1991) Quantitative importance and trophic role of heterotrophic dinoflagellates in a coastal pelagial food web. *Mar Ecol Prog Ser* **73**:253-261
- Hansen PJ (1992) Prey size selection, feeding rates and growth dynamics of heterotrophic dinoflagellates with special emphasis on *Gyrodinium spirale*. *Mar Biol* **114**:327-334
- Hansen PJ (1995) Growth and grazing response of a ciliate feeding on the red tide dinoflagellate *Gyrodinium aureolum* in monoculture and in mixture with a non-toxic alga. *Mar Ecol Prog Ser* **121**:65-72
- Hansen PJ (2002) Effect of high pH on the growth and survival of marine phytoplankton: implications for species succession. *Aquat Microb Ecol* **28**:279-288
- Hansen PJ, Lundholm N, Rost B (2007) Growth limitation in marine red-tide dinoflagellates: effects of pH versus inorganic carbon availability. *Mar Ecol Prog Ser* **334**:63-71
- Hansen PJ, Nielsen TG (1997) Mixotrophic feeding of *Fragilidium subglobosum* (Dinophyceae) on three species of *Ceratium*: effects of prey concentration, prey species and light intensity. *Mar Ecol Prog Ser* **147**:187-196
- Hansen PJ, Calado AJ (1999) Phagotrophic mechanisms and prey selection in free-living dinoflagellates. *J Eukaryot Microbiol* **46**:382-389
- Heinbokel JF, Coats DW, Henderson KW, Tyler MA (1988) Reproduction rates and secondary production of three species of the rotifer genus *Synchaeta* in the estuarine Potomac River. *J Plankton Res* **10**:659-672

- Houde SEL, Roman M (1987) Effects of food quality on the functional ingestion response of the copepod *Acartia tonsa*. Mar Ecol Prog Ser **40**:69-77
- Huntley M, Barthel K-G, Star JL (1983) Particle rejection by *Calanus pacificus*: discrimination between similarly sized particles. Mar Biol **74**:151-160
- Huntley M, Sykes P, Rohan S, Marin V (1986) Chemically-mediated rejection of dinoflagellate prey by the copepods *Calanus pacificus* and *Paracalanus parvus*: mechanism, occurrence and significance. Mar Ecol Prog Ser **28**:105-120
- Huskin I, Anadon R, Ivarez-Marqus F, Harris RP (2000) Ingestion, faecal pellet and egg production rates of *Calanus helgolandicus* feeding coccolithophorid versus non-coccolithophorid diets. J Exp Mar Biol Ecol **248**:239-254
- Hutchinson GE (1961) The paradox of the plankton. Am Nat **95**:137-145
- Jacobson DM (1987) The ecology and feeding biology of thecate heterotrophic dinoflagellates. Ph.D. Thesis, Woods Hole Oceanographic Institution/Massachusetts Institute of Technology Joint Program, 210 p
- Jacobson DM, Anderson DM (1986) Thecate heterotrophic dinoflagellates: feeding behavior and mechanisms. J Phycol **22**:249-258
- Jacobson DM, Andersen RA (1994) The discovery of mixotrophy in photosynthetic species of *Dinophysis* (Dinophyceae): light and electron microscopical observations of food vacuoles in *Dinophysis acuminata*, *D. norvegica* and two heterotrophic dinophysoid dinoflagellates. Phycologia **33**:97-110
- Jacobson DM, Anderson DM (1996) Widespread phagocytosis of ciliates and other protists by marine mixotrophic and heterotrophic thecate dinoflagellates. J Phycol **32**:279-285
- Jakobsen HH, Hansen PJ (1997) Prey size selection, grazing and growth response of the small heterotrophic dinoflagellate *Gymnodinium* sp. and the ciliate *Balanion comatum*-a comparative study. Mar Ecol Prog Ser **158**:75-86
- Jakobsen HH, Hansen PJ, Larsen J (2000) Growth and grazing responses of two chloroplast-retaining dinoflagellates: Effect of irradiance and prey species. Mar Ecol Prog Ser **201**: 121-128
- Jeong HJ (1994a) Predation effects of the calanoid copepod *Acartia tonsa* on a population of the heterotrophic dinoflagellate *Protoperidinium* cf. *divergens* in the presence of co-occurring red-tide dinoflagellate prey. Mar Ecol Prog Ser **111**:87-97
- Jeong HJ (1994b) Predation by the heterotrophic dinoflagellate *Protoperidinium* cf. *divergens* on copepod eggs and early naupliar stages. Mar Ecol Prog Ser **114**:203-208
- Jeong HJ (1995) The interactions between microzooplanktonic grazers and dinoflagellates causing red tides in the open coastal waters off southern California. Ph.D. Thesis, Scripps Institution of Oceanography, University of California, San Diego, 139 p. Available on microfilm from University of Michigan, Accession Number 223882
- Jeong HJ (1999) The ecological roles of heterotrophic dinoflagellates in marine planktonic community. J Eukaryot Microbiol **46**: 390-396
- Jeong HJ, Latz MI (1994) Growth and grazing rates of the heterotrophic dinoflagellate *Protoperidinium* spp. on red tide dinoflagellates. Mar Ecol Prog Ser **106**:173-185
- Jeong HJ, Lee CW, Yih WH, Kim JS (1997) *Fragilidium* cf. *mexicanum*, a thecate mixotrophic dinoflagellate which is prey for and a predator on co-occurring thecate heterotrophic dinoflagellate *Protoperidinium* cf. *divergens*. Mar Ecol Prog Ser **151**:299-305
- Jeong HJ, Shim JH, Kim JS, Park JY, Lee CW, Lee Y (1999a) The feeding by the thecate mixotrophic dinoflagellate *Fragilidium* cf. *mexicanum* on red tide and toxic dinoflagellate. Mar Ecol Prog Ser **176**:263-277
- Jeong HJ, Shim JH, Lee CW, Kim JS, Koh SM (1999b) Growth and grazing rates of the marine planktonic ciliate *Strombidinopsis* sp. on red-tide and toxic dinoflagellates. J Eukaryot Microbiol **46**:69-76
- Jeong HJ, Kang HJ, Shim JH, Park JG, Kim JS, Song JY, Choi HJ (2001a) Interactions among the toxic dinoflagellate *Amphidinium carterae*, the heterotrophic dinoflagellate *Oxyrrhis marina*, and the calanoid copepods *Acartia* spp. Mar Ecol Prog Ser **218**:77-86
- Jeong HJ, Kim SK, Kim JS, Kim ST, Yoo YD, Yoon JY (2001b) Growth and grazing rates of the heterotrophic dinoflagellate *Polykrikos kofoidii* on red-tide and toxic dinoflagellates. J Eukaryot Microbiol **48**:298-308
- Jeong HJ, Yoon JY, Kim JS, Yoo YD, Seong KA (2002) Growth and grazing rates of the prostomatid ciliate *Tiarina fusus* on red-tide and toxic algae. Aquat Microb Ecol **28**:289-297
- Jeong HJ, Kim JS, Yoo YD, Kim ST, Kim TH, Park MG, Lee CH, Seong KA, Kang NS, Shim JH (2003a) Feeding by the heterotrophic dinoflagellate *Oxyrrhis marina* on the red-tide raphidophyte *Heterosigma akashiwo*: a potential biological method to control red tides using mass-cultured grazers. J Eukaryot Microb **50**:274-282
- Jeong HJ, Park KH, Kim JS, Kang HJ, Kim CH, Choi HJ, Kim YS, Park JY, Park MG (2003b) Reduction in the toxicity caused by a toxic dinoflagellate *Gymnodinium catenatum* by the feeding of the heterotrophic dinoflagellate *Polykrikos kofoidii*. Aquat Microb Ecol **31**:307-312
- Jeong HJ, Song JY, Lee CH, Kim ST (2004a) Feeding by the larvae of the mussel *Mytilus galloprovincialis* on red-tide dinoflagellates. J Shellfish Res **23**:185-195
- Jeong HJ, Yoo YD, Kim JS, Kang NS, Kim TH, Kim JH (2004b) Feeding by the marine planktonic ciliate *Strombidinopsis jeokjo* on common heterotrophic dinoflagellates. Aquat Microb Ecol **36**:181-187
- Jeong HJ, Yoo YD, Kim JS, Kim TH, Kim JH, Kang NS, Yih

- WH (2004c) Mixotrophy in the phototrophic harmful alga *Cochlodinium polykrikoides* (Dinophyceae): prey species, the effects of prey concentration and grazing impact. *J Eukaryot Microbiol* **51**:563-569
- Jeong HJ, Yoo YD, Kim ST, Kang NS (2004d) Feeding by the heterotrophic dinoflagellate *Protoperdinium bipes* on the diatom *Skeletonema costatum*. *Aquat Microb Ecol* **36**:171-179
- Jeong HJ, Kim JS, Kim JH, Kim ST, Seong KA, Kim TH, Song JY, Kim SK (2005a) Feeding and grazing impact by the newly described heterotrophic dinoflagellate *Stoeckeria algicida* on the harmful alga *Heterosigma akashiwo*. *Mar Ecol Prog Ser* **295**:69-78
- Jeong HJ, Park JY, Nho JH, Park MO, Ha JH, Seong KA, Jeng C, Seong CN, Lee KY, Yih WH (2005b) Feeding by red-tide dinoflagellates on the cyanobacterium *Synechococcus*. *Aquat Microb Ecol* **41**:131-143
- Jeong HJ, Yoo YD, Park JY, Song JY, Kim ST, Lee SH, Kim KY, Yih WH (2005c) Feeding by the phototrophic red-tide dinoflagellates: five species newly revealed and six species previously known to be mixotrophic. *Aquat Microb Ecol* **40**:133-155
- Jeong HJ, Yoo YD, Seong KA, Kim JH, Park JY, Kim SH, Lee SH, Ha JH, Yih WH (2005d) Feeding by the mixotrophic dinoflagellate *Gonyaulax polygramma*: mechanisms, prey species, the effects of prey concentration, and grazing impact. *Aquat Microb Ecol* **38**:249-257
- Jeong HJ, Ha JH, Park JY, Kim JH, Kang NS, Kim S, Kim JS, Yoo YD, Yih WH (2006) Distribution of the heterotrophic dinoflagellate *Pfiesteria piscicida* in Korean waters and its feeding on mixotrophic dinoflagellates, raphidophytes, and fish blood cells. *Aquat Microb Ecol* **44**:263-275
- Jeong HJ, Ha JH, Yoo YD, Park JY, Kim JH, Kang NS, Kim TH, Kim HS, Yih WH (2007a) Feeding by the *Pfiesteria*-like heterotrophic dinoflagellate *Luciella masanensis*. *J Eukaryot Microbiol* **54**:231-241
- Jeong HJ, Song JE, Kang NS, Kim S, Yoo YD, Park JY (2007b) Feeding by heterotrophic dinoflagellates on the common marine heterotrophic nanoflagellate *Cafeteria* sp. *Mar Ecol Prog Ser* **333**:151-160
- Jeong HJ, Kim JS, Song JY, Kim JH, Kim TH, Kim SK, Kang NS (2007c) Feeding by heterotrophic protists and copepods on the heterotrophic dinoflagellates *Pfiesteria piscicida*, *Stoeckeria algicida*, and *Luciella masanensis*. *Mar Ecol Prog Ser* **349**:199-211
- Jeong HJ, Seong KA, Yoo YD, Kim TH, Kang NS, Kim S, Park JY, Kim JS, Kim GH, Song JY (2008) Feeding and grazing impact by small marine heterotrophic dinoflagellates on heterotrophic bacteria. *J Eukaryot Microbiol* **55**:271-288
- Jeong HJ, Yoo YD, Kang NS, Rho JR, Seong KA, Park JW, Nam GS, Yih WH (2010) Ecology of *Gymnodinium aureolum*. I. Feeding in western Korean water. *Aquat Microb Ecol* **59**:239-255
- Jost C, Lawrence CA, Campolongo F, van de Bund W, Hill S, DeAngelis DL (2004) The effects of mixotrophy on the stability and dynamics of a simple planktonic food web model. *Theoret Popul Biol* **66**:37-51
- Kamiyama T, Arima S (2001) Feeding characteristics of two tintinnid ciliate species on phytoplankton including harmful species: effects of prey size on ingestion rates and selectivity. *J Exp Mar Biol Ecol* **257**:281-296
- Kang NS, Jeong HJ, Moestrup Ø, Shin WG, Nam SW, Park JY, de Salas MF, Kim KW, Noh JH (2010) Description of a new planktonic mixotrophic dinoflagellate *Paragymnodinium shiwhaense* n. gen., n. sp. from the coastal waters off western Korea: morphology, pigments, and ribosomal DNA gene sequence. *J Eukaryot Microbiol* **57**:121-144
- Kim JS, Jeong HJ (2004) Feeding by the heterotrophic dinoflagellates *Gyrodinium dominans* and *G. spirale* on the red-tide dinoflagellate *Prorocentrum minimum*. *Mar Ecol Prog Ser* **280**:85-94
- Kondo K, Seike Y, Date Y (1990) Red tides in the brackish Lake Nakanoumi. (II). Relationships between the occurrence of *Prorocentrum minimum* red tide and environmental conditions. *Bull Plankt Soc Japan Hiroshima* **37**:19-34
- Koski M, Riser CW (2006) Post-bloom feeding of *Calanus finmarchicus* copepodites: selection for autotrophic versus heterotrophic prey. *Mar Biol Res* **2**:109-119
- Larsen J (1988) An ultrastructural study of *Amphidinium poecilochroum* (Dinophyceae), a phagotrophic dinoflagellate feeding on small species of cryptophytes. *Phycologia* **27**:366-377
- Lee CW (1998) Growth and grazing rates of the heterotrophic dinoflagellate *Oxyrrhis marina* and the ciliate *Strombidinopsis* sp. on *Prorocentrum* spp. M.A. Thesis, Kunsan National University, 36 p (In Korean with English abstract)
- Lee SH (2006) Feeding by mixotrophic red-tide algae on photosynthetic picoeukaryotes. M.A. Thesis, Seoul National University, 55 p (In Korean with English abstract)
- Legrand C, Graneli E, Carlsson P (1998) Induced phagotrophy in the photosynthetic dinoflagellate *Heterocapsa triquetra*. *Aquat Microb Ecol* **15**:65-75
- Lessard EJ (1984) Oceanic heterotrophic dinoflagellates: distribution, abundance and role as microzooplankton. Ph.D. Thesis, University of Rhode Island, Kingstown, 166 p
- Lessard EJ, Swift E (1985) Species-specific grazing rates of heterotrophic dinoflagellates in oceanic waters, measured with a dual-label radioisotope technique. *Mar Biol* **87**:289-296
- Li A, Stoecker DK, Coats DW, Adam EJ (1996) Ingestion of fluorescently-labeled and phycoerythrin-containing prey by photosynthetic dinoflagellates. *Aquat Microb Ecol* **10**:139-147
- Li A, Stoecker DK, Coats DW (2000) Mixotrophy in *Gyrodinium*

- galatheanum* (dinophyceae): grazing responses to light intensity and inorganic nutrients. *J Phycol* **36**:33-45
- Liu S, Wang W-X (2002) Feeding and reproductive responses of marine copepods in South China Sea to toxic and nontoxic phytoplankton. *Mar Biol* **140**:595-603
- Maneiro I, Frangopulos M, Guisande C, Fernandez M, Reguera B, Riveiro I (2000) Zooplankton as a potential vector of diarrhetic shellfish poisoning toxins through the food web. *Mar Ecol Prog Ser* **201**:155-163
- Mason PL, Litaker RW, Jeong HJ, Ha JH, Reece KS, Vogelbein WK, Stokes NA, Park JY, Steidinger KA, Vandersea MW, Kibler S, Tester PA, Vogelbein WK (2007) Description of a new genus of *Pfiesteria*-like dinoflagellate, *Luciella* gen. nov. (dinophyceae), including two new species: *Luciella masanensis* sp. nov. and *Luciella atlantis* sp. nov. *J Phycol* **43**:799-810
- Matsubara T, Nagasoe S, Amasaki Y, Shikata T, Shimasaki Y, Oshima Y, Honjo T (2007) Effects of temperature, salinity, and irradiance on the growth of the dinoflagellate *Akashiwo sanguinea*. *J Exp Mar Biol Ecol* **342**:226-230
- McFadden GI (2001) Primary and secondary endosymbiosis and the origin of plastids. *J Phycol* **37**:951-959
- Menden-Deuer S, Lessard EJ, Sattergerg J, Grnbaum D (2005) Growth rates and starvation survival of three species of the pallium-feeding, thecate dinoflagellate genus *Protoperidinium*. *Aquat Microb Ecol* **41**:145-152
- Mitsui A, Cao S, Takahashi A, Arai T (1986) Growth synchrony and cellular parameters of the unicellular nitrogen-fixing marine cyanobacterium, *Synechococcus* sp. strain Miami BG 043511 under continuous illumination. *Physiol Plant* **69**:1-8
- Nagasoe S, Kim DI, Shimasaki Y, Oshima Y, Yamaguchi M, Honjo T (2006) Effects of temperature, salinity and irradiance on the growth of the red tide dinoflagellate *Gyrodinium instriatum* Freudenthal et Lee. *Harmful algae* **5**:20-25
- Nakamura Y, Suzuki S, Hiromi J (1995) Population dynamics of heterotrophic dinoflagellates during a *Gymnodinium mikimotoi* red tide in the Seto Inland Sea. *Mar Ecol Prog Ser* **125**:269-277
- Nakamura Y, Yamazaki Y, Hiromi J (1992) Growth and grazing of a heterotrophic dinoflagellate, *Gyrodinium dominans*, feeding on a red tide flagellate, *Chattonella antiqua*. *Mar Ecol Prog Ser* **82**:275-279
- Naustvoll L-J (1998) Growth and grazing by the thecate heterotrophic dinoflagellate *Diplopsalis lenticula* (Diplopsalidaceae, Dinophyceae). *Phycologia* **37**:1-9
- Naustvoll L-J (2000) Prey size spectra and food preferences in thecate heterotrophic dinoflagellates. *Phycologia* **39**:187-198
- Navarro JM, Muoz MG, Contreras AM (2006) Temperature as a factor regulating growth and toxin content in the dinoflagellate *Alexandrium catenella*. *Harmful algae* **5**:762-769
- Nygaard K, Tobiesen A (1993) Bacterivory in algae: a survival strategy during nutrient limitation. *Limnol Oceanogr* **38**:273-279
- Park MG, Kim SJ, Kim HS, Myung GO, Kang IG, Yih WH (2006) First successful culture of the marine dinoflagellate *Dinophysis acuminata*. *Aquat Microb Ecol* **45**:101-106
- Parrow MW, Glasgow HB, Burkholder JM, Zhang C (2001) Comparative response to algal prey by *Pfiesteria piscicida*, *Pfiesteria shumwayae* and an estuarine 'lookalike' species. In: Hallegraeff GM, Blackburn S, Bolch C, Lewis R (eds) Intergovernmental Oceanographic Commission of UNESCO, Paris, pp 101-104
- Radi T, Pospelova V, de Vernal A, Vaughn Barrie J (2007) Dinoflagellate cysts as indicators of water quality and productivity in British Columbia estuarine environments. *Mar Micropaleontol* **62**:269-297
- Richardson TL, Pinckney JL, Walker EA, Marshalonis DM (2006) Photopigment radiolabelling as a tool for determining *in situ* growth rates of the toxic dinoflagellate *Karenia brevis* (Dinophyceae) *Eur J Phycol* **41**:415-423
- Roman M, Reaugh M, Zhang X (2006) Ingestion of the dinoflagellate, *Pfiesteria piscicida*, by the calanoid copepod, *Acartia tonsa*. *Harmful algae* **5**:435-441
- Rublee PA, Allen C, Schaefer Rhodes EL, Adamson J, Lapworth C, Burkholder J, Glasgow H (2004) Global distribution of toxic *Pfiesteria* complex species detected by PCR assay. In: Steidinger KA, Landsberg JH, Tomas CR, Vargo G (eds) Harmful algae 2002. Intergovernmental Oceanographic Commission of UNESCO, Paris, pp 320-322
- Scura ED, Jerde C (1977) Various species of phytoplankton as food for larval northern anchovy, *Engraulis mordax*, and relative nutritional values of the dinoflagellate *Gymnodinium splendens* and *Gonyaulax polyedra*. *Fish Bull* **75**:577-583
- Seong KA, Jeong HJ, Kim S, Kim GH, Kang JH (2006) Bacterivory by co-occurring red-tide algae, heterotrophic nanoflagellates, and ciliates on marine bacteria. *Mar Ecol Prog Ser* **322**:85-97
- Sherr EB, Sherr BF (2002) Significance of predation by protists in aquatic microbial food webs. *Ant van Leeuwenh* **81**:293-308
- Siano R, Montresor M (2005) Morphology, ultrastructure and feeding behaviour of *Protoperidinium vorax* sp. nov. (Dinophyceae, Peridinales). *Eur J Phycol* **40**:221-232
- Skovgaard A (1996a) Engulfment of *Ceratium* spp. (Dinophyceae) by the thecate photosynthetic dinoflagellate *Fragilidium subglobosum*. *Phycologia* **35**:490-499
- Skovgaard A (1996b) Mixotrophy in *Fragilidium subglobosum* (Dinophyceae): growth and grazing responses as functions of light intensity. *Mar Ecol Prog Ser* **143**:247-253
- Skovgaard A (1998) Role of chloroplast retention in a marine dinoflagellate. *Aquat Microb Ecol* **15**:293-301
- Skovgaard A (2000) A phagotrophically derivable growth factor in the plastidic dinoflagellate *Gyrodinium resplendens*

- (Dinophyceae). *J Phycol* **36**:1069-1078
- Skovgaard A, Hansen PJ, Stoecker DK (2000) Physiology of the mixotrophic dinoflagellate *Fragilidium subglobosum*. 1. Effects of phagotrophy and irradiance on photosynthesis and carbon content. *Mar Ecol Prog Ser* **201**:129-136
- Sleigh MA (1989) Protozoa and other protists. Edward Arnold, London, 342 p
- Smalley GW, Coats DW (2002) Ecology of the red-tide dinoflagellate *Ceratium furca*: distribution, mixotrophy, and grazing impact on ciliate populations of Chesapeake Bay. *J Eukaryot Microbiol* **49**:63-73
- Smalley GW, Coats DW, Adam EJ (1999) A new method using fluorescent microspheres to determine grazing on ciliates by the mixotrophic dinoflagellate *Ceratium furca*. *Aquat Microb Ecol* **17**:167-179
- Smalley GW, Coats DW, Stoecker DK (2003) Feeding in the mixotrophic dinoflagellate *Ceratium furca* is influenced by intracellular nutrient concentrations. *Mar Ecol Prog Ser* **262**:137-151
- Smayda TJ (1997) Harmful algal blooms: their ecophysiology and general relevance to phytoplankton blooms in the sea. *Limnol Oceanogr* **42**:1137-1153
- Steidinger KA, Burkholder JM, Glasgow HB Jr, Hobbs CW, Garrett JK, Truby EW, Noga EJ, Smith SA (1996) *Pfiesteria piscicida* gen. et sp. nov. (Pfiesteriaceae fam. nov.), a new toxic dinoflagellate with a complex life cycle and behavior. *J Phycol* **32**:157-164
- Stickney HL, Hood RR, Stoecker DK (2000) The impact of mixotrophy on planktonic trophic dynamics in marine ecosystems. *Ecol Model* **125**:203-230
- Stoecker DK (1998) Conceptual models of mixotrophy in planktonic protists and some ecological and evolutionary implications. *Eur J Protistol* **34**:281-290
- Stoecker DK (1999) Mixotrophy among dinoflagellates. *J Eukaryot Microbiol* **46**:397-401
- Stoecker DK, Evans GT (1985) Effects of protozoan herbivory and carnivory in a microplankton food web. *Mar Ecol Prog* **25**:159-167
- Stoecker DK, Guillard RRL, Kavee RM (1981) Selective predation by *Favella ehrenbergii* (Tintinnida) on and among dinoflagellates. *Biol Bull* **160**:136-145
- Stoecker DK, Li A, Coats DW, Gustafson DE, Nannen MK (1997) Mixotrophy in the dinoflagellate *Prorocentrum minimum*. *Mar Ecol Prog Ser* **152**:1-12
- Stoecker DK, Parrow MW, Burkholder JM, Glasgow HB Jr (2002) Grazing by microzooplankton on *Pfiesteria piscicida* cultures with different histories of toxicity. *Aquat Microb Ecol* **28**:79-85
- Strom SL (1991) Growth and grazing rates of the herbivorous dinoflagellate *Gymnodinium* sp. from the open subarctic Pacific Ocean. *Mar Ecol Prog Ser* **78**:103-113
- Strom SL, Buskey EJ (1993) Feeding, growth, and behavior of the thecate heterotrophic dinoflagellate *Oblea rotunda*. *Limnol Oceanogr* **38**:965-977
- Tester PA, Turner JT, Shea D (2000) Vectorial transport of toxins from the dinoflagellate *Gymnodinium breve* through copepods to fish. *J Plankton Res* **22**:47-62
- Tezuka Y (1990) Bacterial regeneration of ammonium and phosphate as affected by the carbon: nitrogen: phosphorus ratio of organic substrates. *Microb Ecol* **19**:227-238
- Tillmann U (2004) Interactions between planktonic microalgae and protozoan grazers. *J Eukaryot Microbiol* **51**:156-168
- Tillmann U, John U (2002) Toxic effects of *Alexandrium* spp. on heterotrophic dinoflagellates: an allelochemical defence mechanism independent of PSP-toxin content. *Mar Ecol Prog Ser* **230**:47-58
- Turner JT, Anderson DM (1983) Zooplankton grazing during dinoflagellate blooms in a Cape Cod embayment, with observations of predation upon tintinnids by copepods. *Mar Ecol* **4**:359-374
- Turner JT, Tester PA (1997) Toxic marine phytoplankton, zooplankton grazers, and pelagic food webs. *Limnol Oceanogr* **42**:1203-1214
- Uchida T, Kamiyama T, Matsuyama Y (1997) Predation by a photosynthetic dinoflagellate *Gyrodinium instriatum* on loricated ciliates. *J Plankton Res* **19**:603-608
- Uye SI, Takamatsu K (1990) Feeding interactions between planktonic copepods and red-tide flagellates from Japanese coastal waters. *Mar Ecol Prog Ser* **59**:97-107
- Verity PG, Paffenhöfer G-A (1996) On assessment of prey ingestion by copepods. *J Plankton Res* **18**:1767-1779
- Watanabe MM, Suda S, Inouye I, Sawaguchi T, Chihara M (1990) *Lepidodinium viride* gen. et sp. nov. (Gymnodinales, Dinophyta), a green dinoflagellate with a chlorophyll a- and b-containing endosymbiont. *J Phycol* **26**:741-751
- Watras CJ, Garcon VC, Olson RJ, Chisholm SW, Anderson DM (1985) The effect of zooplankton grazing on estuarine blooms of the toxic dinoflagellate *Gonyaulax tamarensis*. *J Plankton Res* **7**:891-908
- Yamaguchi M, Shigeru I, Nagasaki K, Matsuyama Y, Uchida T, Imai I (1997) Effects of temperature and salinity on the growth of the red tide flagellates *Heterocapsa circularisquama* (Dinophyceae) and *Chattonella verruculosa* (Raphidophyceae). *J Plankton Res* **19**:1167-1174
- Yamamoto T, Tarutani K (1997) Effects of temperature, salinity and irradiance on the growth of toxic dinoflagellate *Alexandrium tamarense* isolated from Hiroshima bay, Japan. *Japanese J Phycol* **45**:95-101 (In Japanese with English abstract)
- Yoo YD, Jeong HJ, Kim MS, Kang NS, Song JY, Shin WG, Kim KY, Lee KT (2009) Feeding by phototrophic red-tide dinoflagellates on the ubiquitous marine diatom *Skeletonema*

- costatum*. J Eukaryot Microbiol **56**:413-420
- Yoo YD, Jeong HJ, Kang NS, Song JY, Kim KY, Lee KT, Kim JH (2010) Feeding by the newly described mixotrophic dinoflagellate *Paragymnodinium shiwhaense*: feeding mechanism, prey species, and effect of prey concentration. J Eukaryot Microbiol **57**:145-158
- Yoon HS, Hackett JD, Pinto G, Bhattacharya D (2002) The single, ancient origin of chromist plastids. Proc Natl Acad Sci USA **99**:15507-15512