The CLEAN project in the context of CO₂ storage and enhanced gas recovery

Michael Kühn · Uwe-Jens Görke · Jens T. Birkholzer ·
Olaf Kolditz

Received: 12 June 2012 / Accepted: 14 June 2012 / Published online: 7 July 2012
© Springer-Verlag 2012

Anthropogenic emissions of carbon dioxide (CO₂) into the atmosphere have a significant impact on the Earth's carbon cycle. While efforts are made to reduce the release of greenhouse gases via reduced energy consumption, more efficient energy production, and a shift to renewable energy supplies, it is generally expected that fossil fuels will continue to provide a major part of the world's energy portfolio during the twenty-first century (IPCC 2005). This is particularly true for coal which is relatively inexpensive and abundantly available in existing or emerging industrial power houses such as the United States, Brazil, Russia, India, and China. Carbon Capture and Storage (CCS) has been developed as an interim measure to allow energy production from coal without CO₂ emissions. This currently available technology entails capturing CO₂ from large-scale industrial sources such as coal-fired power plants or cement, steel and petrochemical factories. The captured CO₂ gas is then compressed to a smaller volume, hence higher density, transported in pipelines, and eventually injected for long-term storage into deep geologic

formations, such as deep saltwater-bearing aquifers or depleted oil and gas fields (IPCC 2005).

Extensive research is currently carried out to (1) improve the technology and efficiency of capturing CO₂ and (2) to ensure the long-term safety of storing CO₂ underground. This volume of "*Environmental Earth Sciences*" is a *topical issue* (TI) focusing on the geologic storage aspect of CCS. The idea of a TI as a new type of publication within this series to assemble manuscripts dealing with a specific topic comprising its state-of-the art, providing recent research results, and discussing future work from an international perspective. This TI is dedicated to the recent progress made in Germany's CO₂ storage R&D programme GEOTECHNOLOGIEN funded by the Federal Ministry of Education and Research (BMBF) and the German Research Foundation (DFG) (GEOTECHNOLOGIEN 2006, 2007, 2009). A specific focus therein was the joint research project CLEAN—CO₂ large-scale enhanced gas recovery in the Altmark natural gas field (Germany), which was conducted by 16 German institutions from academia and industry within the period from July 2008 until December 2011 (Kühn and Münch 2012; Kühn et al. 2011, 2012). In order to place the findings into a broader context, the TI also covers other related CO₂ storage R&D in Germany and provides an international point of view with contributions from the United States (Mukhopadhyay et al. 2012), Korea (Park et al. 2012) and Norway (Mykkeltvedt and Nordbotten 2012). The topical issue contributions are structured into four major thematic areas: (1) Introduction of Joint Initiatives for Field-Based Research; (2) Laboratory Experiments and Field Tests; (3) Monitoring Concepts, and (4) Modelling Studies.

Under the topic Introduction of Joint Initiatives for Field-Based Research, the current issue collates research

M. Kühn (✉)
Helmholtz Centre Potsdam, GFZ, German Research Centre
for Geosciences, Telegrafenberg, 14473 Potsdam, Germany
e-mail: michael.kuehn@gfz-potsdam.de

U.-J. Görke · O. Kolditz
Helmholtz Centre for Environmental Research, UFZ,
Permoserstraße 15, 04318 Leipzig, Germany
e-mail: olaf.kolditz@ufz.de

J. T. Birkholzer
Earth Sciences Division, Lawrence Berkeley National
Laboratory, One Cyclotron Road, Berkeley, CA 94720, USA

O. Kolditz
TU Dresden, Helmholtzstraße 10, 01062 Dresden, Germany

results from several CO₂ joint research projects, directly connected to test sites. Most of the material is provided by the CLEAN project (Kühn et al. 2012) which aimed to inject around 100,000 t of CO₂ into the depleted gas field in the Altmark, Germany. Another pilot site on the same scale is “Europe’s longest-operating on-shore CO₂ storage site at Ketzin, Germany” (Schilling et al. 2009, Würdemann et al. 2010) which is presented in this TI by Martens et al. (2012). A small scale injection test was performed to study the geochemical impact of CO₂ on shallow groundwater through an injection test in Northeast Germany (Peter et al. 2012). A virtual test site is modelled in the CO₂-MoPa project, which concentrates on basic research independent from a specific test site in order to develop generic theoretical and experimental methods (Bauer et al. 2012).

Laboratory Experiments and Field Tests are a integral part to deepen the understanding within the framework of CO₂ storage and enhanced gas recovery (EGR). Pudlo et al. (2012) show results about “the impact of diagenetic fluid-rock reactions on Rotliegend sandstone composition and petrophysical properties (Altmark area, central Germany)”, Huq et al. (2012) investigate “chemical changes in fluid composition due to CO₂ injection in the Altmark gas field” and Hou et al. (2012a) present a developed long-term wellbore sealing concept which was tested in situ in the Altmark natural gas field.

A major factor for the success of any CO₂ storage project is that the subsurface response to CO₂ injection and storage can be reliably and efficiently monitored. This TI introduces several studies describing development and application of Monitoring Concepts. Schütze et al. (2012) study natural analogues to develop reliable monitoring methods to understand subsurface CO₂ migration processes. Park et al. (2012) present “a pressure-monitoring method to warn CO₂ leakage in geological storage sites”. Lempp et al. (2012) outline “methodological approaches in the laboratory with respect to the in situ conditions of the Altmark gas field”. Lamert et al. (2012) test the “feasibility of geoelectrical monitoring and multiphase modelling for process understanding of gaseous CO₂ injection into a shallow aquifer”. Baumann and Henningses (2012) focus on well logging for injection and saturation profiling with special emphasis on the CO₂ injection in depleted gas fields. Myrtilinen et al. (2012) conclude this section with their work on stable isotope applications with regard to CCS and EGR.

Modelling Studies are an important tool to improve the fundamental understanding of subsurface processes related to CO₂ storage but also to support more practical investigations of CO₂ migration, risk assessment, and reservoir management. The challenges for modelling lie in both describing the complexity of coupled thermo-hydro-mechanical–chemical processes and capturing the

structural geology and heterogeneity of real systems (Nordbotten and Celia 2012). First step for any dynamic simulation model is the development of a static geological model. With regard to the CLEAN project, Norden et al. (2012) outline the basis to do that, the “geological and thermal structure of the larger Altensalzwedel area”. Böttcher et al. (2012) evaluate in this TI the thermal equations of state for CO₂ in numerical simulations. Singh et al. (2012a) conducted a “thermal analysis of the Altmark gas field for carbon dioxide injection with enhanced gas recovery”. Hydraulic processes are the focus of Mykkeltvedt and Nordbotten (2012) as they study convective mixing in response to commercial-scale injection. Singh et al. (2012b) describe “numerical simulation of tracer transport in the Altmark gas field”. Hou et al. (2012b) focus their modelling on mechanical processes with special emphasis on enhanced gas recovery technology. The chemical processes of CO₂-induced fluid-rock interactions are evaluated in the modelling studies by De Lucia et al. (2012) and Beyer et al. (2012).

Benchmarking by code or model comparison is an important strategy for demonstrating the accuracy and predictive capability of models. Benchmarks with special emphasis on CO₂ storage have been discussed by Class et al. (2009) and Kempka et al. (2010). In this topical issue, Kolditz et al. (2012a) suggest “a systematic benchmarking approach for geologic CO₂ injection and storage” and Mukhopadhyay et al. (2012) present “a model comparison initiative for a CO₂ injection field test”. In order to better deal with the complexity of simulating coupled processes in complex systems, an on-going effort is to develop modelling platforms allowing efficient cooperation of distributed developer groups (Flemisch et al. 2011, Kalbacher et al. 2012). Here, Kolditz et al. (2012b) present one such effort introducing “OpenGeoSys: an open-source initiative for numerical simulation of thermo-hydro-mechanical/chemical (THM/C) processes in porous media”.

Acknowledgments We owe many thanks to the authors of this topical issue. These include participants of the CLEAN project who contributed to a large extent to this volume. Furthermore we acknowledge manuscripts from other projects in Europe and the world and the additional input from partners around the globe.

References

- Bauer S, Class H, Ebert M, Feeser V, Goetze H, Holzheid A, Kolditz O, Rosenbaum S, Rabbel W, Schaefer D, Dahmke A (2012) Modeling and parameterization of CO₂ storage in deep saline formations for dimension and risk analyses—the CO₂-MoPa project. *Environ Earth Sci*. doi:10.1007/s12665-012-1707-y
- Baumann G, Henningses J (2012) Well logging for injection and saturation profiling during CO₂ injection in depleted gas fields. *Environ Earth Sci*. doi:10.1007/s12665-012-1708-x

- Beyer C, Li D, de Lucia M, Kühn M, Bauer S (2012) Modelling CO₂-induced fluid-rock interactions in the Altensalzswedel gas reservoir. Part II—coupled reactive transport simulation. *Environ Earth Sci*. doi:10.1007/s12665-012-1684-1
- Böttcher N, Taron J, Kolditz O, Park CH, Liedl R (2012) Evaluation of thermal equations of state for CO₂ in numerical simulations. *Environ Earth Sci*. doi:10.1007/s12665-012-1704-1
- Class H, Ebigo A, Helmig R, Dahle HK, Nordbotten JM, Celia MA, Audigane P, Darcis M, Ennis-King J, Fan Y, Flemisch B, Gasda SE, Jin M, Krug S, Labregere D, Naderi Beni A, Pawar RJ, Sbai A, Thomas SG, Trenty L, Wei L (2009) A benchmark study on problems related to CO₂ storage in geologic formations: summary and discussion of the results. *Comput Geosci* 13(4): 409–434. doi:10.1007/s10596-009-9146-x
- De Lucia M, Albrecht D, Bauer S, Beyer C, Kühn M, Nowak T, Pudlo D, Stadler S (2012) Modelling CO₂-induced fluid-rock interactions in the Altensalzswedel gas reservoir. Part I—From experimental data to a reference geochemical model. *Environ Earth Sci*. doi:10.1007/s12665-012-1725-9
- Flemisch B, Darcis M, Erbertseder K, Faigle B, Lauser A, Mosthaf K, Müthing S, Nuske P, Tatomir A, Wolff M, Helmig R (2011) DuMu^x: DUNE for Multi-{phase, component, scale, physics, ...} flow and transport in porous media. *Adv Water Resour* 34(9):1102–1112. doi:10.1016/j.advwatres.2011.03.007
- GEOTECHNOLOGIEN (2006) Investigation, Utilization and protection of the underground kick-off-meeting, BGR Hannover, 22–23 September 2005. Programme & Abstracts Koordinierungsbüro GEOTECHNOLOGIEN, Science Report No. 6, p 144, ISSN: 1619–7399. <http://www.geotechnologien.de>
- GEOTECHNOLOGIEN (2007) 1st French–German Symposium on geological storage of CO₂—abstracts from the CO₂-Symposium in Potsdam, June 21st and 22nd, 2007. Koordinierungsbüro GEOTECHNOLOGIEN, Science Report No. 9, p 202, ISSN: 1619–7399. <http://www.geotechnologien.de>
- GEOTECHNOLOGIEN (2009) Die dauerhafte geologische Speicherung von CO₂ in Deutschland—Aktuelle Forschungsergebnisse und Perspektiven. Sonderband Koordinierungsbüro GEOTECHNOLOGIEN, Science Report No. 14, p 141, ISSN: 1619–7399. <http://www.geotechnologien.de>
- Hou Z, Gou Y, Taron J, Görke UJ, Kolditz O (2012a) Thermo-hydro-mechanical modeling of carbon dioxide injection for enhanced gas-recovery (CO₂-EGR): a benchmarking study for code comparison. *Environ Earth Sci*. doi:10.1007/s12665-012-1703-2
- Hou Z, Wundram L, Meyer R, Schmidt M, Schmitz S, Were P (2012b) Development of a long-term wellbore sealing concept based on numerical simulations and in situ-testing in the Altmark natural gas field. *Environ Earth Sci*. doi:10.1007/s12665-012-1670-7
- Huq F, Blum P, Marks M, Nowak M, Haderlein SB, Grathwohl P (2012) Chemical changes in fluid composition due to CO₂ injection in the Altmark Gas Field: preliminary results from batch experiments. *Environ Earth Sci*. doi:10.1007/s12665-012-1687-y
- IPCC (2005) In: Metz B, Davidson O, de Coninck HC, Loos M, Meyer LA (eds) Special report on carbon dioxide capture and storage prepared by working group III of the intergovernmental panel on climate change. Cambridge University Press, Cambridge, p 442
- Kalbacher T, Delfs JO, Shao H, Wang W, Walther M, Samaniego L, Schneider C, Musloff A, Centler F, Sun F, Hildebrandt A, Liedl R, Borchardt D, Krebs P, Kolditz O (2012) The IWAS-ToolBox: software coupling for an integrated water resources management. *Environ Earth Sci* 65(5):1367–1380. doi:10.1007/s12665-011-1270-y
- Kempka T, Kühn M, Class H, Frykman P, Kopp A, Nielsen CM, Probst P (2010) Modelling of CO₂ arrival time at Ketzin—part I. *Int J Greenhouse Gas Control* 4(6):1007–1015. doi:10.1016/j.ijggc.2010.07.005
- Kolditz O, Bauer S, Beyer C, Böttcher N, Dietrich P, Görke U-J, Kalbacher T, Park C-H, Sauer U, Schütze C, Shao HB, Singh AK, Taron J, Wang W, Watanabe N (2012a) A systematic benchmarking approach for geologic CO₂ injection and storage. *Environ Earth Sci*. doi:10.1007/s12665-012-1656-5
- Kolditz O, Bauer S, Bilke L, Böttcher N, Delfs JO, Fischer T, Görke UJ, Kalbacher T, Kosakowski G, McDermott CI, Park CH, Radu F, Rink K, Shao H, Shao HB, Sun F, Sun YY, Singh AK, Taron J, Walther M, Wang W, Watanabe N, Wu N, Xie M, Xu W, Zehner B (2012b) OpenGeoSys: an open-source initiative for numerical simulation of thermo-hydro-mechanical/chemical (THM/C) processes in porous media. *Environ Earth Sci*. doi:10.1007/s12665-012-1546-x
- Kühn M, Münch U (eds) (2012) CLEAN: CO₂ large-scale enhanced gas recovery. GEOTECHNOLOGIEN Science Report No. 19. Series: Advanced Technologies in Earth Sciences, p 199, ISBN 978-3-642-31676-0 (in press)
- Kühn M, Förster A, Großmann J, Meyer R, Reinicke K, Schäfer D, Wendel H (2011) CLEAN: preparing for a CO₂-based enhanced gas recovery in a depleted gas field in Germany. *Energy Procedia* 4:5520–5526. doi:10.1016/j.egypro.2011.02.538
- Kühn M, Tesmer M, Pilz P, Meyer R, Reinicke K, Förster A, Kolditz O, Schäfer D, CLEAN Partners (2012) CLEAN: project overview on CO₂ large-scale enhanced gas recovery in the Altmark natural gas field (Germany). *Environ Earth Sci*. doi:10.1007/s12665-012-1714-z
- Lamert H, Geistlinger H, Werban U, Schütze C, Peter A, Hornbruch G, Schulz A, Pohlert M, Kalia S, Beyer M, Großmann J, Dahmke A, Dietrich P (2012) Feasibility of geoelectrical monitoring and multiphase modelling for process understanding of gaseous CO₂ injection into a shallow aquifer. *Environ Earth Sci*. doi:10.1007/s12665-012-1669-0
- Lempp CFH, Shams KM, Jahr N (2012) Stress monitoring in deep boreholes as a task in future CCS projects: Methodological approaches in the laboratory with respect to the in situ conditions of the Altmark Gas Field. *Environ Earth Sci*. doi:10.1007/s12665-012-1706-z
- Martens S, Kempka T, Liebscher A, Lüth S, Möller F, Myrntinen A, Norden B, Schmidt-Hattenberger C, Zimmer M, Kühn M (2012) Europe's longest-operating on-shore CO₂ storage site at Ketzin, Germany: a progress report after three years of injection. *Environ Earth Sci*. doi:10.1007/s12665-012-1672-5
- Mukhopadhyay S, Birkholzer JT, Nicot JP, Hosseini SA (2012) A single site multi-model comparative study for CO₂ injection field test: An introduction to Sim-SEQ. *Environ Earth Sci*. doi:10.1007/s12665-012-1668-1
- Mykkeltvedt TS, Nordbotten JM (2012) Representing effective rates of convective mixing from commercial-scale injection. *Environ Earth Sci*. doi:10.1007/s12665-012-1674-3
- Myrntinen A, Becker V, Nowak M, Zimmer M, Pilz P, Barth JAC (2012) Analyses of pre-injection reservoir data for stable carbon isotope trend predictions in CO₂ monitoring: preparing for CO₂ injection. *Environ Earth Sci*. doi:10.1007/s12665-012-1710-3
- Nordbotten JM, Celia MA (2012) Geological storage of CO₂—modeling approaches for large scale simulations. Wiley, New York
- Norden B, Foerster A, Behrends K, Prokoph K, Stecken L, Meyer R (2012) Geological and thermal structure of the larger Altensalzswedel area: Inputs for a shared earth model of the CLEAN site. *Environ Earth Sci*. doi:10.1007/s12665-012-1709-9
- Park YC, Huh DG, Park CH (2012) A pressure-monitoring method to warn CO₂ leakage in geological storage sites. *Environ Earth Sci*. doi:10.1007/s12665-012-1667-2
- Peter A, Lamert H, Beyer M, Hornbruch G, Heinrich B, Schulz A, Geistlinger H, Schreiber P, Dietrich P, Werban U, Vogt C,

- Richnow HH, Großmann J, Dahmke A (2012) Investigation of the geochemical impact of CO₂ on shallow groundwater: design and implementation of a CO₂ injection test in Northeast Germany. *Environ Earth Sci*. doi:[10.1007/s12665-012-1700-5](https://doi.org/10.1007/s12665-012-1700-5)
- Pudlo D, Reitenbach V, Albrecht D, Ganzer L, Gernert U, Wienand J, Kohlhepp B, Gaupp R (2012) The impact of diagenetic fluid-rock reactions on Rotliegend sandstone composition and petrophysical properties (Altmark area, central Germany). *Environ Earth Sci*. doi:[10.1007/s12665-012-1723-y](https://doi.org/10.1007/s12665-012-1723-y)
- Schilling F, Borm G, Würdemann H, Möller F, Kühn M, CO₂SINK Group (2009) Status Report on the First European on-shore CO₂ Storage Site at Ketzin (Germany). *Greenhouse Gas Control Technologies 9*. *Energy Procedia* 1(1):2029–2035. doi:[10.1016/j.egypro.2009.01.264](https://doi.org/10.1016/j.egypro.2009.01.264)
- Schütze C, Sauer U, Beyer K, Lamert H, Strauch G, Braeuer K, Flechsig C, Kaempf H, Dietrich P (2012) Natural analogues—a potential approach for developing reliable monitoring methods to understand subsurface CO₂ migration processes. *Environ Earth Sci*. doi:[10.1007/s12665-012-1701-4](https://doi.org/10.1007/s12665-012-1701-4)
- Singh AK, Görke U-J, Kolditz O (2012a) Thermal analysis of the Altmark gas field for carbon dioxide injection with enhanced gas recovery. *Environ Earth Sci*. doi:[10.1007/s12665-012-1689-9](https://doi.org/10.1007/s12665-012-1689-9)
- Singh AK, Pilz P, Zimmer M, Kalbacher T, Görke U-J, Kolditz O (2012b) Numerical simulation and geophysical monitoring of tracer transport in the Altmark gas field. *Environ Earth Sci*. doi:[10.1007/s12665-012-1688-x](https://doi.org/10.1007/s12665-012-1688-x)
- Würdemann H, Möller F, Kühn M, Heidug W, Christensen NP, Borm G, Schilling FR (2010) CO₂SINK—from site characterisation and risk assessment to monitoring and verification: one year of operational experience with the field laboratory for CO₂ storage at Ketzin, Germany. *Int J Greenhouse Gas Control* 4(6):938–951. doi:[10.1016/j.ijggc.2010.08.010](https://doi.org/10.1016/j.ijggc.2010.08.010)