ERRATUM

Erratum to: Hydrological modelling of changes in the water balance due to the impact of woody biomass production in the North German Plain

Jens Hartwich¹ · Markus Schmidt² · Jens Bölscher¹ · Christian Reinhardt-Imjela¹ · Dieter Murach² · Achim Schulte¹

Published online: 2 August 2016
© Springer-Verlag Berlin Heidelberg 2016

Erratum to: Environ Earth Sci (2016)75:1071
DOI 10.1007/s12665-016-5870-4

The original article has been published inadvertently with an error in the table head in Table 7. Table 7 with corrected table head is given below.

The online version of the original article can be found under doi:[10.1007/s12665-016-5870-4](https://doi.org/10.1007/s12665-016-5870-4).

✉ Jens Hartwich
jens.hartwich@fu-berlin.de

¹ Institute of Geographical Science, Applied Geography, Environmental Hydrology and Resource Management, Freie Universität Berlin, Berlin, Germany

² Faculty of Forest and Environment, Eberswalde University for Sustainable Development, University of Applied Science, Eberswalde, Germany

Table 7 Values suggested for willow and poplar SRC parametrization based on own measurements and literature; definition and estimation parameters based on the SWAT theoretical documentation (Neitsch et al. 2011)

Changed crop parameter	Implemented values		Former values in SWAT database Willow and poplar ^a
	Willow	Poplar	
Optimal temperature (T_OPT) (°C)	18	18	30
Base temperature (T_BASE) (°C)	5	5	10
Maximum potential LAI (BLAI) (m ² /m ²)	6.2	8.8	5
Fraction of growing season when growth declines (DLAI) (–)	0.701	0.87	0.99
Minimum LAI for plant during dormant period (ALAI_MIN or WAI) (m ² /m ²)	0.9	0.8	0.75
First point fraction of BLAI for optimum growth curve (LAIMX1) (–)	0.225	0.123	0.05
Second point fraction of BLAI for optimum growth curve (LAIMX2) (–)	0.916	0.899	0.95
Fraction of growing season coinciding LAIMX1 (FRGRW1) (–)	0.04	0.04	0.05
Fraction of growing season coinciding LAIMX2 (FRGW2) (–)	0.228	0.228	0.4
Max. canopy height (CHTMX) (m)	8	8	7.5
Max. rooting depth (RDMX) (m)	2	2	3.5
Number of years for tree to reach full development (MAT_YRS) (years)	3	3	10

^a Willow and poplar are separate in the SWAT database, but the values selected are equal for both tree genera