Erratum to: Excited States in P3HT and P3HT/PCBM Blends

K. Müller · M. Richter · S. Philip · M. Kunst ·
D. Schmeißer

Published online: 7 March 2012
© Springer Science+Business Media, LLC 2012

Erratum to: BioNanoSci. (2012) 2:42–51
DOI 10.1007/s12668-011-0031-4

In the original article the π and σ symbols in Figs. 2, 3, and 4 did not appear correctly. Following are the corrected figures with their corresponding captions:

The online version of the original article can be found at <http://dx.doi.org/10.1007/s12668-011-0031-4>.

K. Müller (✉) · M. Richter · S. Philip · D. Schmeißer
Brandenburgische Technische Universität Cottbus,
Angewandte Physik-Sensorik,
03046 Cottbus, Germany
e-mail: muellerk@tu-cottbus.de

M. Kunst
Helmholtz-Zentrum Berlin, Institut für Solare Brennstoffe (E-16),
Hahn-Meitner-Platz 1,
14109 Berlin, Germany

Fig. 2 Combined UPS-and C1s-XAS spectra of a rr-P3HT film, with respect to the Fermi level. The VBM and CBM are assigned as *arrows*; the features *E1* and *E2* are of excitonic; *P0*, *1*, *2*, and *3* are of polaronic nature, as described in text

Fig. 3 2D resPES diagram of rr-P3HT on ITO. The photon energy is plotted as the *x*-axis, and the binding energy is plotted as *y*-axis. The intensity of the signal is given as a contour. Indicated are the two CIS cuts of Fig. 4 with an initial state energy of -1.6 and -6.0 eV, corresponding to the C2p- π and C2p- π - σ bands, respectively. Also, the C1s XPS line excited by second-order light is marked, appearing in the lower initial state energy region

Fig. 4 XAS spectra in constant initial state mode (*CIS*), for different binding energies and in constant final state mode (*CFS*), for different kinetic energies. The corresponding resPES plot is given in Fig. 3. The *dashed* part of the C2p- π band is the C1s XPS line excited by second-order light appearing in the lower initial state energy region of the 2D resPES spectra