

Basic properties of multiplication and composition operators between distinct Orlicz spaces

T. Chawziuk¹ · Y. Estaremi² · H. Hudzik^{1,3} ·
S. Maghsoudi⁴ · I. Rahmani⁴

Received: 31 March 2016 / Accepted: 16 November 2016 / Published online: 15 December 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract First, we present some simple (and easily verifiable) necessary conditions and sufficient conditions for boundedness of the multiplication operator M_u and composition operator C_T acting from Orlicz space $L^{\Phi_1}(\Omega)$ into Orlicz space $L^{\Phi_2}(\Omega)$ over arbitrary complete, σ -finite measure space (Ω, Σ, μ) . Next, we investigate the problem of conditions on the generating Young functions, the function u , and/or the function $h = d(\mu \circ T^{-1})/d\mu$, under which the operators M_u and C_T are of closed range or finite rank. Finally, we give necessary and sufficient conditions for boundedness of the operators M_u and C_T in terms of techniques developed within the theory of Musielak–Orlicz spaces.

✉ H. Hudzik
hudzik@amu.edu.pl

T. Chawziuk
tchawz@gmail.com

Y. Estaremi
yestaremi@pnu.ac.ir

S. Maghsoudi
s_maghsodi@znu.ac.ir

I. Rahmani
i_rahmani@znu.ac.ir

- ¹ Faculty of Mathematics and Computer Science, Adam Mickiewicz University in Poznań, Umultowska 87, 61-614 Poznan, Poland
- ² Department of Mathematics, Payame Noor University (PNU), P. O. Box: 19395-3697, Tehran, Iran
- ³ Faculty of Economics and Information Technology, The State University of Applied Sciences in Plock, Nowe Trzepowo 55, 09-402 Plock, Poland
- ⁴ Department of Mathematics, University of Zanjan, P.O. Box: 45195-313, Zanjan, Iran

Keywords Multiplication operator · Composition operator · Continuous operators · Closed-range operators · Finite-rank operators · Orlicz space · Musielak–Orlicz space · Lebesgue space · Measurable transformation · Radon–Nikodym derivative

Mathematics Subject Classification 47B38 · 47B33 · 46E30

1 Introduction

Given the functions f and T ($f : \Omega \rightarrow \mathbb{R}$, $T : \Omega \rightarrow \Omega$), one way to produce, under certain conditions, a new function is to compose them, i.e. to evaluate f at points $T(t)$. The resulting function is denoted by $f \circ T$, and the operator $f \mapsto f \circ T$, as f runs through a linear function space, is a linear operator C_T called the composition operator induced by a mapping T . Another way to produce a new function out of given functions u and f (each from Ω into \mathbb{R}) is to multiply them, whenever it makes sense, and this gives rise to the operator called the multiplication operator M_u induced by a function u . A combination of the two methods leads to yet another operator, named the weighted composition operator, defined by $M_u C_T f = u(f \circ T)$. Composition and multiplication operators received much attention over the past several decades, especially in spaces of measurable functions such as L^p -spaces, Bergman spaces, and, to a lesser degree, Orlicz spaces. They also play an important role in the study of operators on Hilbert spaces. Consequently, by now there exists a vast literature on the properties of these transformations in various function spaces; we refer the interested reader to the beautiful books [5, 35, 36], and, for more recent studies, to [9–11, 14, 23, 25], for instance.

As said, composition and multiplication operators between classical L^p -spaces have been the subject-matter of intensive and extensive study and they feature prominently in operator theory, operator algebras, dynamical systems, and endomorphisms of Banach algebras. Composition operators were used by Banach [2] in his study of isometries between function spaces. In this paper, we aim at a generalization to Orlicz spaces of some results of the paper [38] concerning the classical Lebesgue spaces. Apparently, the paper [24] by Kumar in 1997 was the first attempt at the study of composition operators between Orlicz spaces. In 2004, Cui, Hudzik, Kumar, and Maligranda in [6] considered the composition operator between Orlicz spaces induced by a non-singular measurable transformation and studied its boundedness and compactness. Some other results on the boundedness and compactness of the composition operator between Orlicz spaces were published in [6–8, 12, 13, 16, 21, 25, 33].

Our concern in this paper is to state and prove some necessary conditions, sufficient conditions, and some simultaneously necessary and sufficient conditions for the composition and multiplication operator between distinct Orlicz spaces to be bounded or to have closed-range or finite rank. Our results generalize and improve on some recent results to be found in the literature.

2 Preliminaries and basic lemmas

In this section, for the convenience of the reader, we gather some essential facts on Orlicz spaces and prove two basic lemmas for later use. For more detail on Orlicz spaces, see [22,34].

A function $\Phi : \mathbb{R} \rightarrow [0, \infty]$ is called a *Young function* if Φ is convex, even, and $\Phi(0) = 0$; we will also assume that Φ is neither identically zero nor identically infinite on $(0, \infty)$. The fact that $\Phi(0) = 0$, along with the convexity of Φ , implies that $\lim_{x \rightarrow 0^+} \Phi(x) = 0$; while $\Phi \neq 0$, again along with the convexity of Φ , implies that $\lim_{x \rightarrow \infty} \Phi(x) = \infty$. We set $a_\Phi := \sup\{x \geq 0 : \Phi(x) = 0\}$ and $b_\Phi := \sup\{x > 0 : \Phi(x) < \infty\}$. Then it can be checked that Φ is continuous and nondecreasing on $[0, b_\Phi)$ and strictly increasing on $[a_\Phi, b_\Phi)$ if $a_\Phi < b_\Phi$. We also assume the left-continuity of the function Φ at b_Φ , i.e. $\lim_{x \rightarrow b_\Phi^-} \Phi(x) = \Phi(b_\Phi) \in [0, \infty]$.

To each Young function Φ is associated another convex function $\Psi : \mathbb{R} \rightarrow [0, \infty)$ with similar properties, defined by

$$\Psi(y) = \sup\{|xy| - \Phi(x) : x \geq 0\} \quad (y \in \mathbb{R}).$$

The function Ψ is called the *function complementary* to Φ in the sense of Young. Any pair of complementary functions (Φ, Ψ) satisfies Young’s inequality $xy \leq \Phi(x) + \Psi(y)$ ($x, y \in \mathbb{R}$).

The generalized inverse of the Young function Φ is defined by

$$\Phi^{-1}(y) = \inf\{x \geq 0 : \Phi(x) > y\} \quad (y \in [0, \infty)).$$

Notice that if $x \geq 0$, then $\Phi(\Phi^{-1}(x)) \leq x$, and if $\Phi(x) < \infty$, we also have $x \leq \Phi^{-1}(\Phi(x))$. There are equalities in both cases when Φ is a Young function vanishing only at zero and taking only finite values. Also, if (Φ, Ψ) is a pair of complementary Young functions, then

$$x < \Phi^{-1}(x)\Psi^{-1}(x) \leq 2x \tag{2.1}$$

for all $x \geq 0$ (Proposition 2.1.1(ii) in [34]).

By an *N-function* we mean a Young function vanishing only at zero, taking only finite values, and such that $\lim_{x \rightarrow \infty} \Phi(x)/x = \infty$ and $\lim_{x \rightarrow 0^+} \Phi(x)/x = 0$. Note that then $a_\Phi = 0$, $b_\Phi = \infty$, and, as we said above, Φ is continuous and strictly increasing on $[0, \infty)$. Moreover, a function complementary to an *N-function* is again an *N-function*.

A Young function Φ is said to satisfy the Δ_2 -condition at ∞ if $\Phi(2x) \leq K\Phi(x)$ ($x \geq x_0$) for some constants $K > 0$ and $x_0 > 0$ with $\Phi(x_0) < \infty$. A Young function Φ satisfies the Δ_2 -condition globally if $\Phi(2x) \leq K\Phi(x)$ ($x \geq 0$) for some $K > 0$.

A Young function Φ is said to satisfy the Δ' -condition (respectively, the ∇' -condition) at ∞ , if there exist $c > 0$ (respectively, $b > 0$) and $x_0 > 0$ with $\Phi(x_0) < \infty$ such that

$$\begin{aligned} &\Phi(xy) \leq c\Phi(x)\Phi(y) \quad (x, y \geq x_0) \\ &(\text{respectively, } \Phi(bxy) \geq \Phi(x)\Phi(y) \quad (x, y \geq x_0)). \end{aligned}$$

If $x_0 = 0$, these conditions are said to hold globally. Notice that if $\Phi \in \Delta'$, then $\Phi \in \Delta_2$ (both at ∞ and globally).

Let Φ_1, Φ_2 be Young functions. Then Φ_1 is called stronger than Φ_2 at ∞ , which is denoted by $\Phi_1 \overset{\ell}{>} \Phi_2$ [or $\Phi_2 \overset{\ell}{<} \Phi_1$], if

$$\Phi_2(x) \leq \Phi_1(ax) \quad (x \geq x_0)$$

for some $a \geq 0$ and $x_0 > 0$ with $\Phi_2(x_0) < \infty$; if $x_0 = 0$, this condition is said to hold globally and is then denoted by $\Phi_1 \overset{a}{>} \Phi_2$ [or $\Phi_2 \overset{a}{<} \Phi_1$]. We record the following observation for later use.

Lemma 2.1 *If Φ_1, Φ_2, Φ_3 are Young functions vanishing only at zero, taking only finite values, and such that*

$$\Phi_1(xy) \leq \Phi_2(x) + \Phi_3(y) \quad (x, y \geq 0),$$

then $\Phi_2 \overset{\ell}{\not>} \Phi_1$, and hence also $\Phi_2 \overset{a}{\not>} \Phi_1$.

Proof The inequality in the assumptions of the lemma implies that $\Phi_1(\Phi_2^{-1}(x)\Phi_3^{-1}(x)) \leq 2x$ for all $x \geq 0$, and this, together with subadditivity of Φ_1^{-1} , leads to

$$\Phi_2^{-1}(x)\Phi_3^{-1}(x) \leq \Phi_1^{-1}(2x) \leq 2\Phi_1^{-1}(x) \quad (x \geq 0). \tag{2.2}$$

Now suppose, towards a contradiction, that there are $a > 0$ and $x_0 > 0$ such that

$$\Phi_2(x) < \Phi_1(ax) \quad (x \geq x_0).$$

Then, since Φ_1^{-1} is increasing on $[0, \infty)$, we have $\Phi_1^{-1}(\Phi_2(x)) < \Phi_1^{-1}(\Phi_1(ax)) = ax$, and so, replacing x with $\Phi_2(x)$ in (2.2), we get

$$x \Phi_3^{-1}(\Phi_2(x)) = \Phi_2^{-1}(\Phi_2(x))\Phi_3^{-1}(\Phi_2(x)) \leq 2\Phi_1^{-1}(\Phi_2(x)) < 2ax \quad (x \geq x_0).$$

This means, since Φ_3 is increasing on $[0, \infty)$, that $\Phi_1(x) < \Phi_3(2a) = \text{const.}$ for all $x \geq x_0$, a contradiction, because $\Phi_2(x) \rightarrow \infty$ as $x \rightarrow \infty$. □

Let (Ω, Σ, μ) be a complete σ -finite measure space and let $L^0(\Omega)$ be the linear space of equivalence classes of Σ -measurable real-valued functions on Ω , that is, we identify functions equal μ -almost everywhere on Ω . The support S of a measurable function f is defined by $S(f) := \{t \in \Omega : f(t) \neq 0\}$. For a Young function Φ , the space

$$L^\Phi(\Omega) = \left\{ f \in L^0(\Omega) : \exists k > 0, I_\Phi(kf) < \infty \right\}$$

is a Banach space if it is equipped with the norm

$$\|f\|_\Phi = \inf \{k > 0 : I_\Phi(f/k) \leq 1\},$$

where $I_\Phi(f) = \int_\Omega \Phi(f(t))d\mu(t)$. The couple $(L^\Phi(\Omega), \|\cdot\|_\Phi)$ is called the Orlicz space generated by a Young function Φ . For any $f \in L^0(\Omega)$, we have $I_\Phi(f) \leq 1$ if

and only if $\|f\|_\Phi \leq 1$. In particular, if $I_\Phi(f) \leq a$ ($a > 0$), then $\|f\|_\Phi \leq \max(a, 1)$. If $\Phi \in \Delta_2$ and $f \in L^\Phi(\Omega)$, then $I_\Phi(kf) < \infty$ for all $k > 0$.

Let $\Phi(x) = |x|^p/p$ with $1 < p < \infty$; Φ is then a Young function and $\Psi(x) = |x|^{p'}/p'$, with $1/p + 1/p' = 1$, is the Young function complementary to Φ . Thus, with this function Φ we retrieve the classical Lebesgue space $L^p(\Omega)$, i.e. $L^\Phi(\Omega) = L^p(\Omega)$.

Recall that an atom of the measure space (Ω, Σ, μ) is a set $\mathfrak{A} \in \Sigma$ with $\mu(\mathfrak{A}) > 0$ such that if $F \in \Sigma$ and $F \subset \mathfrak{A}$, then either $\mu(F) = 0$ or $\mu(F) = \mu(\mathfrak{A})$. A measure space (Ω, Σ, μ) with no atoms is called a non-atomic measure space [29].

Remark 2.2 By saying that $\Phi \in \Delta_2$ we will always mean that the Young function Φ satisfies the Δ_2 -condition globally, since we place no restrictions on the underlying measure space (Ω, Σ, μ) , which can be finite or infinite, non-atomic, purely atomic, or mixed.

The following lemma is a key tool in some of our investigations.

Lemma 2.3 *Let Φ_1, Φ_2 be Young functions such that $\Phi_2 \not\prec_\ell \Phi_1$. If E is a non-atomic Σ -measurable set with positive measure, then there exists $f \in L^{\Phi_1}(\Omega)$ such that $f|_E \notin L^{\Phi_2}(E)$.*

Proof It is well-known that if $L^{\Phi_1}(\Omega, \Sigma, \mu) \subset L^{\Phi_2}(\Omega, \Sigma, \mu)$ for a non-atomic measure space (Ω, Σ, μ) , then $\Phi_2 \prec_\ell \Phi_1$ (see, for instance, [26] Theorem 3.4, and [28,30,31] for some extensions). Therefore, if $\Phi_2 \not\prec_\ell \Phi_1$ and E is a Σ -measurable subset of Ω with $\mu(E) > 0$, then we can find a function $g \in L^{\Phi_1}(E)$ such that $g \notin L^{\Phi_2}(E)$. Defining $f(t) = g(t)$ for $t \in E$, and $f(t) = 0$ for $t \notin E$, we have the desired function. □

Remark 2.4 We will use Lemmas 2.1 and 2.3 in conjunction since the thesis of the former is the assumption of the latter.

Given a measurable function $u \in L^0_+(\Omega)$ the mapping $f \mapsto uf$ is a linear transformation on $L^0(\Omega)$ called the *multiplication operator* induced by u and denoted M_u .

Let $T : \Omega \rightarrow \Omega$ be a measurable transformation. The transformation T is said to be non-singular if $\mu(T^{-1}(A)) = 0$ for all $A \in \Sigma \cap T(\Omega)$ with $\mu(A) = 0$. The condition ensures that the measure $\mu \circ T^{-1}$ defined by $\mu \circ T^{-1}(A) = \mu(T^{-1}(A))$ for all $A \in \Sigma \cap T(\Omega)$ is absolutely continuous with respect to the measure μ (this fact is usually denoted by $\mu \circ T^{-1} \ll \mu$). Then the Radon–Nikodym theorem guarantees the existence of a non-negative locally integrable function $h : T(\Omega) \rightarrow \mathbb{R}_+$, called the Radon–Nikodym derivative of $\mu \circ T^{-1}$ with respect to μ , such that

$$\mu \circ T^{-1}(A) = \int_A h d\mu \quad (A \in \Sigma \cap T(\Omega)).$$

We do not necessarily assume that the transformation T is surjective or injective.

The non-singular measurable transformation T induces a linear operator C_T on $L^0(\Omega)$, called the *composition operator*, defined by

$$C_T f(t) = f(T(t)) \quad (t \in \Omega, f \in L^0(\Omega)).$$

Finally, we recall two well-known facts from measure theory; for proofs see, for example, Section 11.9 in [29]:

- (i) If (Ω, Σ, μ) be a σ -finite measure space, then for every measurable real-valued function f on Ω and every atom \mathfrak{A} , there is a real number, denoted by $f(\mathfrak{A})$, such that $f = f(\mathfrak{A}) \mu$ -a.e. on \mathfrak{A} .
- (ii) If (Ω, Σ, μ) is a σ -finite measure space that fails to be non-atomic, there is a non-empty countable set of pairwise disjoint atoms $\{\mathfrak{A}_n\}_{n \in \mathbb{N}}$ with the property that $\mathfrak{B} := \Omega \setminus \bigcup_{n \in \mathbb{N}} \mathfrak{A}_n$ contains no atoms. We assume that both the atoms and their counterimages under T have strictly positive measure. The latter fact means that the Radon–Nikodym derivative h is strictly positive on all atoms from $T(\Omega)$.

We keep the above notation throughout the paper.

3 Boundedness of the multiplication operator

In this section we state various necessary conditions and sufficient conditions under which the multiplication operator M_u between distinct Orlicz spaces is bounded.

Recall the well-known theorem due to O’Neil [32] on the multiplication operator between Orlicz spaces (see also [1] and Theorem 5.4.1 in [33]): suppose that Φ_1, Φ_2, Φ_3 are Young functions such that either (a) there exist $c > 0$ and $x_0 > 0$ with $\Phi_2(x_0) < \infty$ such that $\Phi_2(cxy) \leq \Phi_1(x) + \Phi_3(y)$ for all $x, y \geq x_0$ in the case when $\mu(\Omega) < \infty$ or (b) there exists $c > 0$ such that $\Phi_2(cxy) \leq \Phi_1(x) + \Phi_3(y)$ for all $x, y > 0$ in the case when $\mu(\Omega) = \infty$. Let Ω contain a non-atomic set of positive measure. Then, whenever $u \in L^{\Phi_3}(\Omega)$, the multiplication operator M_u is bounded from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$.

Before going any further, let us consider some related examples.

Example 3.1 (i) Let Φ and Ψ be complementary Young functions. Since $\frac{|x|^p}{p} \leq \frac{1}{p}(\Phi(x) + \Psi(x^{p-1}))$ and $\frac{|x|^p}{p} \leq \frac{1}{p}(\Phi(x^{p-1}) + \Psi(x))$, for $x \geq 0$ and $p > 2$, by O’Neil’s theorem we get that if $u^{\frac{1}{p-1}} \in L^\Psi(\Omega)$ then the operator M_u is bounded from $L^\Phi(\Omega)$ into $L^p(\Omega)$, and if $v^{\frac{1}{p-1}} \in L^\Phi(\Omega)$ then the operator M_v is bounded from $L^\Psi(\Omega)$ into $L^p(\Omega)$.

(ii) Let $\Omega = [a, b]$, $a, b > 1$, $p > 1$, and let μ be Lebesgue measure. If we take $\Phi_1(x) = e^{|x|^p} - |x|^p - 1$, $\Phi_2(x) = \frac{|x|^p}{p}$, and $\Phi_3(x) = (1 + |x|^p) \log(1 + |x|^p) - |x|^p$, then, since $\phi_1(x) = \Phi_1(|x|^{1/p})$ and $\phi_3(x) = \Phi_3(|x|^{1/p})$ are complementary Young functions, Young’s inequality yields $\Phi_2(xy) \leq \Phi_1(x) + \Phi_3(y)$ for all $x, y \geq 0$. Now, if $u(t) = \sqrt[p]{t^p - 1}$, it is clear that $I_{\Phi_3}(u) < \infty$, i.e. $u \in L^{\Phi_3}(\Omega)$. O’Neil’s theorem implies that M_u is a bounded operator from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$.

(iii) Let $A = (0, a]$, $B = \{\ln t : t \in \mathbb{N}, t > a\}$, for $a > 0$, and let $\Omega = A \cup B$. Take $\Phi_1(x) = e^{|x|} - |x| - 1$, $\Phi_2(x) = (1 + |x|) \ln(1 + |x|) - |x|$, and for every Lebesgue measurable $C \subset \Omega$ define $\mu(C) = \mu_1(C \cap A) + \mu_2(C \cap B)$, where μ_1 is Lebesgue measure and $\mu_2(\{\ln t\}) = 1/t^3$ for $\ln t \in B$. If we set $u(t) = 1/t^2$, then M_u is not bounded from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$. Indeed, for $f(t) = t$ we

have

$$\begin{aligned} I_{\Phi_1}(f) &= \int_{\Omega} (\Phi_1 \circ f) d\mu = \int_{\Omega} (e^t - t - 1) d\mu \\ &= \int_A (e^t - t - 1) d\mu + \int_B (e^t - t - 1) d\mu < \infty, \end{aligned}$$

where we used the following calculation

$$\int_B (e^t - t - 1) d\mu = \sum_{n>a} (e^{\ln n} - \ln n - 1) \frac{1}{n^3} < \sum_{n>a} \frac{1}{n^2} < \infty.$$

Hence $f \in L^{\Phi_1}(\Omega)$. But for any $\lambda > 0$ we have

$$\begin{aligned} I_{\Phi_2}(\lambda M_u f) &= \int_{\Omega} (\Phi_2 \circ \lambda M_u f) d\mu = \int_{\Omega} \Phi_2 \left(\frac{\lambda}{t} \right) d\mu \\ &= \int_{\Omega} \left(\left(1 + \frac{\lambda}{t} \right) \ln \left(1 + \frac{\lambda}{t} \right) - \frac{\lambda}{t} \right) d\mu \\ &= \int_A \left(\left(1 + \frac{\lambda}{t} \right) \ln \left(1 + \frac{\lambda}{t} \right) - \frac{\lambda}{t} \right) d\mu \\ &\quad + \int_B \left(\left(1 + \frac{\lambda}{t} \right) \ln \left(1 + \frac{\lambda}{t} \right) - \frac{\lambda}{t} \right) d\mu \\ &> \int_0^a \left(\left(1 + \frac{\lambda}{t} \right) \ln \left(1 + \frac{\lambda}{t} \right) - \frac{\lambda}{t} \right) dt \\ &= \lambda \int_{\lambda/a}^{\infty} ((1 + s) \ln(1 + s) - s) \frac{1}{s^2} ds \\ &= \lambda \int_{\lambda/a}^{\infty} \left(\frac{\ln(1 + s)}{s^2} + \frac{\ln(1 + s) - 1}{s} \right) ds \\ &> \lambda \int_{\lambda/a}^{\infty} \left(\frac{\ln(1 + s) - 1}{s} \right) ds = \infty, \end{aligned}$$

and so $M_u f \notin L^{\Phi_2}(\Omega)$. The operator M_u does even act from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$ and, by O’Neil’s theorem, we conclude that if a Young function Φ_3 satisfies $\Phi_2(xy) \leq \Phi_1(x) + \Phi_3(y)$ for all $x, y \geq 0$, then u does not belong to $L^{\Phi_3}(\Omega)$.

Maligranda and Persson in [27] proved a theorem which can be stated in the following way: assume that Φ_1, Φ_2, Φ_3 are Young functions with values in $[0, \infty)$ such that $\Phi_2(xy) \leq \Phi_1(x) + \Phi_3(y)$ for all $x, y \geq 0$ and $\Phi_2^{-1}(x) \leq \Phi_1^{-1}(x)\Phi_3^{-1}(x)$ for all $x \geq 0$, and assume further that either (A) Ω is a non-atomic measure space or (B) $\Phi_2(x) = x^r$ and $x^{-r}\Phi_1(x)$ is nondecreasing. Then M_u is a bounded multiplication operator from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$ if and only if $u \in L^{\Phi_3}(\Omega)$.

These results have been generalized and extended to the more general setting of Calderón–Lozanovskii spaces in two recent papers [19,20] by Kolwicz et al.

We give another necessary condition on the function u so that it induces a bounded multiplication operator M_u between distinct Orlicz spaces; it will be used in the proof of Theorem 4.11.

Proposition 3.2 *Let Φ_1 and Φ_2 be Young functions vanishing only at zero, taking only finite values, and such that Φ_1 is an N -function, $\Phi_1 \in \Delta'$, $\Phi_2 \in \Delta_2$, and $\Phi_3 := \Psi_2 \circ \Psi_1^{-1}$ is a Young function. If $u \in L^0_+(\Omega)$ and M_u is a bounded multiplication operator from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$, then $u \in L^{\Psi_3 \circ \Psi_1}(\Omega)$.*

Proof Let $u \in L^0_+(\Omega)$ and let M_u be a bounded multiplication operator from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$ induced by the function u . Since $\Phi_1 \in \Delta_2$ (as a consequence of the fact that $\Phi_1 \in \Delta'$) and $\Phi_2 \in \Delta_2$, the dual spaces of $L^{\Phi_1}(\Omega)$ and $L^{\Phi_2}(\Omega)$ are equal to $L^{\Psi_1}(\Omega)$ and $L^{\Psi_2}(\Omega)$, respectively. Hence the adjoint operator M_u^* to the bounded operator M_u is bounded and acts from $L^{\Psi_2}(\Omega)$ into $L^{\Psi_1}(\Omega)$.

It is easy to check that $\Phi_1 \in \Delta'$ implies that $\Psi_1 \in \nabla'$, i.e. there is $b > 0$ such that $\Psi_1(bxy) \geq \Psi_1(x)\Psi_1(y)$ for all $x, y \geq 0$.

Let f be any function from $L^{\Phi_3}(\Omega)$. By the definition of Φ_3 , we have that the function $\Psi_1^{-1} \circ |f|$ is contained in the space $L^{\Psi_2}(\Omega)$, and so $M_u^*(\Psi_1^{-1} \circ |f|)$ is contained in the space $L^{\Psi_1}(\Omega)$. Therefore, there is $\lambda > 0$ such that $I_{\Psi_1}(\lambda M_u^*(\Psi_1^{-1} \circ |f|)) < \infty$. We have

$$\begin{aligned} \left| \int_{\Omega} (\Psi_1 \circ \frac{\lambda}{b} u) f d\mu \right| &\leq \int_{\Omega} (\Psi_1 \circ \frac{\lambda}{b} u) |f| d\mu \leq \int_{\Omega} (\Psi_1 \circ \frac{\lambda}{b} u) (\Psi_1 \circ \Psi_1^{-1} \circ |f|) d\mu \\ &\leq \int_{\Omega} \Psi_1(b \frac{\lambda}{b} u \Psi_1^{-1} \circ |f|) d\mu = \int_{\Omega} \Psi_1(\lambda M_u^*(\Psi_1^{-1} \circ |f|)) d\mu < \infty. \end{aligned}$$

This means that $\Psi_1 \circ \frac{\lambda}{b} u$ belongs to the Köthe dual of $L^{\Phi_3}(\Omega)$, which is equal to $L^{\Psi_3}(\Omega)$. Consequently, there is $\alpha \in (0, 1)$ such that $I_{\Psi_3}(\alpha \Psi_1 \circ \frac{\lambda}{b} u) < \infty$, whence $I_{\Psi_3 \circ \Psi_1}(\alpha \frac{\lambda}{b} u) = I_{\Psi_3}(\Psi_1 \circ \alpha \frac{\lambda}{b} u) \leq I_{\Psi_3}(\alpha \Psi_1 \circ \frac{\lambda}{b} u) < \infty$. Hence $u \in L^{\Psi_3 \circ \Psi_1}(\Omega)$. \square

In the case of a non-atomic complete and σ -finite measure space, some necessary and sufficient conditions for the boundedness of M_u and C_T from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$ were established in [3]. It turns out that these are equivalent to the conditions for the inclusion of the Orlicz space $L^{\Phi_1}(\Omega)$ into the Musielak–Orlicz space $L^{\Phi_2, u}(\Omega)$ and of the Orlicz space $L^{\Phi_1}(T(\Omega))$ into the Musielak–Orlicz space $L^{\Phi_2}_h(T(\Omega))$, where $h = d(\mu \circ T^{-1})/d\mu$, respectively. The spaces $L^{\Phi_2, u}(\Omega)$ and $L^{\Phi_2}_h(T(\Omega))$ are generated over the measure spaces (Ω, Σ, μ) and $(T(\Omega), \Sigma \cap T(\Omega), \mu|_{\Sigma \cap T(\Omega)})$ by the Musielak–Orlicz functions $\Phi_2(xu(t))$ ($t \in \Omega$ and $x \in \mathbb{R}$) and $\Phi_2(x)h(t)$ ($t \in T(\Omega)$ and $x \in \mathbb{R}$), respectively, $t \in \Omega$ and $x \in \mathbb{R}$.

We shall now prove a theorem from which it follows that, if the measure space (Ω, Σ, μ) is non-atomic, for the above inclusions to hold it is necessary that $\Phi_2 \stackrel{\ell}{<} \Phi_1$. This might come as a surprise since it means that the simple condition $\Phi_2 \stackrel{\ell}{<} \Phi_1$, which does not involve the Radon–Nikodym derivative h , is necessary for the inclusion $L^{\Phi_1}(T(\Omega)) \subset L^{\Phi_2}_h(T(\Omega))$, which does.

Theorem 3.3 *Let Φ_1, Φ_2 be Young functions such that $\Phi_2 \not\stackrel{\ell}{\prec} \Phi_1$ and let (Ω, Σ, μ) be a non-atomic measure space. Then there are no non-zero bounded operators $C_T M_u$ from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$.*

Proof Suppose, to the contrary, that $C_T M_u$ is a non-zero bounded linear operator from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$ and let

$$E_n = \left\{ t \in \Omega : u(t) > \frac{1}{n} \right\} \cap \left\{ t \in T(\Omega) : h(t) > \frac{1}{n} \right\}$$

For any $n \in \mathbb{N}$. Then $\{E_n\}_{n \in \mathbb{N}}$ is an increasing sequence of Σ -measurable sets. Since $C_T M_u$ is non-zero, $\mu(E_m) > 0$ for some $m \in \mathbb{N}$, whence also $\mu(E_n) > 0$ for all $n \geq m$. We assume without loss of generality that $\mu(E_n) > 0$ for all $n \in \mathbb{N}$. Let

$F \subset E := \bigcup_n E_n$ and $0 < \mu(F) < \infty$. The assumption that $\Phi_2 \not\stackrel{\ell}{\prec} \Phi_1$ implies that an increasing sequence of positive numbers $\{y_n\}$ can be found such that $\Phi_2(y_n) > \Phi_1(2^n n^3 y_n)$. Since the measure space (Ω, Σ, μ) is non-atomic, we can choose a sequence $\{F_n\}$ of pairwise disjoint measurable subsets of F such that $F_n \subset E_n$ and $\mu(F_n) = \frac{\Phi_1(y_1)\mu(F)}{2^n \Phi_1(n^3 y_n)}$ For any $n \in \mathbb{N}$. This is possible because ν is non-atomic and $\Phi_1(n^3 y_n) \geq \Phi_1(y_n) \geq \Phi_1(y_1)$, and so $\mu(F_n) \leq \frac{\mu(F)}{2^n}$, and $\sum_{n=1}^{\infty} \frac{\mu(F)}{2^n} = \mu(F)$.

Define the function $f := \sum_{n=1}^{\infty} b_n \chi_{F_n}$, where $b_n := n^2 y_n$, and take arbitrary $\alpha > 0$. Then for a natural number $n_0 > \alpha$ we have

$$\begin{aligned} I_{\Phi_1}(\alpha f) &= \int_{\Omega} \Phi_1(\alpha f) d\mu = \sum_{n=1}^{\infty} \int_{\Omega} \Phi_1(\alpha b_n) \chi_{F_n} d\mu \\ &= \sum_{n=1}^{n_0} \Phi_1(\alpha b_n) \mu(F_n) + \sum_{n>n_0} \Phi_1(\alpha b_n) \mu(F_n) \\ &= \sum_{n=1}^{n_0} \Phi_1(\alpha b_n) \mu(F_n) + \mu(F) \sum_{n>n_0} \frac{\Phi_1(\alpha b_n) \Phi_1(y_1)}{2^n \Phi_1(n^3 y_n)} \\ &\leq \sum_{n=1}^{n_0} \Phi_1(\alpha b_n) \mu(F_n) + \mu(F) \sum_{n>n_0} \frac{\Phi_1(n^3 y_n) \Phi_1(y_1)}{2^n \Phi_1(n^3 y_n)} < \infty. \end{aligned}$$

This implies that $f \in L^{\Phi_1}(\Omega)$. But for $m_0 > 0$ such that $\frac{1}{m_0} < \alpha$, we obtain

$$\begin{aligned} I_{\Phi_2}(\alpha C_T M_u f) &= \int_{\Omega} \Phi_2(\alpha C_T M_u f) d\mu = \int_{T(\Omega)} h(t) \Phi_2(\alpha u f) d\mu \\ &\geq \sum_{n \geq m_0} \int_{F_n} h(t) \Phi_2(\alpha b_n u) d\mu \geq \sum_{n \geq m_0} \int_{F_n} \frac{1}{n} \Phi_2(b_n/n^2) d\mu \\ &\geq \sum_{n \geq m_0} \int_{F_n} \frac{1}{n} \Phi_2(y_n) d\mu \geq \sum_{n \geq m_0} \frac{1}{n} \Phi_1(2^n n^3 y_n) \mu(F_n) \\ &\geq \sum_{n \geq m_0} \frac{1}{n} 2^n \Phi_1(n^3 y_n) \mu(F_n) \geq \mu(F) \sum_{n \geq m_0} \frac{1}{n} \Phi_1(y_1) = \infty, \end{aligned}$$

which, by the arbitrariness of $\alpha > 0$, contradicts the boundedness of $C_T M_u$. (In fact, we even proved that $C_T M_u$ does not act from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$). \square

We provide some necessary conditions for the boundedness of the multiplication operator M_u from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$ under the assumption that $\Phi_1(xy) \leq \Phi_2(x) + \Phi_3(y)$ for some Young function Φ_3 and for all $x, y \geq 0$.

Theorem 3.4 *Let Φ_1, Φ_2, Φ_3 be Young functions vanishing only at zero, taking only finite values, and such that $\Phi_1(xy) \leq \Phi_2(x) + \Phi_3(y)$ for all $x, y \geq 0$. If $u \in L^0_+(\Omega)$ induces a bounded multiplication operator $M_u : L^{\Phi_1}(\Omega) \rightarrow L^{\Phi_2}(\Omega)$, then*

(i) $u(t) = 0$ for μ -a.e. $t \in \mathfrak{B}$, the non-atomic part of Ω ;

(ii) $\sup_{n \in \mathbb{N}} u(\mathfrak{A}_n) \Phi_3^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right) < \infty$.

Proof Let the assumptions about $\Phi_i(i=1,2,3)$ be satisfied and M_u be bounded. First we prove (i). If $\mu\{t \in \mathfrak{B} : u(t) \neq 0\} > 0$, then there exists a constant $\delta > 0$ such that the set $E = \{t \in \mathfrak{B} : u(t) > \delta\}$ has positive measure. Since E is non-atomic, $\mu(E) > 0$, and $\Phi_1(xy) \leq \Phi_2(x) + \Phi_3(y)$ for all $x, y \geq 0$, by Lemmas 2.1 and 2.3 we have that there exists $f \in L^{\Phi_1}(\Omega)$ such that $f|_E \notin L^{\Phi_2}(E)$, and so

$$\infty = \int_E \Phi_2\left(\frac{\delta f(t)}{\|M_u f\|_{\Phi_2}}\right) d\mu \leq \int_{\Omega} \Phi_2\left(\frac{u(t)f(t)}{\|M_u f\|_{\Phi_2}}\right) d\mu \leq 1,$$

which is a contradiction. Thus (i) holds.

Now we prove (ii). We may assume that the function u is not identically zero. For each $n \in \mathbb{N}$, put $f_n = \Phi_1^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right)\chi_{\mathfrak{A}_n}$. It is clear that $f_n \in L^{\Phi_1}(\Omega)$ and $I_{\Phi_1}(f_n) = 1$, whence $\|f_n\|_{\Phi_1} = 1$. Since the operator M_u is bounded, we have

$$\begin{aligned} 1 &\geq \int_{\Omega} \Phi_2\left(\frac{u(t)f_n(t)}{\|M_u f_n\|_{\Phi_2}}\right) d\mu = \int_{\mathfrak{A}_n} \Phi_2\left(\frac{u(t)\Phi_1^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right)}{\|M_u f_n\|_{\Phi_2}}\right) d\mu \\ &= \Phi_2\left(\frac{u(\mathfrak{A}_n)\Phi_1^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right)}{\|M_u f_n\|_{\Phi_2}}\right) \mu(\mathfrak{A}_n). \end{aligned}$$

Therefore

$$\frac{u(\mathfrak{A}_n)\Phi_1^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right)}{\Phi_2^{-1}(1/\mu(\mathfrak{A}_n))} \leq \|M_u f_n\|_{\Phi_2}. \tag{3.1}$$

Put $x = 1/\mu(\mathfrak{A}_n)$ in the inequality $\Phi_2^{-1}(x)\Phi_3^{-1}(x) \leq 2\Phi_1^{-1}(x)$ derived in lines 2–3 of the proof of Lemma 2.1 and use inequality (3.1) to obtain

$$u(\mathfrak{A}_n) \Phi_3^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right) \leq u(\mathfrak{A}_n) \frac{2\Phi_1^{-1}(1/\mu(\mathfrak{A}_n))}{\Phi_2^{-1}(1/\mu(\mathfrak{A}_n))} \leq 2\|M_u f_n\|_{\Phi_2} \leq 2\|M_u\| < \infty.$$

This completes the proof. \square

We have a straightforward consequence.

Corollary 3.5 *Under the assumptions of Theorem 3.4, if (Ω, Σ, μ) is a non-atomic measure space, then the multiplication operator M_u is bounded from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$ if and only if $M_u = 0$.*

Now we present some sufficient conditions for the continuity of the operator M_u from one Orlicz space into another.

Theorem 3.6 *Let Φ_1, Φ_2 be Young functions vanishing only at zero, taking only finite values, and such that $\Phi_1, \Phi_2 \in \Delta'$ and $\Phi_2 \circ \Phi_1^{-1}$ is a Young function. Then $u \in L^{\Phi_1}_+(\Omega)$ induces a bounded multiplication operator $M_u : L^{\Phi_1}(\Omega) \rightarrow L^{\Phi_2}(\Omega)$ if*

- (i) $u(t) = 0$ for μ -a.e. $t \in \mathfrak{B}$;
- (ii) $\sup_{n \in \mathbb{N}} \Phi_2\left[\frac{u(\mathfrak{A}_n)}{\Phi_1^{-1}(\mu(\mathfrak{A}_n))}\right] \mu(\mathfrak{A}_n) < \infty$.

Proof Suppose that (i) and (ii) hold and set $M := \sup_{n \in \mathbb{N}} \Phi_2\left[\frac{u(\mathfrak{A}_n)}{\Phi_1^{-1}(\mu(\mathfrak{A}_n))}\right] \mu(\mathfrak{A}_n)$. Then for each $f \in L^{\Phi_1}(\Omega)$ we have

$$\begin{aligned} I_{\Phi_2}(M_u f) &= \int_{\Omega} \Phi_2(M_u f) d\mu = \int_{\mathfrak{B}} \Phi_2(u(t) f(t)) d\mu + \int_{\bigcup_n \mathfrak{A}_n} \Phi_2(u(t) f(t)) d\mu \\ &= \sum_{n \in \mathbb{N}} \int_{\mathfrak{A}_n} \Phi_2(u(t) f(t)) d\mu = \sum_{n \in \mathbb{N}} \Phi_2(u(\mathfrak{A}_n) f(\mathfrak{A}_n)) \mu(\mathfrak{A}_n) \\ &= \sum_{n \in \mathbb{N}} \Phi_2 \left[\Phi_1^{-1}(\mu(\mathfrak{A}_n)) \Phi_1^{-1} \circ \Phi_1(f(\mathfrak{A}_n)) \frac{u(\mathfrak{A}_n)}{\Phi_1^{-1}(\mu(\mathfrak{A}_n))} \right] \mu(\mathfrak{A}_n) \\ &\leq \sum_{n \in \mathbb{N}} \Phi_2 \left[\Phi_1^{-1} \left[c_1 \mu(\mathfrak{A}_n) \Phi_1(f(\mathfrak{A}_n)) \right] \frac{u(\mathfrak{A}_n)}{\Phi_1^{-1}(\mu(\mathfrak{A}_n))} \right] \mu(\mathfrak{A}_n) \\ &\leq c_2 \sum_{n \in \mathbb{N}} \Phi_2 \circ \Phi_1^{-1} \left[c_1 \mu(\mathfrak{A}_n) \Phi_1(f(\mathfrak{A}_n)) \right] \Phi_2 \left[\frac{u(\mathfrak{A}_n)}{\Phi_1^{-1}(\mu(\mathfrak{A}_n))} \right] \mu(\mathfrak{A}_n) \\ &\leq c_2 \sum_{n \in \mathbb{N}} \Phi_2 \circ \Phi_1^{-1} \left[c_1 \mu(\mathfrak{A}_n) \Phi_1(f(\mathfrak{A}_n)) \right] M \\ &\leq c_2 M \Phi_2 \circ \Phi_1^{-1} \left[c_1 \sum_{n \in \mathbb{N}} \mu(\mathfrak{A}_n) \Phi_1(f(\mathfrak{A}_n)) \right], \end{aligned}$$

where c_1, c_2 are positive constants related to conditions $\Phi_1 \in \Delta'$ and $\Phi_2 \in \Delta'$, respectively, and the last inequality follows from the superadditivity of the convex function $\Phi_2 \circ \Phi_1^{-1}$ on the interval $[0, \infty)$. Now let $\|f\|_{\Phi_1} \leq 1$. We get

$$I_{\Phi_2}(M_u f) \leq c_2 M \Phi_2 \circ \Phi_1^{-1}(c_1) < \infty.$$

This implies that $\|M_u f\|_{\Phi_2} \leq \max(c_2 M \Phi_2 \circ \Phi_1^{-1}(c_1), 1)$, and so M_u is bounded. \square

Remark 3.7 The following results on the classical Lebesgue spaces, presented in [38] (see also [18]), are an immediate consequence of Theorem 3.6.

1. Taking $\Phi_1(x) = |x|^p/p$ and $\Phi_2(x) = |x|^q/q$, where $1 < p < q < \infty$, by Theorems 3.4 and 3.6, we obtain that the multiplication operator M_u induced by a function $u \in L^0_+(\Omega)$ is bounded from $L^p(\Omega)$ into $L^q(\Omega)$ if and only if the following conditions hold:
 - (i) $u(t) = 0$ for μ -almost all $t \in \mathfrak{B}$;
 - (ii) $\sup_{n \in \mathbb{N}} \frac{u(\mathfrak{A}_n)^r}{\mu(\mathfrak{A}_n)^r} < \infty$, where $q^{-1} + r^{-1} = p^{-1}$.
2. Similarly, taking $\Phi_1(x) = |x|^p/p$ and $\Phi_2(x) = |x|^q/q$, where $1 < q < p < \infty$, by O’Niel’s theorem (see the beginning of Sect. 3) combined with Proposition 3.2 we obtain that the multiplication operator M_u induced by a function $u \in L^0_+(\Omega)$ is bounded from $L^p(\Omega)$ into $L^q(\Omega)$ if and only if $u \in L^r(\Omega)$, where $p^{-1} + r^{-1} = q^{-1}$.

4 Boundedness of the composition operator

In this section, we give necessary conditions and sufficient conditions under which the composition operator C_T acts continuously between distinct Orlicz spaces.

Theorem 4.1 *Let $T : \Omega \rightarrow \Omega$ be a non-singular measurable transformation and let Φ_1, Φ_2, Φ_3 be Young functions vanishing only at zero, taking only finite values, and such that $\Phi_1(xy) \leq \Phi_2(x) + \Phi_3(y)$ for all $x, y \geq 0$. If T induces a bounded composition operator $C_T : L^{\Phi_1}(\Omega) \rightarrow L^{\Phi_2}(\Omega)$, then $h(t) = 0$ for μ -almost all $t \in T(B)$.*

Proof Suppose, towards a contradiction, that $\mu(\{t \in T(\mathfrak{B}) : h(t) \neq 0\}) > 0$. Then there is a constant $\delta > 0$ such that the set $E = \{t \in T(\mathfrak{B}) : h(t) > \delta\}$ has positive measure. Since $\mu(E) > 0$, E is non-atomic, and $\Phi_1(xy) \leq \Phi_2(x) + \Phi_3(y)$ for all $x, y \geq 0$, by Lemmas 2.1 and 2.3 we have that there exists $f \in L^{\Phi_1}(\Omega)$ such that $f|_E \notin L^{\Phi_2}(E)$. Since $E \subset T(\mathfrak{B})$, we obtain that $C_T f \neq 0$ (otherwise we would have $f|_E \in L^{\Phi_2}(E)$). Together with the fact that C_T is bounded, the above gives

$$\begin{aligned} \infty &= \int_E \delta \Phi_2 \left(\frac{f(t)}{\|C_T f\|_{\Phi_2}} \right) d\mu \leq \int_E h(t) \Phi_2 \left(\frac{f(t)}{\|C_T f\|_{\Phi_2}} \right) d\mu \\ &\leq \int_{\Omega} \Phi_2 \left(\frac{C_T f(t)}{\|C_T f\|_{\Phi_2}} \right) d\mu \leq 1, \end{aligned}$$

which is a contradiction finishing the proof. □

Remark 4.2 Although the function Φ_3 does not feature in the thesis of Theorem 4.1, it was actually used in the proof via Lemma 2.1.

Theorem 4.3 *Let $T : \Omega \rightarrow \Omega$ be a non-singular measurable transformation, and let Φ_1, Φ_2 be Young functions vanishing only at zero, taking only finite values, and such that $\Phi_1, \Phi_2 \in \Delta'$ and $\Phi_2 \circ \Phi_1^{-1}$ is a Young function. Then T induces a bounded composition operator $C_T : L^{\Phi_1}(\Omega) \rightarrow L^{\Phi_2}(\Omega)$ if*

- (i) $h(t) = 0$ for μ -almost all $t \in T(\mathfrak{B})$;
- (ii) $\sup_{n \in \mathbb{N}, \mathfrak{A}_n \subset T(\Omega)} \Phi_2 \left[\frac{1}{\Phi_1^{-1}(\mu(\mathfrak{A}_n))} \right] h(\mathfrak{A}_n) \mu(\mathfrak{A}_n) < \infty$.

Proof Assume that (i) and (ii) hold and put

$$M := \sup_{n \in \mathbb{N}, \mathfrak{A}_n \subset T(\Omega)} \Phi_2 \left[\frac{1}{\Phi_1^{-1}(\mu(\mathfrak{A}_n))} \right] h(\mathfrak{A}_n) \mu(\mathfrak{A}_n).$$

Then for each $f \in L^{\Phi_1}(\Omega)$ we have

$$\begin{aligned} I_{\Phi_2}(C_T f) &= \int_{\Omega} \Phi_2(C_T f) d\mu = \int_{T(\mathfrak{B})} h(t) \Phi_2(f(t)) d\mu + \int_{\bigcup_n T(\mathfrak{A}_n)} h(t) \Phi_2(f(t)) d\mu \\ &= \sum_{\mathfrak{A}_n \subset T(\Omega)} \int_{\mathfrak{A}_n} h(t) \Phi_2(f(t)) d\mu = \sum_{\mathfrak{A}_n \subset T(\Omega)} h(\mathfrak{A}_n) \Phi_2(f(\mathfrak{A}_n)) \mu(\mathfrak{A}_n) \\ &= \sum_{\mathfrak{A}_n \subset T(\Omega)} h(\mathfrak{A}_n) \Phi_2 \left[\Phi_1^{-1}(\mu(\mathfrak{A}_n)) \Phi_1^{-1} \circ \Phi_1(f(\mathfrak{A}_n)) \frac{1}{\Phi_1^{-1}(\mu(\mathfrak{A}_n))} \right] \mu(\mathfrak{A}_n) \\ &\leq \sum_{\mathfrak{A}_n \subset T(\Omega)} h(\mathfrak{A}_n) \Phi_2 \left[\Phi_1^{-1} \left[c_1 \mu(\mathfrak{A}_n) \Phi_1(f(\mathfrak{A}_n)) \right] \frac{1}{\Phi_1^{-1}(\mu(\mathfrak{A}_n))} \right] \mu(\mathfrak{A}_n) \\ &\leq c_2 \sum_{\mathfrak{A}_n \subset T(\Omega)} h(\mathfrak{A}_n) \Phi_2 \circ \Phi_1^{-1} \left[c_1 \mu(\mathfrak{A}_n) \Phi_1(f(\mathfrak{A}_n)) \right] \Phi_2 \left[\frac{1}{\Phi_1^{-1}(\mu(\mathfrak{A}_n))} \right] \mu(\mathfrak{A}_n) \\ &\leq c_2 \sum_{\mathfrak{A}_n \subset T(\Omega)} \Phi_2 \circ \Phi_1^{-1} \left[c_1 \mu(\mathfrak{A}_n) \Phi_1(f(\mathfrak{A}_n)) \right] M \\ &\leq c_2 M \Phi_2 \circ \Phi_1^{-1} \left[c_1 \sum_{n \in \mathbb{N}} \mu(\mathfrak{A}_n) \Phi_1(f(\mathfrak{A}_n)) \right], \end{aligned}$$

where c_1, c_2 are positive constants related to conditions $\Phi_1 \in \Delta'$ and $\Phi_2 \in \Delta'$, respectively, and the last inequality follows from the superadditivity of the convex function $\Phi_2 \circ \Phi_1^{-1}$ on the interval $[0, \infty)$. Now let $\|f\|_{\Phi_1} \leq 1$. We obtain

$$I_{\Phi_2}(C_T f) \leq c_2 M \Phi_2 \circ \Phi_1^{-1}(c_1) < \infty.$$

Hence $\|C_T f\|_{\Phi_2} \leq \max(c_2 M \Phi_2 \circ \Phi_1^{-1}(c_1), 1)$, and so C_T is bounded. □

Theorem 4.4 *Let $T : \Omega \rightarrow \Omega$ be a non-singular measurable transformation, and let Φ_1, Φ_2, Φ_3 be Young functions vanishing only at zero, taking only finite values, and such that $\Phi_1(xy) \leq \Phi_2(x) + \Phi_3(y)$ for all $x, y \geq 0$. Consider the following statements:*

- (i) T induces a bounded composition operator $C_T : L^{\Phi_1}(\Omega) \rightarrow L^{\Phi_2}(\Omega)$.
- (ii) $\mu(\mathfrak{B}) = 0$ and there is a constant $M > 0$ such that $\Phi_1^{-1} \left(\frac{1}{\mu(\mathfrak{A}_n)} \right) \leq M \Phi_2^{-1} \left(\frac{1}{h(\mathfrak{A}_n) \mu(\mathfrak{A}_n)} \right)$ for all $n \in \mathbb{N}$ such that $\mathfrak{A}_n \subset T(\Omega)$.

(iii) $h(t) = 0$ for μ -a.e. $t \in T(\mathfrak{B})$ and

$$\sup_{n \in \mathbb{N}, \mathfrak{A}_n \subset T(\Omega)} \frac{\Phi_2^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right) \Phi_3^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right)}{\Phi_2^{-1}\left(\frac{1}{h(\mathfrak{A}_n)\mu(\mathfrak{A}_n)}\right)} < \infty.$$

(iv) $h(t) = 0$ for μ -a.e. $t \in T(\mathfrak{B})$ and

$$\sup_{n \in \mathbb{N}, \mathfrak{A}_n \subset T(\Omega)} \Phi_2^{-1}(h(\mathfrak{A}_n)) \Phi_3^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right) < \infty.$$

Then (i) \Rightarrow (ii) \Rightarrow (iii). Moreover, if $\Phi_2 \in \Delta'$, then (ii) \Rightarrow (iv).

Proof To prove that (i) \Rightarrow (ii), assume that C_T is a bounded operator from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$. It implies that there is a constant $M > 0$ such that for every $n \in \mathbb{N}$,

$$\|C_T \chi_{\mathfrak{A}_n}\|_{\Phi_2} \leq M \|\chi_{\mathfrak{A}_n}\|_{\Phi_1},$$

where $\chi_{\mathfrak{A}_n}$ is the characteristic function of the atom $\mathfrak{A}_n \subset T(\Omega)$. Calculating the norm of the characteristic function $\chi_{\mathfrak{A}_n}$, we obtain

$$\frac{1}{\Phi_2^{-1}\left(\frac{1}{\mu \circ T^{-1}(\mathfrak{A}_n)}\right)} \leq \frac{M}{\Phi_1^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right)},$$

which yields the desired inequality.

By Theorem 4.1, we have $h(t) = 0$ for μ -a.e. $t \in T(\mathfrak{B})$, hence $\mu(\mathfrak{B}) = \mu \circ T^{-1}(T(\mathfrak{B})) = \int_{T(\mathfrak{B})} h(t) d\mu(t) = 0$.

Next we show that (ii) \Rightarrow (iii). For each $\mathfrak{A}_n \subset T(\Omega)$ let

$$\Phi_1^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right) \leq M \Phi_2^{-1}\left(\frac{1}{h(\mathfrak{A}_n)\mu(\mathfrak{A}_n)}\right). \tag{4.1}$$

Since $\Phi_1(xy) \leq \Phi_2(x) + \Phi_3(y)$ for all $x, y \geq 0$, we have $\Phi_3^{-1}(x)\Phi_2^{-1}(x) \leq 2\Phi_1^{-1}(x)$ for all $x \geq 0$. Plugging $x = \frac{1}{\mu(\mathfrak{A}_n)}$ in this inequality and using (4.1), we have

$$\begin{aligned} \Phi_3^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right) \Phi_2^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right) &\leq 2\Phi_1^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right) \\ &\leq 2M \Phi_2^{-1}\left(\frac{1}{h(\mathfrak{A}_n)\mu(\mathfrak{A}_n)}\right). \end{aligned} \tag{4.2}$$

Hence

$$\sup_{n \in \mathbb{N}, \mathfrak{A}_n \subset T(\Omega)} \frac{\Phi_2^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right) \Phi_3^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right)}{\Phi_2^{-1}\left(\frac{1}{h(\mathfrak{A}_n)\mu(\mathfrak{A}_n)}\right)} \leq 2M < \infty.$$

Moreover, since $\mu \circ T^{-1}(T(\mathfrak{B})) = \mu(\mathfrak{B}) = 0$, we have $h(t) = 0$ for μ -a.e. $t \in T(\mathfrak{B})$.

Finally, we prove implication (ii) \Rightarrow (iv) under the assumption that $\Phi_2 \in \Delta'$, i.e. that there is a constant $c > 0$ such that $\Phi_2^{-1}(x)\Phi_2^{-1}(y) \leq \Phi_2^{-1}(cxy)$ ($x, y \geq 0$). We may assume that $c \geq 1$. Plugging $x = 1/h(\mathfrak{A}_n)\mu(\mathfrak{A}_n)$ and $y = h(\mathfrak{A}_n)/c$ in this inequality, we get

$$\Phi_2^{-1}\left(\frac{1}{h(\mathfrak{A}_n)\mu(\mathfrak{A}_n)}\right) \leq \frac{\Phi_2^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right)}{\Phi_2^{-1}\left(\frac{h(\mathfrak{A}_n)}{c}\right)}.$$

Combining inequality (4.2) with this, yields

$$\Phi_3^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right) \Phi_2^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right) \leq 2M \Phi_2^{-1}\left(\frac{1}{h(\mathfrak{A}_n)\mu(\mathfrak{A}_n)}\right) \leq 2M \frac{\Phi_2^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right)}{\Phi_2^{-1}\left(\frac{h(\mathfrak{A}_n)}{c}\right)}.$$

Therefore

$$\Phi_2^{-1}\left(\frac{h(\mathfrak{A}_n)}{c}\right) \Phi_3^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right) \leq 2M.$$

Since Φ_2^{-1} is a concave function and $c \geq 1$, we have

$$\frac{1}{c} \Phi_2^{-1}(h(\mathfrak{A}_n)) \Phi_3^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right) \leq 2M,$$

hence $\sup_{n \in \mathbb{N}, \mathfrak{A}_n \subset T(\Omega)} \Phi_2^{-1}(h(\mathfrak{A}_n)) \Phi_3^{-1}(1/\mu(\mathfrak{A}_n)) \leq 2M c < \infty$, and this completes the proof. □

Now we present two lemmas on the relationship between multiplication operators and composition operators, which will be needed later on.

Lemma 4.5 *Let $T : \Omega \rightarrow \Omega$ be a non-singular measurable transformation, and let Φ_1, Φ_2 be Young functions vanishing only at zero and taking only finite values, with $\Phi_2 \in \nabla'$. For any function $f \in L^{\Phi_1}(\Omega)$ such that $M_{\Phi_2^{-1} \circ h} f \in L^{\Phi_2}(T(\Omega))$, we have $C_T f \in L^{\Phi_2}(\Omega)$ and the following inequality holds*

$$\|C_T f\|_{\Phi_2} \leq b \|M_{\Phi_2^{-1} \circ h} f\|_{\Phi_2},$$

where b is some positive constant.

Proof Let $f \in L^{\Phi_1}(\Omega)$ be such that $M_{\Phi_2^{-1} \circ h} f \in L^{\Phi_2}(T(\Omega))$. By the definition of the norm $\|\cdot\|_{\Phi_2}$ and the assumption that $\Phi_2 \in \nabla'$, for $b > 0$ such that $\Phi_2(bxy) \geq \Phi_2(x)\Phi_2(y)$ ($x, y \geq 0$) we have

$$\|C_T f\|_{\Phi_2} = \inf \left\{ k : \int_{\Omega} \Phi_2\left(\frac{f(T(t))}{k}\right) d\mu \leq 1 \right\}$$

$$\begin{aligned}
 &= \inf \left\{ k : \int_{T(\Omega)} h(t) \Phi_2 \left(\frac{f(t)}{k} \right) d\mu \leq 1 \right\} \\
 &= \inf \left\{ k : \int_{T(\Omega)} \Phi_2(\Phi_2^{-1}(h(t))) \Phi_2 \left(\frac{f(t)}{k} \right) d\mu \leq 1 \right\} \\
 &\leq \inf \left\{ k : \int_{T(\Omega)} \Phi_2 \left(\frac{b\Phi_2^{-1}(h(t))f(t)}{k} \right) d\mu \leq 1 \right\} \\
 &= b \inf \left\{ k/b : \int_{T(\Omega)} \Phi_2 \left(\frac{\Phi_2^{-1}(h(t))f(t)}{k/b} \right) d\mu \leq 1 \right\} \\
 &= b \|M_{\Phi_2^{-1} \circ h} f\|_{\Phi_2}.
 \end{aligned}$$

□

Corollary 4.6 *Under the assumptions of Lemma 4.5, if $M_{\Phi_2^{-1} \circ h}$ is a bounded multiplication operator from $L^{\Phi_1}(T(\Omega))$ into $L^{\Phi_2}(T(\Omega))$, then C_T is a bounded composition operator from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$.*

Lemma 4.7 *Let $T : \Omega \rightarrow \Omega$ be a non-singular measurable transformation and let Φ_1, Φ_2 be Young functions vanishing only at zero and taking only finite values, with $\Phi_2 \in \Delta'$. For any function $f \in L^{\Phi_1}(\Omega)$ such that $C_T f \in L^{\Phi_2}(\Omega)$, we have $M_{\Phi_2^{-1} \circ h} f \in L^{\Phi_2}(T(\Omega))$ and the following inequality holds*

$$\|M_{\Phi_2^{-1} \circ h} f\|_{\Phi_2} \leq c \|C_T f\|_{\Phi_2},$$

where c is some positive constant.

Proof Let $f \in L^{\Phi_1}(\Omega)$ be such that $C_T f \in L^{\Phi_2}(\Omega)$, and let $c \geq 1$ be a constant in the inequality $\Phi_2(xy) \leq c\Phi_2(x)\Phi_2(y)$ for all $x, y \geq 0$. Then

$$\begin{aligned}
 \|M_{\Phi_2^{-1} \circ h} f\|_{\Phi_2} &= \inf \left\{ k : \int_{T(\Omega)} \Phi_2 \left(\frac{\Phi_2^{-1}(h(t))f(t)}{k} \right) d\mu \leq 1 \right\} \\
 &\leq \inf \left\{ k : \int_{T(\Omega)} c \Phi_2(\Phi_2^{-1}(h(t))) \Phi_2 \left(\frac{f(t)}{k} \right) d\mu \leq 1 \right\} \\
 &= \inf \left\{ k : \int_{T(\Omega)} c h(t) \Phi_2 \left(\frac{f(t)}{k} \right) d\mu \leq 1 \right\} \\
 &= \inf \left\{ k : \int_{\Omega} c \Phi_2 \left(\frac{f(T(t))}{k} \right) d\mu \leq 1 \right\} \\
 &\leq \inf \left\{ k : \int_{\Omega} \Phi_2 \left(\frac{f(T(t))}{k/c} \right) d\mu \leq 1 \right\} \\
 &= c \inf \left\{ k/c : \int_{\Omega} \Phi_2 \left(\frac{f(T(t))}{k/c} \right) d\mu \leq 1 \right\} \\
 &= c \|C_T f\|_{\Phi_2},
 \end{aligned}$$

as required. □

Corollary 4.8 *Under the assumptions of Lemma 4.7, if C_T is a bounded composition operator from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$, then $M_{\Phi_2^{-1} \circ h}$ is a bounded multiplication operator from $L^{\Phi_1}(T(\Omega))$ into $L^{\Phi_2}(T(\Omega))$.*

Following [38], for any F in the σ -algebra $\Sigma \cap T(\Omega)$, we define

$$Q_T(F) := \inf\{b \geq 0 : \mu \circ T^{-1}(E) \leq b \mu(E), E \in \Sigma \cap T(\Omega), E \subset F\}.$$

To prove our next results, we will need the following two lemmas.

Lemma 4.9 [38, Lemma 3.6] *For any $F \in \Sigma \cap T(\Omega)$, we have $Q_T(F) = \text{ess sup}_{t \in F} h(t)$, where $h = d(\mu \circ T^{-1})/d\mu$.*

Lemma 4.10 *Let T be a measurable transformation of the measure space (Ω, Σ, μ) into itself and let Φ be a Young function. Then*

$$\int_{T(\Omega)} \Phi \circ h \, d\mu = \inf \left\{ \sum_{j=1}^{\infty} \Phi(Q_T(F_j))\mu(F_j) : \{F_j\} \in \mathcal{P}_{T(\Omega)} \right\},$$

where $\mathcal{P}_{T(\Omega)}$ is the family of all measurable partitions of $T(\Omega)$ and the case $\infty = \infty$ is admissible.

Proof Let $I := \inf\{\sum_{j=1}^{\infty} \Phi(Q_T(F_j))\mu(F_j) : \{F_j\} \in \mathcal{P}_{T(\Omega)}\}$. By Lemma 4.9, for the partition $\{F_j\}$ of $T(\Omega)$, we have

$$\begin{aligned} \int_{T(\Omega)} \Phi \circ h \, d\mu &= \sum_{j=1}^{\infty} \int_{F_j} \Phi(h(t)) \, d\mu \leq \sum_{j=1}^{\infty} \Phi(\text{ess sup}_{t \in F_j} h(t))\mu(F_j) \\ &= \sum_{j=1}^{\infty} \Phi(Q_T(F_j))\mu(F_j). \end{aligned}$$

Hence

$$\int_{T(\Omega)} \Phi \circ h \, d\mu \leq I.$$

To prove the reverse inequality, let $a > 1$ be an arbitrary number. For each $m \in \mathbb{Z}$ define the set

$$G_m := \{t \in T(\Omega) : a^{m-1} \leq \Phi(h(t)) < a^m\}.$$

Let $\{F_j\}_{j=1}^{\infty}$ be any rearrangement of the sets $\{G_m\}_{m=-\infty}^{\infty}$ and the set $\{t \in T(\Omega) : h(t) = 0\}$. Clearly, $\{F_j\}_{j=1}^{\infty}$ is a partition of $T(\Omega)$. By Lemma 4.9, we have

$$\begin{aligned}
 I &\leq \sum_{j=1}^{\infty} \Phi(Q_T(F_j))\mu(F_j) = \sum_{j=1}^{\infty} \Phi(\operatorname{ess\,sup}_{t \in F_j} h(t))\mu(F_j) \\
 &= \sum_{m=-\infty}^{\infty} \Phi(\operatorname{ess\,sup}_{t \in G_m} h(t))\mu(G_m) \leq \sum_{m=-\infty}^{\infty} a^m \mu(G_m) \\
 &= a \sum_{m=-\infty}^{\infty} a^{m-1} \mu(G_m) = a \sum_{m=-\infty}^{\infty} \int_{G_m} a^{m-1} d\mu \\
 &\leq a \sum_{m=-\infty}^{\infty} \int_{G_m} \Phi \circ h d\mu = a \sum_{j=1}^{\infty} \int_{F_j} \Phi \circ h d\mu \\
 &= a \int_{T(\Omega)} \Phi \circ h d\mu.
 \end{aligned}$$

Since this holds for any $a > 1$, the result follows. □

Proposition 3.2 and Lemma 4.7 allow us to give a necessary condition for the boundedness of the composition operator C_T . Statement (ii) of the next theorem is interesting, because under some assumptions on the Young functions Φ_1, Φ_2 it gives the information that if C_T acts continuously from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$, then the Radon–Nikodym derivative h must be from a concrete Orlicz space $L^{\Phi}(\Omega)$, where Φ depends on Φ_1 and Φ_2 .

Theorem 4.11 *Let T be a non-singular measurable transformation of the measure space (Ω, Σ, μ) into itself. Then the following assertions hold.*

- (i) *If Φ is a Young function, then the Radon–Nikodym derivative $h = d(\mu \circ T^{-1})/d\mu$ belongs to $L^{\Phi}(T(\Omega))$ if and only if there exists a partition $\{F_j\}_{j=1}^{\infty}$ of $T(\Omega)$ and $\lambda > 0$ such that $\sum_{j=1}^{\infty} \Phi(\lambda Q_T(F_j))\mu(F_j) < \infty$.*
- (ii) *Let Φ_1 be an N -function, Φ_2 a Young function vanishing only at zero and taking only finite values, such that $\Phi_1 \in \Delta', \Phi_2 \in \Delta_2$, and $\Phi_3 := \Psi_2 \circ \Psi_1^{-1}$ is a Young function. If T induces a bounded composition operator $C_T : L^{\Phi_1}(\Omega) \rightarrow L^{\Phi_2}(\Omega)$, then the Radon–Nikodym derivative $h = d(\mu \circ T^{-1})/d\mu$ belongs to $L^{\Psi_3 \circ \Psi_1 \circ \Phi_2^{-1}}(T(\Omega))$.*

Proof (i) It is an easy consequence of Lemma 4.10 with the function $\Phi_{\lambda}(x) := \Phi(\lambda x)$ in place of the Young function Φ .

(ii) Since, by assumption, C_T is a bounded composition operator from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$, Lemma 4.7 implies that

$$M_{\Phi_2^{-1} \circ h} : L^{\Phi_1}(T(\Omega)) \rightarrow L^{\Phi_2}(T(\Omega))$$

is a bounded multiplication operator. Thus, by Proposition 3.2, for the Young function $\Phi_3 = \Psi_2 \circ \Psi_1^{-1}$, we have $\Phi_2^{-1} \circ h \in L^{\Psi_3 \circ \Psi_1}(T(\Omega))$, and so

$$\int_{T(\Omega)} \Psi_3 \circ \Psi_1 \circ \lambda \Phi_2^{-1} \circ h d\mu < \infty,$$

for some constant $\lambda > 0$. Since $\Psi_3 \circ \Psi_1$ is a nondecreasing function on the interval $[0, \infty)$, we may assume that $0 < \lambda < 1$. It can be easily shown that the Δ_2 -condition assumed for the function Φ_2 is equivalent to the condition: for all $\lambda \in (0, 1)$ there is a constant $K > 0$ such that $\Phi_2^{-1}(Kx) \leq \lambda \Phi_2^{-1}(x)$ ($x \geq 0$). Since $\Psi_3 \circ \Psi_1$ is a nondecreasing function on the interval $[0, \infty)$, we then get

$$\int_{T(\Omega)} \Psi_3 \circ \Psi_1 \circ \Phi_2^{-1} \circ Kh \, d\mu < \infty.$$

Thus $h \in L^{\Psi_3 \circ \Psi_1 \circ \Phi_2^{-1}}(T(\Omega))$. □

Finally, we give a sufficient condition for the composition operator C_T to be bounded from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$ when $\Phi_1 \circ \Phi_2^{-1}$ is a Young function.

Theorem 4.12 *Let Φ_1, Φ_2 be Young functions vanishing only at zero and taking only finite values. If $\Phi_3 = \Phi_1 \circ \Phi_2^{-1}$ is a Young function and $h \in L^{\Psi_3}(T(\Omega))$, then T induces a bounded composition operator $C_T : L^{\Phi_1}(\Omega) \rightarrow L^{\Phi_2}(\Omega)$.*

Proof Since

$$\begin{aligned} I_{\Phi_3}(\Phi_2 \circ f) &= \int_{\Omega} \Phi_3 \circ \Phi_2 \circ f \, d\mu = \int_{\Omega} \Phi_1 \circ \Phi_2^{-1} \circ \Phi_2 \circ f \, d\mu \\ &= \int_{\Omega} \Phi_1 \circ f \, d\mu = I_{\Phi_1}(f), \end{aligned}$$

we have $f \in L^{\Phi_1}(\Omega)$ if and only if $\Phi_2 \circ f \in L^{\Phi_3}(\Omega)$, and furthermore that $\|\Phi_2 \circ f\|_{\Phi_3} = \|f\|_{\Phi_1}$. Hence an application of Hölder’s inequality yields

$$\begin{aligned} I_{\Phi_2}(C_T f) &= \int_{\Omega} \Phi_2 \circ f \circ T \, d\mu = \int_{T(\Omega)} h \Phi_2 \circ f \, d\mu \\ &\leq 2\|h\|_{\Psi_3} \|\Phi_2 \circ f\|_{\Phi_3} = 2\|h\|_{\Psi_3} \|f\|_{\Phi_1}. \end{aligned}$$

Thus $\|C_T\| \leq \max(2\|h\|_{\Psi_3}, 1)$, and this completes the proof. □

Remark 4.13 We conclude this section with some applications of the theorems proved.

1. Taking $\Phi_1(x) = |x|^p/p$ and $\Phi_2(x) = |x|^q/q$, where $1 < p < q < \infty$, by Theorems 4.1 and 4.4, we obtain that if C_T is a composition operator induced by the non-singular measurable transformation $T : \Omega \rightarrow \Omega$, then the following statements are equivalent:
 - (a) C_T is bounded from $L^p(\Omega)$ into $L^q(\Omega)$.
 - (b) $h(t) = 0$ for μ -a.e. $t \in T(\mathfrak{B})$ and $\sup_{n \in \mathbb{N}, \mathfrak{A}_n \subset T(\Omega)} \frac{h(\mathfrak{A}_n)^p}{\mu(\mathfrak{A}_n)^{q-p}} < \infty$.
 - (c) $\mu(\mathfrak{B}) = 0$ and there is a constant $k > 0$ such that $\mu \circ T^{-1}(\mathfrak{A}_n)^p \leq k\mu(\mathfrak{A}_n)^q$ for all $n \in \mathbb{N}, \mathfrak{A}_n \subset T(\Omega)$.
2. Similarly, taking $\Phi_1(x) = |x|^p/p$ and $\Phi_2(x) = |x|^q/q$, where $1 < q < p < \infty$, by Theorems 4.11 and 4.12, we obtain that if C_T is a composition operator induced by the non-singular measurable transformation $T : \Omega \rightarrow \Omega$, then the following statements are equivalent:

- (a) C_T is a bounded operator from $L^p(\Omega)$ into $L^q(\Omega)$.
- (b) $h \in L^{\frac{r}{q}}(T(\Omega))$, where $p^{-1} + r^{-1} = q^{-1}$.
- (c) There exists a partition $\{F_j\}_{j=1}^\infty$ of $T(\Omega)$ such that $\sum_{j=1}^\infty Q_T(F_j)^{\frac{r}{q}} \mu(F_j) < \infty$.

These characterizations are due to Takagi and Yokouchi [38].

5 Multiplication and composition operators with closed-range and/or finite rank

In this section we are going to investigate closed-range multiplication and composition operators between distinct Orlicz spaces.

We start with a basic observation concerning Young functions.

Lemma 5.1 *Let Φ_1, Φ_2 be Young functions such that Φ_1 is an N -function and $\Phi_3 := \Psi_2 \circ \Psi_1^{-1}$ is a Young function. If $\Psi_1 \in \Delta'$, then there is a constant $b > 0$ such that $b \Psi_1(xy) \leq \Psi_2(x) + \Psi_3(\Psi_1(y))$ for all $x, y \geq 0$.*

Proof By the definition of the function Φ_3 , we have

$$\Psi_3(y) = \sup\{x|y| - \Phi_3(x) : x \geq 0\} = \sup\{x|y| - \Psi_2 \circ \Psi_1^{-1}(x) : x \geq 0\}.$$

Since Φ_1 is an N -function, Ψ_1 is again an N -function, and $a_{\Psi_1} = 0, b_{\Psi_1} = \infty$. So the usual inverse function Ψ_1^{-1} exists. By substitution, we have

$$\Psi_3(\Psi_1(y)) = \sup\{x\Psi_1(y) - \Psi_2 \circ \Psi_1^{-1}(x) : x \geq 0\},$$

hence

$$\Psi_3(\Psi_1(y)) = \sup\{\Psi_1(x)\Psi_1(y) - \Psi_2(x) : x \geq 0\}. \tag{5.1}$$

Since $\Psi_1 \in \Delta'$, i.e. $\Psi_1(xy) \leq c\Psi_1(x)\Psi_1(y)$ for some $c > 0$ and all $x, y \geq 0$, we get from (5.1) that

$$\Psi_3(\Psi_1(y)) \geq \frac{1}{c}\Psi_1(xy) - \Psi_2(x),$$

for all $x, y \geq 0$, whence the desired inequality with $b = 1/c$ follows. □

Now we characterize closed-range multiplication operators $M_u : L^{\Phi_1}(\Omega) \rightarrow L^{\Phi_2}(\Omega)$ under the assumption $\Phi_2(xy) \leq \Phi_1(x) + \Phi_3(y)$ for all $x, y \geq 0$.

Theorem 5.2 *Let Φ_1, Φ_2, Φ_3 , be Young functions vanishing only at zero, taking only finite values, and such that $\Phi_2, \Phi_3 \in \Delta_2$ and $\Phi_2(xy) \leq \Phi_1(x) + \Phi_3(y)$ for all $x, y \geq 0$. If $u \in L^{\Phi_3}(\Omega)$, then M_u is a bounded multiplication operator from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$ and the following assertions are equivalent:*

- (a) $u(t) = 0$ for μ -a.e. $t \in \mathfrak{B}$ and the set $E = \{n \in \mathbb{N} : u(\mathfrak{A}_n) \neq 0\}$ is finite.
- (b) M_u has finite rank.
- (c) M_u has closed-range.

Proof Let $u \in L^{\Phi_3}(\Omega)$ and $f \in L^{\Phi_1}(\Omega)$, i.e. there are constants $\lambda, \alpha > 0$ such that $I_{\Phi_3}(\lambda u) < \infty$ and $I_{\Phi_1}(\alpha f) < \infty$. Therefore, by the inequality $\Phi_2(xy) \leq \Phi_1(x) + \Phi_3(y)$ ($x, y \geq 0$), we have

$$I_{\Phi_2}(\alpha \lambda u f) \leq I_{\Phi_1}(\alpha f) + I_{\Phi_3}(\lambda u) < \infty,$$

which implies that $u f \in L^{\Phi_2}(\Omega)$.

Let S be the support of u . We may assume that $\mu(S) > 0$ since otherwise M_u is a zero operator and there is nothing to prove.

We prove the implication (a) \Rightarrow (b). Assume that (a) holds. Hence there is $r \in \mathbb{N}$ such that

$$S = \bigcup_{n \in E} \mathfrak{A}_n = \mathfrak{A}_{n_1} \cup \dots \cup \mathfrak{A}_{n_r}.$$

Since $\Phi_2 \in \Delta_2$, the set $\{\chi_{\mathfrak{A}_{n_1}}, \dots, \chi_{\mathfrak{A}_{n_r}}\}$ of characteristic functions generates the subspace

$$\{g \in L^{\Phi_2}(\Omega) : g(x) = 0 \text{ for } \mu - \text{a.e. } t \in \Omega \setminus S\} \cong L^{\Phi_2}(S).$$

The range of M_u is contained in the r -dimensional subspace $L^{\Phi_2}(S)$, hence M_u has finite rank.

- (b) \Rightarrow (c) . If the range of the operator M_u in $L^{\Phi_2}(\Omega)$ is finite-dimensional, then it is also closed, as any finite-dimensional subspace of a Banach space is a closed subspace of this space.
- (c) \Rightarrow (a) . Let M_u have closed-range and assume that $\mu\{t \in \mathfrak{B} : u(t) \neq 0\} > 0$. Then there is $\delta > 0$ such that the set $G = \{t \in \mathfrak{B} : u(t) \geq \delta\}$ has positive measure. It is easy to see that the restriction $u|_G$ induces a bounded multiplication operator $M_{u|_G}$ from $L^{\Phi_1}(G)$ into $L^{\Phi_2}(G)$, and that if M_u has closed-range, then $M_{u|_G}$ has closed-range as well.

We will show that $M_{u|_G}(L^{\Phi_1}(G)) = L^{\Phi_2}(G)$. Let A be any measurable subset of G with $\mu(A) < \infty$, and define the function $f_A := \frac{1}{u|_G} \chi_A$. We get

$$I_{\Phi_1}(f_A) = \int_G \Phi_1 \circ f_A \, d\mu = \int_A \Phi_1 \circ \frac{1}{u|_G} \, d\mu \leq \Phi_1(1/\delta) \mu(A) < \infty,$$

and so $f_A \in L^{\Phi_1}(G)$. Moreover, $M_{u|_G} f_A = \chi_A$, which implies that the linear space $M_{u|_G}(L^{\Phi_1}(G))$ contains the set \mathcal{F} of all linear combinations of characteristic functions of measurable subsets of G with positive and finite measure.

Now \mathcal{F} is a dense subset of $L^{\Phi_2}(G)$ ([26, Lemma 4.1]) and, by assumption, $M_{u|_G}(L^{\Phi_1}(G))$ is a closed subspace of $L^{\Phi_2}(G)$, therefore

$$L^{\Phi_2}(G) = \overline{\mathcal{F}} \subset M_{u|_G}(L^{\Phi_1}(G)) \subset L^{\Phi_2}(G),$$

and so $M_{u|_G}(L^{\Phi_1}(G)) = L^{\Phi_2}(G)$, as claimed.

Consequently, we can define the inverse multiplication operator

$$M_{\frac{1}{u|_G}} : L^{\Phi_2}(G) \rightarrow L^{\Phi_1}(G), \quad M_{\frac{1}{u|_G}} f := \frac{1}{u|_G} f.$$

The operator $M_{\frac{1}{u|_G}}$ is bounded and, since $\Phi_2(xy) \leq \Phi_1(x) + \Phi_3(y)$ ($x, y \geq 0$), we can apply Theorem 3.4 to conclude that $\frac{1}{u(t)} = 0$ for μ -a.e. $t \in G$, which is absurd. This contradiction shows that $u(t) = 0$ for μ -a.e. $t \in \mathfrak{B}$.

Next we show that the set $E = \{n \in \mathbb{N} : u(\mathfrak{A}_n) \neq 0\}$ is finite if M_u has closed range. If $E = \emptyset$, we have nothing to prove. So let us assume that $E \neq \emptyset$. Define $S = \bigcup_{n \in E} \mathfrak{A}_n$.

Analogously as above, we can show that $M_{u|_S}(L^{\Phi_1}(S)) = L^{\Phi_2}(S)$. Indeed, let A be a subset of S with $\mu(A) < \infty$. Define the function $f_A := \frac{1}{u|_S} \chi_A$. The set A having finite measure, we get

$$I_{\Phi_1}(f_A) = \int_A \Phi_1 \circ \frac{1}{u|_S} d\mu = \sum_{\mathfrak{A}_n \subset A} \Phi_1(1/u(\mathfrak{A}_n))\mu(\mathfrak{A}_n) < \infty.$$

Hence $f_A \in L^{\Phi_1}(S)$. Since $M_{u|_S} f_A = \chi_A$, we conclude that $M_{u|_S}(L^{\Phi_1}(S))$ contains the set ℓ_f^0 of all linear combinations of characteristic functions of subsets of S with positive and finite measure.

Now ℓ_f^0 is a dense subset of $L^{\Phi_2}(S)$, and $M_{u|_S}(L^{\Phi_1}(S))$ is a closed subspace of $L^{\Phi_2}(S)$, which implies that

$$L^{\Phi_2}(S) = \overline{\ell_f^0} \subset M_{u|_S}(L^{\Phi_1}(S)) \subset L^{\Phi_2}(S),$$

and so $M_{u|_S}(L^{\Phi_1}(S)) = L^{\Phi_2}(S)$.

We can thus define a bounded multiplication operator $M_{\frac{1}{u|_S}}$ from $L^{\Phi_2}(S)$ into $L^{\Phi_1}(S)$. Applying Theorem 3.4 to the operator $M_{\frac{1}{u|_S}}$, we obtain

$$\sup_{n \in E} \frac{1}{u(\mathfrak{A}_n)} \Phi_3^{-1} \left(\frac{1}{\mu(\mathfrak{A}_n)} \right) < \infty.$$

Let $C = \sup_{n \in E} \frac{1}{u(\mathfrak{A}_n)} \Phi_3^{-1} \left(\frac{1}{\mu(\mathfrak{A}_n)} \right) > 0$. Since $E \neq \emptyset$ and $1 \leq \Phi_3(Cu(\mathfrak{A}_n)) \mu(\mathfrak{A}_n)$ for all $n \in E$, we have

$$\sum_{n \in E} 1 \leq \sum_{n \in E} \Phi_3(Cu(\mathfrak{A}_n)) \mu(\mathfrak{A}_n) = \sum_{n \in E} \int_{\mathfrak{A}_n} \Phi_3 \circ Cu \, d\mu \leq \int_{\Omega} \Phi_3 \circ Cu \, d\mu < \infty,$$

where the final inequality follows from the assumption that $\Phi_3 \in \Delta_2$. Thus E must be finite. □

In the next theorem we characterize closed-range multiplication operators $M_u : L^{\Phi_1}(\Omega) \rightarrow L^{\Phi_2}(\Omega)$ under the condition that $\Phi_1(xy) \leq \Phi_2(x) + \Phi_3(y)$ for all $x, y \geq 0$.

Theorem 5.3 *Let Φ_1, Φ_2, Φ_3 be Young functions vanishing only at zero, taking only finite values, and such that $\Phi_2, \Phi_3 \in \Delta_2$ and $\Phi_1(xy) \leq \Phi_2(x) + \Phi_3(y)$ for all $x, y \geq 0$. If M_u is a bounded multiplication operator from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$ and $\frac{1}{u} \in L^{\Phi_3}_+(\Omega)$, then the following statements are equivalent:*

- (a) *The set $E = \{n \in \mathbb{N} : u(\mathfrak{A}_n) \neq 0\}$ is finite.*
- (b) *M_u has finite rank.*
- (c) *M_u has closed-range.*

Proof By Theorem 3.4, we have that $u(t) = 0$ for μ -a.e. $t \in \mathfrak{B}$. The proofs of implications (a) \Rightarrow (b) and (b) \Rightarrow (c) are as in the proof of Theorem 5.2.

We prove the implication (c) \Rightarrow (a). Using the same notation as in the proof of Theorem 5.2, let $S = \bigcup_{n \in E} \mathfrak{A}_n$ and $E \neq \emptyset$. Since $M_u : L^{\Phi_1}(S) \rightarrow L^{\Phi_2}(S)$ is bounded, by Theorem 3.4, we have

$$\sup_{n \in E} u(\mathfrak{A}_n) \Phi_3^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right) < \infty$$

Let $C = \sup_{n \in E} u(\mathfrak{A}_n) \Phi_3^{-1}\left(\frac{1}{\mu(\mathfrak{A}_n)}\right) > 0$. Since $E \neq \emptyset, 1 \leq \Phi_3\left(\frac{C}{u(\mathfrak{A}_n)}\right) \mu(\mathfrak{A}_n)$ for all $n \in E$, and $\Phi_3 \in \Delta_2$, we may write

$$\sum_{n \in E} 1 \leq \sum_{n \in E} \Phi_3\left(\frac{C}{u(\mathfrak{A}_n)}\right) \mu(\mathfrak{A}_n) = \sum_{n \in E} \int_{\mathfrak{A}_n} \Phi_3 \circ \frac{C}{u} \, d\mu \leq \int_{\Omega} \Phi_3 \circ \frac{C}{u} \, d\mu < \infty.$$

Thus E is finite. □

Now we characterize closed-range composition operators $C_T : L^{\Phi_1}(\Omega) \rightarrow L^{\Phi_2}(\Omega)$ under the assumption that $\Phi_2(xy) \leq \Phi_1(x) + \Phi_3(y)$ for all $x, y \geq 0$.

We will need an elementary lemma with an easy proof which we omit.

Lemma 5.4 *Let Φ_1, Φ_2 be Young functions vanishing only at zero and taking only finite values, and let T be a non-singular measurable transformation of Ω such that C_T is a bounded composition operator from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$. If T is surjective, then C_T is injective.*

Theorem 5.5 *Let Φ_1, Φ_2, Φ_3 be Young functions vanishing only at zero, taking only finite values, and such that $\Phi_2 \in \nabla' \cap \Delta_2, \Phi_3 \in \Delta_2$, and $\Phi_2(xy) \leq \Phi_1(x) + \Phi_3(y)$ for*

all $x, y \geq 0$. Let T be a surjective non-singular measurable transformation of Ω such that $\Phi_2^{-1} \circ h \in L^{\Phi_3}(T(\Omega))$. If $C_T : L^{\Phi_1}(\Omega) \rightarrow L^{\Phi_2}(\Omega)$ is a bounded composition operator, then the following assertions are equivalent:

- (a) C_T has closed-range.
- (b) $h(t) = 0$ for μ -a.e. $t \in T(\mathfrak{B})$ and the set $\{n \in \mathbb{N} : \mathfrak{A}_n \subset T(\Omega), h(\mathfrak{A}_n) \neq 0\}$ is finite.
- (c) $\mu(\mathfrak{B}) = 0$ and the set $\{n \in \mathbb{N} : \mathfrak{A}_n \subset T(\Omega), \mu \circ T^{-1}(\mathfrak{A}_n) \neq 0\}$ is finite.
- (d) C_T has finite rank.

Proof The implications (b) \Rightarrow (c) and (d) \Rightarrow (a) are obvious.

We prove the implication (a) \Rightarrow (b). Assume that C_T has closed-range. Since T is surjective, by Lemma 5.4, C_T is injective. It is a well-known fact that an injective operator has closed-range if and only if it is bounded away from zero (see [4], III.12.5), hence C_T is bounded away from zero. This, combined with Lemma 4.5, yields that the multiplication operator $M_{\Phi_2^{-1} \circ h}$ is bounded away from zero and therefore (being also injective) has closed-range. Now an application of Theorem 5.2 shows that $\Phi_2^{-1} \circ h(t) = 0$ for μ -a.e. $t \in T(\mathfrak{B})$ and the set $\{n \in \mathbb{N} : \mathfrak{A}_n \subset T(\Omega), \Phi_2^{-1} \circ h(\mathfrak{A}_n) \neq 0\}$ is finite. We conclude that $h(t) = 0$ for μ -a.e. $t \in T(\mathfrak{B})$ and the set $\{n \in \mathbb{N} : \mathfrak{A}_n \subset T(\Omega), h(\mathfrak{A}_n) \neq 0\}$ is finite.

Finally, we prove implication (c) \Rightarrow (d). Assume (c) holds. It is easy to see that the range of the operator C_T is then contained in a subspace generated by the functions $\{\chi_{T^{-1}(\mathfrak{A}_n)} : \mathfrak{A}_n \subset T(\Omega), \mu \circ T^{-1}(\mathfrak{A}_n) \neq 0\}$. Since the set $\{n \in \mathbb{N} : \mathfrak{A}_n \subset T(\Omega), \mu \circ T^{-1}(\mathfrak{A}_n) \neq 0\}$ is finite, the subspace $C_T(L^{\Phi_1}(\Omega))$ is finite-dimensional and so C_T has finite rank. □

As an easy consequence of this theorem we state

Corollary 5.6 *If Ω is non-atomic, then, under the assumptions of Theorem 5.5, there are no non-zero closed-range composition operators from $L^{\Phi_1}(\Omega)$ into $L^{\Phi_2}(\Omega)$.*

In the next theorem we characterize closed-range composition operators $C_T : L^{\Phi_1}(\Omega) \rightarrow L^{\Phi_2}(\Omega)$ in the case when $\Phi_1(xy) \leq \Phi_2(x) + \Phi_3(y)$ for all $x, y \geq 0$ (note that compared to Theorem 5.5 the roles of the functions Φ_1 and Φ_2 are reversed).

Theorem 5.7 *Let Φ_1, Φ_2, Φ_3 be Young functions vanishing only at zero, taking only finite values, and such that $\Phi_2 \in \nabla' \cap \Delta_2, \Phi_3 \in \Delta_2$, and $\Phi_1(xy) \leq \Phi_2(x) + \Phi_3(y)$ for all $x, y \geq 0$. Let T be a surjective non-singular measurable transformation of Ω such that $\frac{1}{\Phi_2^{-1} \circ h} \in L^{\Phi_3}(\Omega)$. If $C_T : L^{\Phi_1}(\Omega) \rightarrow L^{\Phi_2}(\Omega)$ is a bounded composition operator, then the following assertions are equivalent:*

- (a) C_T has closed-range.
- (b) The set $\{n \in \mathbb{N} : \mathfrak{A}_n \subset T(\Omega), h(\mathfrak{A}_n) \neq 0\}$ is finite.
- (c) The set $\{n \in \mathbb{N} : \mathfrak{A}_n \subset T(\Omega), \mu \circ T^{-1}(\mathfrak{A}_n) \neq 0\}$ is finite.
- (d) C_T has finite rank.

Proof Using Lemma 4.5 and Theorem 5.3 and a method similar as in the proof of Theorem 5.5 proves the theorem. □

Remark 5.8 As an applications of our results we derive characterizations of bounded and closed-range multiplication and composition operators in the special case of L^p -spaces. These results recover the main theorems in [38].

1. If for $1 < p < q < \infty$ the multiplication operator M_u is bounded from $L^p(\Omega)$ into $L^q(\Omega)$, then the following assertions are equivalent:
 - (a) M_u has closed-range.
 - (b) M_u has finite rank.
 - (c) The set $\{n \in \mathbb{N} : u(\mathfrak{A}_n) \neq 0\}$ is finite.
2. If for $1 < q < p < \infty$ the multiplication operator M_u is bounded from $L^p(\Omega)$ into $L^q(\Omega)$, then the following assertions are equivalent:
 - (a) M_u has closed-range.
 - (b) M_u has finite rank.
 - (c) $u(t) = 0$ for μ -almost all $t \in \mathfrak{B}$, and the set $\{n \in \mathbb{N} : u(\mathfrak{A}_n) \neq 0\}$ is finite.
3. If for $1 < p < q < \infty$ the composition operator C_T is bounded from $L^p(\Omega)$ into $L^q(\Omega)$, then the following assertions are equivalent:
 - (a) C_T has closed-range.
 - (b) C_T has finite rank.
 - (c) The set $\{n \in \mathbb{N} : h(\mathfrak{A}_n) \neq 0\}$ is finite.
 - (d) The set $\{n \in \mathbb{N} : \mu \circ T^{-1}(\mathfrak{A}_n) \neq 0\}$ is finite.
4. If for $1 < q < p < \infty$ the composition operator C_T is bounded from $L^p(\Omega)$ into $L^q(\Omega)$, then the following assertions are equivalent:
 - (a) C_T has closed-range.
 - (b) C_T has finite rank.
 - (c) $h(t) = 0$ for μ -almost all $t \in \mathfrak{B}$, and the set $\{n \in \mathbb{N} : h(\mathfrak{A}_n) \neq 0\}$ is finite.
 - (d) $\mu \circ T^{-1}(\mathfrak{B}) = 0$ and the set $\{n \in \mathbb{N} : \mu \circ T^{-1}(\mathfrak{A}_n) \neq 0\}$ is finite.

6 Simultaneously necessary and sufficient conditions for continuity of the multiplication and composition operators between distinct Orlicz spaces

Up to this point no conditions that would be simultaneously necessary and sufficient for the continuity of the operators M_u and C_T have been given; the provided necessary conditions and sufficient conditions though have the merit of being quite easy to verify for a pair or a triple of Young functions involved. In order to obtain simultaneously necessary and sufficient conditions for continuity of the operators M_u and C_T , it is convenient to take advantage of the theory of embeddings between Musielak–Orlicz spaces. In the case of non-atomic measure spaces such conditions were presented in [3]; now we will present them for purely atomic and mixed measure spaces. However, there is a price: the simultaneously necessary and sufficient conditions for the continuity of the operators M_u and C_T are more complicated and not as easy to verify as those presented in the preceding part of this paper.

A function Φ acting from $\Omega \times \mathbb{R}$ into $[0, \infty)$ such that $\Phi(t, \cdot)$ is an Orlicz function (i.e. a finite-valued continuous Young function) for μ -a.e. $t \in \Omega$ and $\Phi(\cdot, x)$ is a Σ -measurable function for every $x \in \mathbb{R}$, is called a *generalized Orlicz function* or a *Musielak–Orlicz function*. The Musielak–Orlicz space $L^\Phi = L^\Phi(\Omega, \Sigma, \mu)$ is

the space of all (equivalence classes of) Σ -measurable functions $f : \Omega \rightarrow \mathbb{R}$ such that

$$I_\Phi(kf) := \int_\Omega \Phi(t, kf(t))d\mu < \infty$$

for some $k > 0$ depending on f . The Musielak–Orlicz space furnished with the Luxemburg norm

$$\|f\|_\Phi = \inf \left\{ \lambda > 0 : I_\Phi \left(\frac{f}{\lambda} \right) \leq 1 \right\}$$

is a Banach space (cf. [26, 28]). It is obvious that Orlicz functions are Musielak–Orlicz functions and, consequently, Orlicz spaces are Musielak–Orlicz spaces. Moreover, an Orlicz weighted space $L_w^\Phi(\Omega)$ is a Musielak–Orlicz space $L^{\Phi_w}(\Omega)$ with $\Phi_w(t, x) = \Phi(x)w(t)$ for any $t \in \Omega$ and any $x \in \mathbb{R}$.

When the measure space (Ω, Σ, μ) is purely atomic with infinite number of atoms, we may restrict ourselves to counting measure. Since (Ω, Σ, μ) is σ -finite, it has countably many atoms, and so, without loss of generality, it may be viewed as having the form $(\mathbb{N}, 2^\mathbb{N}, \mu_c)$, where μ_c is the counting measure on $2^\mathbb{N}$. Indeed, considering atoms as singletons, we write $\mu(\{n\}) = a_n$ for any $n \in \mathbb{N}$, where $\{a_n\}_{n \in \mathbb{N}}$ is a sequence of positive real numbers. If ℓ^0 denotes the space of all real sequences, then, given the Musielak–Orlicz function $\Phi = (\Phi_n)_{n=1}^\infty$, i.e. a sequence of Young functions Φ_n , we can define on ℓ^0 the absolutely convex modular

$$I_\Phi(f) = \sum_{n=1}^\infty \Phi_n(f_n)a_n \quad (f = (f_n) \in \ell^0).$$

Defining

$$B_\Phi = \{f \in \ell^0 : I_\Phi(f) \leq 1\}$$

and

$$\ell^\Phi(\mathbb{N}) = \{f \in \ell^0 : I_\Phi(kf) < \infty \text{ for some } k > 0\},$$

we easily see that B_Φ is an absolutely convex set in $\ell^\Phi(\mathbb{N})$ and that all elements of $\ell^\Phi(\mathbb{N})$ are absorbed by B_Φ . Consequently, the Minkowski functional generated by B_Φ

$$m_\Phi(f) := \inf\{\lambda > 0 : f/\lambda \in B_\Phi\}$$

is a seminorm on $\ell^\Phi(\mathbb{N})$. If we assume that for all $n \in \mathbb{N}$ the functions Φ_n are not identically equal to zero, then it is easy to prove that $m_\Phi(f) = 0$ if and only if $f = 0$. Of course,

$$m_\Phi(f) = \inf\{\lambda > 0 : I_\Phi(f/\lambda) \leq 1\}.$$

This is the Luxemburg norm $\|f\|_\Phi = m_\Phi(f)$.

Let us define a new Musielak–Orlicz function $\Psi = (\Psi_n)_{n=1}^\infty$ with

$$\Psi_n(x) = \Phi_n(x) \mu(\{n\}) = \Phi_n(x) a_n,$$

where $\{n\}$ is identified with the n th atom of the measure μ . Then the Musielak–Orlicz space generated by the original Musielak–Orlicz function Φ and the original purely atomic measure with $\mu(\{n\}) = a_n$ is isometric to the Musielak–Orlicz space generated by the Musielak–Orlicz function Ψ defined above and corresponding to counting measure with $\mu_c(\{n\}) = 1$ for all $n \in \mathbb{N}$, because of the equality

$$I_\Psi(f, \mu_c) = \sum_{n=1}^\infty \Psi_n(f_n) = \sum_{n=1}^\infty \Phi_n(f_n) a_n = I_\Phi(f, \mu).$$

For these reasons Musielak–Orlicz sequence spaces, without loss of generality, may be considered (and usually are) over the counting measure space $(\mathbb{N}, 2^\mathbb{N}, \mu_c)$.

Assume that $(\mathbb{N}, 2^\mathbb{N}, \mu)$ is a purely atomic measure space with $\mu(\{n\}) > 0$ for all $n \in \mathbb{N}$. If $T : \mathbb{N} \rightarrow \mathbb{N}$ is a non-singular measurable transformation and $u : \mathbb{N} \rightarrow [0, \infty)$ a measurable function, then we can define a composition operator C_T by

$$C_T f(n) := f(T(n)) \quad (f \in \ell^0, n \in \mathbb{N}),$$

and a multiplication operator M_u by

$$M_u f(n) := u(n) f(n) \quad (f \in \ell^0, n \in \mathbb{N}).$$

Given Orlicz functions Φ and Ψ , we can look for conditions on the triples (Φ, Ψ, T) and (Φ, Ψ, u) , so that C_T or M_u act continuously from $\ell^\Phi(\mathbb{N})$ into $\ell^\Psi(\mathbb{N})$. Note that for every function f on \mathbb{N} we have

$$\begin{aligned} I_\Psi(C_T f, \mu) &= \sum_{n=1}^\infty \Psi(f(T(n)))\mu(\{n\}) = \sum_{m \in T(\mathbb{N})} \sum_{s \in T^{-1}(m)} \Psi(f(m))\mu(\{s\}) \\ &= \sum_{m \in T(\mathbb{N})} \Psi(f(m)) \sum_{s \in T^{-1}(m)} \mu(\{s\}) = \sum_{m \in T(\mathbb{N})} \Psi(f(m))\mu \circ T^{-1}(\{m\}) \\ &= I_\Psi(f|_{T(\mathbb{N})}, \mu \circ T^{-1}) = I_\Psi^\omega(f|_{T(\mathbb{N})}, \mu), \end{aligned} \tag{6.1}$$

where $\omega(n) = \frac{\mu \circ T^{-1}(\{n\})}{\mu(\{n\})}$. This means, of course, that $C_T f \in \ell^\Psi(\mathbb{N})$ if and only if $f|_{T(\mathbb{N})} \in \ell_\omega^\Psi(T(\mathbb{N}))$.

Similarly, for every function f on \mathbb{N} we have

$$I_\Psi(M_u f, \mu) = \sum_{n=1}^\infty \Psi(u(n) f(n))\mu(\{n\}) = I_\Psi^u(f, \mu), \tag{6.2}$$

which implies that $M_u f \in \ell^\Psi(\mathbb{N})$ if and only if $f \in \ell^{\Psi_u}(\mathbb{N})$, where $\Psi_u(n, x) := \Psi(u(n)x)$.

To work with purely atomic measures, we need to use the discrete version of Ishii’s theorem due to Shragin [37] (see also [15, 17]).

Theorem 6.1 (Shragin 1976) *Let $(\mathbb{N}, 2^{\mathbb{N}}, \mu_c)$ be the counting measure space. Let $\Phi = (\Phi_n)_{n=1}^\infty$ and $\Psi = (\Psi_n)_{n=1}^\infty$ be Musielak–Orlicz functions, and let $\ell^\Phi(\mathbb{N})$ and $\ell^\Psi(\mathbb{N})$ be the Musielak–Orlicz spaces corresponding to Φ, Ψ , and the counting measure μ_c . Then $\ell^\Phi(\mathbb{N}) \subset \ell^\Psi(\mathbb{N})$ if and only if there exist $K, \delta > 0$ and a sequence of non-negative numbers $(c_n)_{n=1}^\infty \in \ell^1(\mathbb{N})$ such that*

$$\Psi_n(Kx) \leq \Phi_n(x) + c_n \tag{6.3}$$

for all $x \in \mathbb{R}$ and $n \in \mathbb{N}$ satisfying $\Phi_n(x) \leq \delta$.

We use Shragin’s result to prove the necessity part of the following theorem.

Theorem 6.2 *Let $(\mathbb{N}, 2^{\mathbb{N}}, \mu)$ be a purely atomic measure space, and let Φ and Ψ be Young functions vanishing only at zero and taking only finite values. The composition operator C_T acts continuously from $\ell^\Phi(\mathbb{N})$ into $\ell^\Psi(\mathbb{N})$ if and only if there exist $K, \delta > 0$ and a sequence of non-negative numbers $(c_n)_{n=1}^\infty \in \ell^1(T(\mathbb{N}))$ such that*

$$\Psi(Kx) \mu \circ T^{-1}(\{n\}) \leq \Phi(x)\mu(\{n\}) + c_n \tag{6.4}$$

for all $x \in \mathbb{R}$ and $n \in T(\mathbb{N})$ satisfying $\Phi(x)\mu(\{n\}) \leq \delta$.

Remark 6.3 In the preceding two theorems δ can be taken in the interval $(0, 1)$.

Proof (of Theorem 6.2) Assume that there are $K > 0, 1 > \delta > 0$, and a sequence of non-negative numbers $(c_n)_{n=1}^\infty \in \ell^1(T(\mathbb{N}))$ such that condition (6.4) is satisfied whenever $\Phi(x)\mu(\{n\}) \leq \delta$. Let f be a function in the unit sphere of $\ell^\Phi(\mathbb{N})$. Then $I_\Phi(f) \leq 1$ and further $I_\Phi(\delta f) \leq \delta I_\Phi(f) \leq \delta$, hence $\Phi(\delta f(n))\mu(\{n\}) \leq \delta$ for all $n \in \mathbb{N}$. Now setting

$$b := K \delta \quad \text{and} \quad c := \sum_{n \in T(\mathbb{N})} c_n,$$

by (6.4) we obtain

$$\begin{aligned} I_\Psi(b C_T f) &= \sum_{n=1}^\infty \Psi(b f(T(n)))\mu(\{n\}) = \sum_{n \in T(\mathbb{N})} \Psi(b f(n))\mu \circ T^{-1}(\{n\}) \\ &\leq \sum_{n \in T(\mathbb{N})} \Phi(\delta f(n))\mu(\{n\}) + \sum_{n \in T(\mathbb{N})} c_n \leq \delta \sum_{n \in T(\mathbb{N})} \Phi(f(n))\mu(\{n\}) + c \\ &\leq \delta + c \end{aligned}$$

Denoting $d = \max(\delta + c, 1)$, we get from this that

$$I_\Psi\left(\frac{b}{d} C_T f\right) \leq \frac{1}{d} I_\Psi(b C_T f) \leq 1.$$

Hence, for all functions f in the unit sphere of $\ell^\Phi(\mathbb{N})$ we have

$$\left\| \frac{b}{d} C_T f \right\|_{\ell^\Psi(\mathbb{N})} \leq 1,$$

or

$$\|C_T f\|_{\ell^\Psi(\mathbb{N})} \leq \frac{d}{b},$$

i.e. the composition operator C_T acts continuously from $\ell^\Phi(\mathbb{N})$ into $\ell^\Psi(\mathbb{N})$.

Now we shall prove that condition (6.4) follows from the continuity of the composition operator $C_T : \ell^\Phi(\mathbb{N}) \rightarrow \ell^\Psi(\mathbb{N})$. To this end, first note that, by Shragin’s result, condition (6.4) is equivalent to the inclusion $\ell^\Phi(T(\mathbb{N})) \subset \ell^\Psi_\omega(T(\mathbb{N}))$. Indeed, it is sufficient to plug $\Phi_n(x) = \Phi(x)\mu(\{n\})$ and $\Psi_n(x) = \Psi(x)\mu \circ T^{-1}(\{n\})$ in the inequality (6.3). Hence, we only have to show that the inclusion $\ell^\Phi(T(\mathbb{N})) \subset \ell^\Psi_\omega(T(\mathbb{N}))$ follows from the continuity of C_T .

Let $f \in \ell^\Phi(T(\mathbb{N}))$. Then $\tilde{f} \in \ell^\Phi(\mathbb{N})$, where \tilde{f} is the extension of f to \mathbb{N} with zero values outside of $T(\mathbb{N})$. Since C_T is continuous, we have $C_T \tilde{f} \in \ell^\Psi(\mathbb{N})$, but, as we noticed above, this means that $f \in \ell^\Psi_\omega(T(\mathbb{N}))$. □

Remark 6.4 We could have proved the sufficiency part of Theorem 6.2 via Shragin’s theorem and the fact that $(f \in \ell^\Phi(\mathbb{N}) \implies C_T f \in \ell^\Psi(\mathbb{N}))$ implies the continuity of the operator C_T since any non-negative linear operator (C_T is such an operator) between Banach lattices is continuous. Let $\ell^\Phi(T(\mathbb{N})) \subset \ell^\Psi_\omega(T(\mathbb{N}))$ [which, as we have pointed out, is equivalent to condition (6.4)] and let $f \in \ell^\Phi(\mathbb{N})$. Then $f|_{T(\mathbb{N})} \in \ell^\Phi(T(\mathbb{N}))$ and so $f|_{T(\mathbb{N})} \in \ell^\Psi_\omega(T(\mathbb{N}))$, which means that indeed $C_T f \in \ell^\Psi(\mathbb{N})$. This method of proof, however, does not provide any estimate of the norm of the operator C_T .

A similar theorem can be stated for the multiplication operator.

Theorem 6.5 *Let $(\mathbb{N}, \Sigma, \mu)$ be a purely atomic measure space, and let Φ and Ψ be Young functions vanishing only at zero and taking only finite values. The multiplication operator M_u acts continuously from $\ell^\Phi(\mathbb{N})$ into $\ell^\Psi(\mathbb{N})$ if and only if there exist $K, \delta > 0$ and a sequence of non-negative numbers $(c_n)_{n=1}^\infty \in \ell^1(\mathbb{N})$ such that*

$$\Psi(Ku(n)x) \leq \Phi(x) + c_n \tag{6.5}$$

for all $x \in \mathbb{R}$ and $n \in \mathbb{N}$ satisfying $\Phi(x)\mu(\{n\}) \leq \delta$.

Proof The proof proceeds along the same lines as the proof of Theorem 6.2, and we will omit it. □

Remark 6.6 As in the proof of Theorem 6.2, the sufficiency part can also be proved via Shragin’s theorem with the substitutions $\Phi_n(x) = \Phi(x)$ and $\Psi_n(x) = \Psi(u(n)x)$ in condition 6.3, which then becomes equivalent to the inclusion $\ell^\Phi(\mathbb{N}) \subset \ell^{\Psi_u}(\mathbb{N})$.

The same techniques can be applied in all theorems in which $h|_{T(\mathfrak{B})} = 0$ μ -a.e., where $h = d(\mu \circ T^{-1})/d\mu$, and/or $u|_{\mathfrak{B}} = 0$ μ -a.e.

In the case of a general measure space Ω , the non-atomic part of Ω and the purely atomic part of Ω can be treated separately, and this allows us to state the following.

Theorem 6.7 *Let (Ω, Σ, μ) be a measure space, and let Φ and Ψ be Young functions vanishing only at zero and assuming only finite values.*

- (a) *Assume that a non-singular measurable transformation T maps \mathfrak{B} into itself and $\Omega \setminus \mathfrak{B}$ into itself. The composition operator C_T acts continuously from $L^\Phi(\Omega)$ into $L^\Psi(\Omega)$ if and only if the following conditions are jointly satisfied:*
 - (i) *there exist a constant $K > 0$, a set $A \in \Sigma \cap T(\mathfrak{B})$ with $\mu(A) = 0$, and a function $g \in L^1_+(T(\mathfrak{B}))$, such that*

$$\Psi(Kx)h(t) \leq \Phi(x) + g(t)$$

for all $x \geq 0$ and all $t \in T(\mathfrak{B}) \setminus A$;

- (ii) *condition (6.4) (we identify \mathbb{N} with $\Omega \setminus \mathfrak{B}$).*
- (b) *The multiplication operator M_u acts continuously from $L^\Phi(\Omega)$ into $L^\Psi(\Omega)$ if and only if the following conditions are jointly satisfied:*
 - (i) *there exist a constant $K > 0$, a set $A \in \Sigma \cap \mathfrak{B}$ with $\mu(A) = 0$, and a function $g \in L^1_+(\mathfrak{B})$, such that*

$$\Psi(u(t)x) \leq \Phi(x) + g(t)$$

for all $x \geq 0$ and all $t \in \mathfrak{B} \setminus A$;

- (ii) *condition (6.5) (again, we identify \mathbb{N} with $\Omega \setminus \mathfrak{B}$).*

Proof We give the proof for both operators concurrently. We can represent $L^\Phi(\Omega)$ and $L^\Psi(\Omega)$ as the direct sums of the spaces over the non-atomic part of Ω and the purely atomic part of Ω , i.e. $L^\Phi(\Omega) = L^\Phi(\mathfrak{B}) \oplus L^\Phi(\Omega \setminus \mathfrak{B})$ and $L^\Psi(\Omega) = L^\Psi(\mathfrak{B}) \oplus L^\Psi(\Omega \setminus \mathfrak{B})$. Therefore, every function $f \in L^\Phi(\Omega)$ can be uniquely written as $f = f\chi_{\mathfrak{B}} + f\chi_{\Omega \setminus \mathfrak{B}}$, and we have

$$\begin{aligned} C_T f(t) &= C_T(f\chi_{\mathfrak{B}} + f\chi_{\Omega \setminus \mathfrak{B}})(t) \\ &= (f\chi_{\mathfrak{B}} + f\chi_{\Omega \setminus \mathfrak{B}})(T(t)) \\ &= f(T(t))\chi_{T^{-1}(\mathfrak{B})}(t) + f(T(t))\chi_{T^{-1}(\Omega \setminus \mathfrak{B})}(t) \\ &= f(T(t))\chi_{\mathfrak{B}}(t) + f(\chi(t))\chi_{\Omega \setminus \mathfrak{B}}(t) \end{aligned}$$

and

$$\begin{aligned} M_u f(t) &= M_u(f\chi_{\mathfrak{B}} + f\chi_{\Omega \setminus \mathfrak{B}})(t) = u(f\chi_{\mathfrak{B}} + f\chi_{\Omega \setminus \mathfrak{B}})(t) \\ &= u(t)f(t)\chi_{\mathfrak{B}}(t) + u(t)f(t)\chi_{\Omega \setminus \mathfrak{B}}(t). \end{aligned}$$

Hence, if we define the operators $C_{T_{\mathfrak{B}}}$, $M_{u_{\mathfrak{B}}}$ on $L^\Phi(\mathfrak{B})$ and the operators $C_{T_{\Omega \setminus \mathfrak{B}}}$, $M_{u_{\Omega \setminus \mathfrak{B}}}$ on $L^\Phi(\Omega \setminus \mathfrak{B})$ by the obvious formulas, we are justified to write

$$C_T = C_{T_{\mathfrak{B}}} \oplus C_{T_{\Omega \setminus \mathfrak{B}}} \quad \text{and} \quad M_u = M_{u_{\mathfrak{B}}} \oplus M_{u_{\Omega \setminus \mathfrak{B}}}.$$

Of course, we have the following inequalities for the norms

$$\begin{aligned} \max\{\|C_{T_{\mathfrak{B}}}\|, \|C_{T_{\Omega \setminus \mathfrak{B}}}\|\} &\leq \|C_T\| \leq \|C_{T_{\mathfrak{B}}}\| + \|C_{T_{\Omega \setminus \mathfrak{B}}}\| \\ \max\{\|M_{u_{\mathfrak{B}}}\|, \|M_{u_{\Omega \setminus \mathfrak{B}}}\|\} &\leq \|M_u\| \leq \|M_{u_{\mathfrak{B}}}\| + \|M_{u_{\Omega \setminus \mathfrak{B}}}\|. \end{aligned}$$

Hence it follows that the operator C_T acts continuously from $L^\Phi(\Omega)$ into $L^\Psi(\Omega)$ if and only if the operator $C_{T_{\mathfrak{B}}}$ acts continuously from $L^\Phi(\mathfrak{B})$ into $L^\Psi(\mathfrak{B})$ and the operator $C_{T_{\Omega \setminus \mathfrak{B}}}$ acts continuously from $L^\Phi(\Omega \setminus \mathfrak{B})$ into $L^\Psi(\Omega \setminus \mathfrak{B})$. The same applies to the multiplication operator: M_u acts continuously from $L^\Phi(\Omega)$ into $L^\Psi(\Omega)$ if and only if $M_{u_{\mathfrak{B}}}$ acts continuously from $L^\Phi(\mathfrak{B})$ into $L^\Psi(\mathfrak{B})$ and $M_{u_{\Omega \setminus \mathfrak{B}}}$ acts continuously from $L^\Phi(\Omega \setminus \mathfrak{B})$ into $L^\Psi(\Omega \setminus \mathfrak{B})$. Therefore, the condition for boundedness of the operator C_T is exactly the conjunction of the conditions for the boundedness of the operators $C_{T_{\mathfrak{B}}}$ and $C_{T_{\Omega \setminus \mathfrak{B}}}$. Analogously, the condition for boundedness of the operator M_u is the conjunction of conditions for the boundedness of the operators $M_{u_{\mathfrak{B}}}$ and $M_{u_{\Omega \setminus \mathfrak{B}}}$. Recall that in [3] it was shown that (a)(i) is a sufficient condition for the continuity of C_T and that (b)(i) is a necessary and sufficient condition for the continuity of M_u in the case of a non-atomic measure space. By analogous arguments to the ones we have used in the proof of the necessity part of Theorem 6.2 (applying Ishii's theorem rather than Shragin's) it can be shown that (a)(i) is also a necessary condition for the continuity of C_T in the case of a non-atomic measure space, which generalizes [3], where (a)(i) was shown to be a necessary condition for the continuity of C_T in the non-atomic case under the additional assumption that the mapping T is surjective up to sets of measure zero. Moreover, Theorems 6.2 and 6.5 provide parallel sufficient and necessary conditions (a)(ii) and (b)(ii) in the case of a purely atomic measure space. This completes the proof of Theorem 6.7 \square

Acknowledgements The authors wish to thank the anonymous referee for providing important corrections and insightful suggestions that helped to improve the text significantly.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. Ando, T.: On products of Orlicz spaces. *Math. Ann.* **140**, 174–186 (1960)
2. Banach, S.: *Théorie des opérations linéaires*. Monografie Matematyczne, Warsaw (1932)

3. Chawziuk, T., Cui, Y., Estaremi, Y., Hudzik, H., Kaczmarek, R.: Composition and multiplication operators between Orlicz function spaces. *J. Inequal. Appl.* **2016**, 52 (2016). doi:[10.1186/s13660-016-0972-9](https://doi.org/10.1186/s13660-016-0972-9)
4. Conway, J.B.: *A course in functional analysis*. Springer, New York (1990)
5. Cowen, C.C., MacCluer, B.D.: *Composition Operators on Spaces of Analytic Functions*. Studies in Advanced Mathematics. CRC Press, Boca Raton (1995)
6. Cui, Y., Hudzik, H., Kumar, R., Maligranda, L.: Composition operators in Orlicz spaces. *J. Aust. Math. Soc.* **76**, 189–206 (2004)
7. Cui, Y., Hudzik, H., Kumar, R., Kumar, R.: Compactness and essential norms of composition operators on Orlicz-Lorentz spaces. *Appl. Math. Lett.* **25**(11), 1778–1783 (2012)
8. Cui, Y., Hudzik, H., Kumar, R., Kumar, R.: Erratum and addendum to the paper Compactness and essential norms of composition operators on Orlicz-Lorentz spaces [*Applied Mathematics Letters* 25 (2012) 1778–1783]. *Comment. Math.* **54**(2), 259–260 (2014)
9. del Campo, R., Fernandez, A., Ferrando, I., Mayoral, F., Naranjo, F.: Multiplication operators on spaces on integrable functions with respect to a vector measure. *J. Math. Anal. Appl.* **343**, 514–524 (2008)
10. Domański, P., Langenbruch, M.: Composition operators on spaces of real analytic functions. *Math. Nachr.* **254**(255), 68–86 (2003)
11. Eryilmaz, I.: Weighted composition operators on weighted Lorentz spaces. *Colloq. Math.* **128**, 143–151 (2012)
12. Estaremi, Y.: Multiplication and composition operators between two diferent Orlicz spaces. [arXiv:1301.4830v2](https://arxiv.org/abs/1301.4830v2). 20 Oct 2013
13. Gupta, S., Komal, B.S., Suri, N.: Weighted composition operators on Orlicz spaces. *Int. J. Contemp. Math. Sci.* **1**, 11–20 (2010)
14. Hencl, S., Kleprlík, L., Malý, J.: Composition operator and Sobolev-Lorentz spaces $WL^{n,q}$. *Studia Math.* **221**, 197–208 (2014)
15. Hudzik, H., Kamińska, A.: On uniformly convexifiable and B-convex Musielak-Orlicz spaces. *Comment. Math. Prace Mat.* **25**(1), 59–75 (1985)
16. Hudzik, H., Krbec, M.: On non-effective weights in Orlicz spaces. *Indag. Mathem. (N.S.)* **18**(2), 215–231 (2007)
17. Ishii, J.: On equivalence of modular function spaces. *Proc. Jpn. Acad.* **35**, 551–556 (1959)
18. Jabbarzadeh, M.R.: Weighted composition operators between L^p -spaces. *Bull. Korean Math. Soc.* **42**, 369–378 (2005)
19. Kolwicz, P., Leśniak, K., Maligranda, L.: Pointwise multipliers of Calderón-Lozanovskii spaces. *Math. Nachr.* **286**, 876–907 (2013)
20. Kolwicz, P., Leśniak, K., Maligranda, L.: Pointwise product of some Banach function spaces and factorization. *J. Funct. Anal.* **266**, 616–659 (2014)
21. Komal, B.S., Gupta, S.: Multiplication operators between Orlicz spaces. *Integr. Equ. Oper. Theory* **41**, 324–330 (2001)
22. Krasnoselskii, M.A., Rutickii, B.: *Convex functions and Orlicz spaces*. Noordhoff, Netherlands (1961) [Translated from Russian]
23. Kumar, R.: Weighted composition operators between two L^p -spaces. *Mat. Vesnik* **61**, 111–118 (2009)
24. Kumar, R.: Composition operators on Orlicz spaces. *Integr. Equ. Oper. Theory* **29**, 17–22 (1997)
25. Kumar, R., Kumar, R.: Composition operators on Orlicz-Lorentz spaces. *Integr. Equ. Oper. Theory* **60**, 79–88 (2008)
26. Maligranda, L.: *Orlicz spaces and interpolation*, Seminars in Math. 5, Univ. Estadual de Campinas, Campinas SP, Brazil (1989)
27. Maligranda, L., Persson, L.E.: Generalized duality of some Banach function spaces. *Indag. Math.* **92**(3), 323–338 (1989)
28. Musielak, J.: *Orlicz spaces and modular spaces*. Lecture Notes in Math, vol. 1034. Springer, Berlin (1983)
29. Nielsen, O.A.: *An introduction to integration and measure theory*. Wiley-Interscience, New York (1997)
30. Nowak, M.: On inclusions between Orlicz spaces. *Comment. Math. Prace Mat.* **26**, 265–277 (1986)
31. Nowak, M.: Some equalities among Orlicz spaces. I. *Comment. Math. Prace Mat.* **29**, 255–275 (1990)
32. O’Neil, R.: Fractional integration in Orlicz spaces. I. *Trans. Am. Math. Soc.* **115**, 300–328 (1965)
33. Rao, M.M.: Convolutions of vector fields II: random walk models. *Nonlinear Anal.* **47**, 3599–3615 (2001)
34. Rao, M.M., Ren, Z.D.: *Theory of Orlicz spaces*. Marcel Dekker, New York (1991)

35. Shapiro, J.H.: *Composition Operators and Classical Function Theory*. Universitext Tracts in Mathematics. Springer, New York (1993)
36. Singh, R.K., Manhas, J.S.: *Composition operators on function spaces*, North-Holland Mathematics Studies, 179. North-Holland Publishing Co., Amsterdam (1993)
37. Shragin, I.V.: The conditions on embeddings of some classes and conclusions from them. *Mat. Zam.* **20**, 681–692 (1976). (in Russian)
38. Takagi, H., Yokouchi, K.: Multiplication and composition operators between two L^p -spaces. *Contemp. Math.* **232**, 321–338 (1999)