PERSPECTIVE

Redemystifying MST1/hippo signaling

Lei Xiao, Zengqiang Yuan✉

State Key Laboratory of Brain and Cognitive Science, Institute of Biophysics, Chinese Academy of Sciences, Beijing 100101, China

✉ Correspondence: zqyuan@ibp.ac.cn

Since the Mammalian Ste20-like kinases (MST) 1/2 were first identified in 1995, our knowledge about MST1/2 and its *Drosophila* ortholog Hippo has expanded to diverse biologic process ranging from cell survival and death, organ size control, to proliferation and tumorigenesis.

MST1 (also known as *Stk4* and *Krs2*) and MST2 (also known as *Stk3* and *Krs1*) were first identified as homologs of the ste20 kinase from *Saccharomyces cerevisiae* (Creasy and Chernoff, 1995). Subsequently, these proteins were also identified in 'in gel' kinase assays as kinases that respond to extreme cellular stress (Taylor et al., 1996). Though a number of apoptotic or stress stimuli have been reported to activate MST, proteolytic cleavage of MST by caspase 3 remains the only and best understood mechanism that regulate MST kinase activity to date (Kakeya et al., 1998). Upon proapoptotic stimuli, the N-terminal proteolytic fragment of MST translocates into the nucleus and phosphorylates histone H2B, which leads to the chromatin condensation and mammalian cell death (Cheung et al., 2003). Interestingly, histone H2B phosphorylation also occurs in *S. cerevisiae*, through the phosphorylation on Ser10, a residue distinct from mammalian histone H2B Ser14 (Ahn et al., 2005). However, growing evidence suggests that full length MST also promotes cell death independently of proteolysis or nuclear translocation (Ahn et al., 2005; Lehtinen et al., 2006). In primary mammalian neurons, oxidative stress-activated-MST phosphorylates the transcription factor FOXO, which could translocate into the nucleus and upregulate the transcriptional activity of pro-death genes, including BIM. Moreover, the characterization of the *C. elegans* ortholog *CST-1* broadens MST functions beyond the control of cell death to the regulation of life span in nematodes (Lehtinen et al., 2006). Despite the flurry of interest in the MST, both the upstream regulators and downstream targets of MST in different scenarios are still in a puzzle.

In contrast to the limited progress of study in mammals, genetic screens for flies with defects in organ size lead to the identification of Hippo complex and the upstream regulators

that modulate cell growth and survival. As a tumor suppressor, The Hippo/MST kinase cooperates with adaptor protein Salvador (Sav)/WW45, phosphorylates and activates the downstream kinase Warts/Lats, which in turn leads to the phosphorylation and inhibition of the transcriptional activator Yorkie (Yki)/YAP through promoting its nuclear exportation in 14-3-3 dependent and-independent manners. Two membrane-associated FERM domain proteins, Merlin (Mer) and Expanded (Ex), have been suggested to function in parallel but differentially regulate the Hippo pathway since Ex predominantly regulates proliferation whereas Mer regulates apoptosis (Pellock et al., 2007). However, McNeill and colleagues showed a physical interaction between Ex and Yki that may directly inhibit Yki activity independent of Hippo (Badouel et al., 2009), implying that there is more complicated cross-regulation in the Hippo/Yki signaling. Recently, the atypical cadherin Fat (Ft) and its ligand Dachshous (Ds) appear to signal through Ex to activate the Hippo pathway (Bennett and Harvey, 2006). Crumbs (Crb) were identified as a novel Hippo pathway regulator via modulating Ex levels and localization, indicating that Hippo/MST might be involved in the establishment of the cellular polarity (Grzeschik et al., 2010; Robinson et al., 2010). Richardson and colleagues also proposed that cell polarity regulators giant-larvae (Lg) I and atypical protein kinase C (aPKC) feed into Hippo pathway. Lgl acts antagonistically to aPKC to regulate Hippo and Ras association family member (RASSF) localization (Grzeschik et al., 2010). Most recently, genetic analysis revealed that WW domain containing protein Kibra physically interacts with Mer and acts upstream of Mer to regulate Hpo activity (Baumgartner et al., 2010). The core components of the Hippo pathway are well conserved, and the understanding of Hippo functions in flies may shed light on the study of MST function in mammals.

To investigate the physiological functions of MST family proteins *in vivo*, MST1/2 knockout mice were generated (Oh et al., 2009; Zhou et al., 2009). MST1 deletion resulted in decreased numbers of peripheral T cells, mainly naive T cells,

which still proliferated after T cell receptor (TCR) ligation *in vitro* and impaired lymphocyte homing to the spleen and lymph nodes (Zhou et al., 2008; Katagiri et al., 2009). In addition, defective egress of mature thymocytes from *Mst1* deficient thymus was also reported (Dong et al., 2009). However, *MST2* knockout mice exhibited no developmental or immunological defects. Due to the redundant functions of *MST1/2*, *MST1/2* single knockout model provided limited information. *MST1/2* double knockout resulted in embryonic lethality, indicating that *MST* is essential for the early development in mice. Single copy of either *MST1* or *MST2* underwent normal organ development. Moreover, by 15 months of age, *MST1*^{-/-} *MST2*^{+/-} mice developed hepatocellular carcinoma (HCC) because of *MST2* loss of heterozygosity. Correspondingly, tissue-specific ablation of both *MST1* and *MST2* in liver leads to HCC (Zhou et al., 2009). Yang and colleagues also reported that tumors developed in the liver with a substantial increase of the stem/progenitor cells by 6 months after removing *MST1* and *MST2* postnatally (Song et al., 2010). Interestingly, apoptosis induced by Fas ligand or TNF α was blocked in the *MST1/2* deficient cells *in vivo* (Zhou et al., 2009; Song et al., 2010). These *in vivo* experiments further supported the physiological significance of *MST1/2* in apoptosis and tumorigenesis.

It is worth noting that *in vivo* experiment also reveals that *MSTs* are differentially regulated in various cell types. For example, The cleaved form of *MST1* are the major form in livers but absent in spleen or MEF cells, indicating that tissue-specific cleavage may be an important mechanism of *MST1/2* regulation *in vivo* (Zhou et al., 2009). Obviously, the upstream regulatory mechanism identified from flies is insufficient for us to understand the variation, especially in mammals. Fortunately, in mammals, we and others have found that *Akt* directly phosphorylates *MST1*, which leads to inhibition of *MST1* cleavage and kinase activity (Cinar et al., 2007; Yuan et al., 2010). Moreover, this phosphorylation inversely correlates with *MST1* auto-phosphorylation/activation and associates with adverse prognosis in human ovarian cancer. Recent study adds a new player *PHLPP* in this model. As a phosphatase, *PHLPP* directly dephosphorylates *MST1* on Thr387, the same site as phosphorylated by *Akt*, resulting in the activation of *MST1* (Qiao et al., 2010). Though this delicate autoinhibitory triangle cannot explain the cell type-dependent functions of *MST1/2*, we can conclude that phosphorylation and cleavage of *MSTs*, alone or cooperated, play an important role in regulating *MST* activity in a variety of settings.

According to many elegant experiments, we have learned a lot about *MST*. However, more questions have been raised than been answered. First, as a tumor suppressor, is mammalian *MST* dysregulated at pathological state indeed and how? While waiting for the next genetic screen in flies, it is crucial to validate whether *Mer*, *Ex*, *Crb* or those cell polarity regulators really work in mammals. Second, in spite of the

redundant roles of *MST1/2*, do they have distinguished functions in different situations? Why does this happen? Does *MST1* or *MST2* fall into unique signaling pathway even though they share most of the signaling components? Finally, at present, oxidative stress remains one of the best known stimuli that activate *MST* kinases, which leading to the phosphorylation of *FOXO* or histone H2B in different type of cells. How does *MST* select the substrates? Some evidences from us and others suggest that the c-terminal domain of *MST1* is dynamically regulated and can exhibit stimulatory or inhibitory activities with specific substrates (Anand et al., 2008). Therefore, more studies related to the substrate specification through the c-terminal modification and regulation will help to solve this puzzle. Nonetheless, an important goal of future studies is to determine the upstream regulators of *MST*, possibly including kinases, phosphatases or cytoskeleton proteins, in the regulation of the *MST* function in cell proliferation and apoptosis, organ size control as well as cellular polarity establishment.

ACKNOWLEDGEMENTS

This study was supported in part by National Natural Science Foundation of China (Grant No. 30870792), and the National Basic Research Program of China (973 Program) (Grant No. 2009CB918704).

REFERENCES

- Ahn, S.H., Cheung, W.L., Hsu, J.Y., Diaz, R.L., Smith, M.M., and Allis, C.D. (2005). Sterile 20 kinase phosphorylates histone H2B at serine 10 during hydrogen peroxide-induced apoptosis in *S. cerevisiae*. *Cell* 120, 25–36.
- Anand, R., Kim, A.Y., Brent, M., and Marmorstein, R. (2008). Biochemical analysis of *MST1* kinase: elucidation of a C-terminal regulatory region. *Biochemistry* 47, 6719–6726.
- Badouel, C., Gardano, L., Amin, N., Garg, A., Rosenfeld, R., Le Bihan, T., and McNeill, H. (2009). The FERM-domain protein Expanded regulates Hippo pathway activity via direct interactions with the transcriptional activator Yorkie. *Dev Cell* 16, 411–420.
- Baumgartner, R., Poembacher, I., Buser, N., Hafen, E., and Stocker, H. (2010). The WW domain protein Kibra acts upstream of Hippo in *Drosophila*. *Dev Cell* 18, 309–316.
- Bennett, F.C., and Harvey, K.F. (2006). Fat cadherin modulates organ size in *Drosophila* via the Salvador/Warts/Hippo signaling pathway. *Curr Biol* 16, 2101–2110.
- Cheung, W.L., Ajiro, K., Samejima, K., Kloc, M., Cheung, P., Mizzen, C.A., Beeser, A., Etkin, L.D., Chernoff, J., Earnshaw, W.C., et al. (2003). Apoptotic phosphorylation of histone H2B is mediated by mammalian sterile twenty kinase. *Cell* 113, 507–517.
- Cinar, B., Fang, P.K., Lutchman, M., Di Vizio, D., Adam, R.M., Pavlova, N., Rubin, M.A., Yelick, P.C., and Freeman, M.R. (2007). The pro-apoptotic kinase *Mst1* and its caspase cleavage products are direct inhibitors of *Akt1*. *EMBO J* 26, 4523–4534.
- Creasy, C.L., and Chernoff, J. (1995). Cloning and characterization of a member of the *MST* subfamily of Ste20-like kinases. *Gene* 167, 303–306.

- Dong, Y., Du, X., Ye, J., Han, M., Xu, T., Zhuang, Y., and Tao, W. (2009). A cell-intrinsic role for Mst1 in regulating thymocyte egress. *J Immunol* 183, 3865–3872.
- Grzeschik, N.A., Parsons, L.M., Allott, M.L., Harvey, K.F., and Richardson, H.E. (2010). Lgl, aPKC, and Crumbs regulate the Salvador/Warts/Hippo pathway through two distinct mechanisms. *Curr Biol* 20, 573–581.
- Takeya, H., Onose, R., and Osada, H. (1998). Caspase-mediated activation of a 36-kDa myelin basic protein kinase during anticancer drug-induced apoptosis. *Cancer Res* 58, 4888–4894.
- Katagiri, K., Katakai, T., Ebisuno, Y., Ueda, Y., Okada, T., and Kinashi, T. (2009). Mst1 controls lymphocyte trafficking and interstitial motility within lymph nodes. *EMBO J* 28, 1319–1331.
- Lehtinen, M.K., Yuan, Z., Boag, P.R., Yang, Y., Villén, J., Becker, E.B., DiBacco, S., de la Iglesia, N., Gygi, S., Blackwell, T.K., *et al.* (2006). A conserved MST-FOXO signaling pathway mediates oxidative-stress responses and extends life span. *Cell* 125, 987–1001.
- Oh, S., Lee, D., Kim, T., Kim, T.S., Oh, H.J., Hwang, C.Y., Kong, Y.Y., Kwon, K.S., and Lim, D.S. (2009). Crucial role for Mst1 and Mst2 kinases in early embryonic development of the mouse. *Mol Cell Biol* 29, 6309–6320.
- Pellock, B.J., Buff, E., White, K., and Hariharan, I.K. (2007). The *Drosophila* tumor suppressors Expanded and Merlin differentially regulate cell cycle exit, apoptosis, and Wingless signaling. *Dev Biol* 304, 102–115.
- Qiao, M., Wang, Y., Xu, X., Lu, J., Dong, Y., Tao, W., Stein, J., Stein, G. S., Iglehart, J.D., Shi, Q., *et al.* (2010). Mst1 is an interacting protein that mediates PHLPPs' induced apoptosis. *Mol Cell* 38, 512–523.
- Robinson, B.S., Huang, J., Hong, Y., and Moberg, K.H. (2010). Crumbs regulates Salvador/Warts/Hippo signaling in *Drosophila* via the FERM-domain protein expanded. *Curr Biol* 20, 582–590.
- Song, H., Mak, K.K., Topol, L., Yun, K., Hu, J., Garrett, L., Chen, Y., Park, O., Chang, J., Simpson, R.M., *et al.* (2010). Mammalian Mst1 and Mst2 kinases play essential roles in organ size control and tumor suppression. *Proc Natl Acad Sci U S A* 107, 1431–1436.
- Taylor, L.K., Wang, H.C., and Erikson, R.L. (1996). Newly identified stress-responsive protein kinases, Krs-1 and Krs-2. *Proc Natl Acad Sci U S A* 93, 10099–10104.
- Yuan, Z., Kim, D., Shu, S., Wu, J., Guo, J., Xiao, L., Kaneko, S., Coppola, D., and Cheng, J.Q. (2010). Phosphoinositide 3-kinase/Akt inhibits MST1-mediated pro-apoptotic signaling through phosphorylation of threonine 120. *J Biol Chem* 285, 3815–3824.
- Zhou, D., Medoff, B.D., Chen, L., Li, L., Zhang, X.F., Praskova, M., Liu, M., Landry, A., Blumberg, R.S., Boussiotis, V.A., *et al.* (2008). The Nore1B/Mst1 complex restrains antigen receptor-induced proliferation of naïve T cells. *Proc Natl Acad Sci U S A* 105, 20321–20326.
- Zhou, D., Conrad, C., Xia, F., Park, J.S., Payer, B., Yin, Y., Lauwers, G.Y., Thasler, W., Lee, J.T., Avruch, J., *et al.* (2009). Mst1 and Mst2 maintain hepatocyte quiescence and suppress hepatocellular carcinoma development through inactivation of the Yap1 oncogene. *Cancer Cell* 16, 425–438.