



CORRECTION

## Correction to: Study Protocol for the Effects of Formula Diet with Dapagliflozin on Metabolic Improvement and Body Composition in Type 2 Diabetes Mellitus

Yasuhiro Watanabe · Nobuichi Kuribayashi · Daigaku Uchida ·  
Daisuke Suzuki · Mitsutoshi Kato · Daiji Nagayama · Hiroshi Ohashi ·  
Masahiro Ohira · Atsuhito Saiki · Ichiro Tatsuno

Published online: February 12, 2020  
© The Author(s) 2020

Correction to: Diabetes Ther (2019) 10:311–321  
<https://doi.org/10.1007/s13300-018-0555-5>

In the original article, the 6th sentence of the section “FD and calorie intake” is incorrect. The correct sentence is “The protein component of FD consists of whey protein, casein, and soy protein”.

There is a correction in the “Inclusion criteria” section, the correct sentence is Estimated glomerular filtration rate (eGFR)  $\geq 45$  ml/min/1.73 m<sup>2</sup>.

---

The original article can be found online at <https://doi.org/10.1007/s13300-018-0555-5>.

---

Y. Watanabe · M. Ohira · A. Saiki · I. Tatsuno (✉)  
Center of Diabetes, Endocrinology and Metabolism,  
Toho University Sakura Medical Center, Chiba,  
Japan  
e-mail: [ichiro.tatsuno@med.toho-u.ac.jp](mailto:ichiro.tatsuno@med.toho-u.ac.jp)

N. Kuribayashi  
Misaki Naika Clinic, Chiba, Japan

D. Uchida  
Hotaruno Central Naika, Chiba, Japan

D. Suzuki  
Suzuki Diabetes Clinic, Kanagawa, Japan

M. Kato  
Kato Clinic of Internal Medicine, Tokyo, Japan

D. Nagayama  
Nagayama Clinic, Tochigi, Japan

H. Ohashi  
Oyama East Clinic, Tochigi, Japan

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits any non-commercial use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.