should be moved to fresh air. Induction of emesis should not be performed. Gastric lavage is not helpful as camphor is rapidly absorbed. Activated charcoal may be considered if there are other ingredients that are effectively absorbed. Short acting anticonvulsants like lorazepam should be used to treat seizures. Barbiturates should be used if benzodiazepines are unsuccessful. Haemodialysis is not of much benefit.

Acknowledgment: Dr Sujata K, Associate Professor in the department.

KSSAHANA AND D RAJIV
*Department of Paediatrics,
 Saphthagiri Institute of Medical Sciences,
 Bangalore,
 Karnataka.
 dr_sahana2003@yahoo.co.in*

Widening the Spectrum of Services Delivered by the NRC

The recent Perspective by Prasad, *et al.* [1] was indeed an interesting and thought provoking article highlighting the issue of managing/preventing and treating malnutrition other than severe acute malnutrition, to be benefitted by the Nutrition rehabilitation centers. It highlights clearly the defined roles of ASHA, AWW and PHCs in managing the larger goal of malnutrition prevention/treatment.

I believe that although secondary/tertiary care hospitals may be targeting management of undernourished children other than severe acute malnutrition optimally, this may not be so at other places where NRC/IYCF are functioning, possibly in the outreach. Just as we have defined guidelines for management of varied multifactorial illnesses as ARI or Asthma, it would indeed be need of the hour to integrate updated comprehensive guidelines/protocols for the management of stunting, moderate malnutrition, growth faltering and acute weight loss needing urgent attention, also by special nutritional counseling; imparting specific knowledge about locally formulated energy dense, home based foods; identifying who and when to investigate; along with specific follow-up and monitoring at prescribed intervals.

REFERENCES

1. Matteucci MJ. One pill can kill: assessing the potential for fatal poisonings in children. *Pediatr Ann.* 2005;34:964-8.
2. Manoguerra AS, Erdman AR, Wax PM, Nelson LS, Caravati M, Cobaugh DJ, *et al.* Camphor poisoning: an evidence based practice guidelines for out of hospital management. *Clin Toxicol.* 2006;44:357-70.
3. Khine H, Weiss D, Graber N, Hoffman RS, Esteban Cruciani N, Avner JR. A cluster of children with seizures caused by camphor poisoning. *Pediatrics.* 2009;123:1269-72.
4. Gilbert J, Flamant C, Hallalel F, Doummar D, Frata A, Renolleau S. Antiflatulence treatment and status epilepticus- a case report of camphor intoxication. *Emerg Med J.* 2007;24:859-60.
5. Aggarwal A, Malhotra HS. Camphor ingestion: an unusual cause of seizures. *J Assoc Physicians India.* 2008;56:123-4.

While the NRC guidelines do observe and note the presence of undernutrition, the treatment focus is on the treatment of SAM (which undoubtedly requires urgent attention). The IYCF services wonderfully focus on the issues of exclusive breastfeeding, timely introduction of complementary foods while handling feeding problems, and complementing mother's knowledge about the right quantity and quality of foods. Though it would be a challenge to address the complex spectrum of various types and degrees of undernutrition, I would strongly believe that such an addition/improvisation in the protocols possibly in the already established NRC may augment the services delivered and have a huge impact on the prevalence of large spectrum of undernutrition. Widening of spectrum of services delivered by the NRC may have logistic implications, but it would be worthwhile for a holistic approach targeting undernutrition.

RUCHIE GULATI
*Specialist Paediatrician,
 Sardar Vallabh Bhai Patel Hospital, Delhi, India.
 ruchiegulati@gmail.com*

REFERENCES

1. Prasad V, Sinha D, Sridhar S. Falling between two stools: Operational inconsistencies between ICDS and NRHM in the management of severe malnutrition. *Indian Pediatr.* 2012;3:181-5.
2. Operational Guidelines on Facility Based Management of Children with Severe Acute Malnutrition. New Delhi: Ministry of Health and Family Welfare; 2011.