Myotonia in a Child with Muscle Hypertrophy

A 5-year-old child was brought with complaints of loss of appetite and not gaining weight. There was no history of contact with tuberculosis and other systemic illness. Diet history revealed inadequate calorie intake.

On examination, child was alert, active, oriented with normal speech output. His weight (13 kg) was <3rd centile and height (110cm) was at 50th centile. His mid arm circumference was 14.5 cm. Detailed examination of all muscle groups revealed hypertrophy of arm, thigh and leg muscles, with normal muscle power and tone. Upper limb and lower limb reflexes were brisk. Plantar response was flexor. On further examination, percussion myotonia and grip release myotonia of upper limb was present (**Web Video 1** and **Fig. 1**). We made a provisional diagnosis of myotonia congenital. He was treated with phenytoin for myotonia, and dietary advice for adequate weight gain.

Myotonia is defined as a disturbance in muscle relaxation after voluntary contraction. Common diseases of children associated with myotonia include Myotonia congenita, Myotonia fluctuans and Neuromyotonia. In these disorders, weakness is not prominent but stiffness may impair muscle function. Abnormalities in the potassium and chloride channels underlie most cases.

FIG. 1 Elicitation of myotonia in a child. (See website for video)

Myotonia associated with muscle hypertrophy is characteristic of Myotonia congenital (Becker's type). EMG establishes the diagnosis. Repetitive discharges at rates of 20 to 80 cycles per second are recorded when needle is inserted into the muscle. The waxing and waning of the amplitude and frequency of potentials produce a characteristic sound. Mexilitine and Phenytoin is used for reducing muscle stiffness in myotonia. This case scenario explains the need of eliciting myotonia in children with muscle hypertrophy to differentiate from muscular dystrophies, as children with myotonia with muscle hypertrophy are easily treatable.

VASANTHAN TANIGASALAM

Department of Pediatrics,
JIPMER, Puducherry 605 006, Tamil Nadu.
vasanthan2k6jipmer@gmail.com

NOTICE

Call for Submission of 'Clinical Videos'

Under this section, *Indian Pediatrics* publishes videos depicting an intricate technique or an interesting clinical manifestation, which are difficult to describe clearly in text or by pictures. A video file submitted for consideration for publication should be of high resolution and should be edited by the author in final publishable format. MPEG or MP4 formats are acceptable. The maximum size of file should be 20 MB. The file should not have been published elsewhere, and will be a copyright of *Indian Pediatrics*, if published. For this section, there should be a write-up of up to 250 words discussing the condition and its differential diagnoses. The write up should also be accompanied by a thumbnail image for publication in the print version and PDF. Submit videos as separate Supplementary files with your main manuscript. A maximum of three authors (not more than two from a single department) are permissible for this section. In case the video shows a patient, he/she should not be identifiable. In case the identification is unavoidable, or even otherwise, each video must be accompanied by written permission of parent/guardian, as applicable. Authors are responsible for obtaining participant consent-to-disclose forms for any videos of identifiable participants, and should edit out any names mentioned in the recording. The consent form should indicate its purpose (publication in the journal in print and online, with the understanding that it will have public access) and the signed consent of the parent/legal guardian. The copy of the consent form must be sent as supplementary file along with the write-up, and original form should be retained by the author. A sample consent form is available at our website www.indianpediatrics.net.