Erratum to: Neonicotinoid insecticides translocated in guttated droplets of seed-treated maize and wheat: a threat to honey bees?

Jana E. REETZ¹, Sebastian ZÜHLKE², Michael SPITELLER², Klaus WALLNER¹

¹Apicultural State Institute, University of Hohenheim, August-von-Hartmann-Str. 13, 70593, Stuttgart, Germany

²Institute of Environmental Research (INFU), Technische Universität Dortmund, Otto-Hahn-Str. 6, 44227, Dortmund, Germany

Erratum to: Apidologie

DOI 10.1007/s13592-011-0049-1

2. MATERIALS AND METHODS

2.2 Plant material

2.2.1 *Triticale experiments*

The submitted specifications of the insecticide Gasur and the fungicide Efa of the treated seed in triticale have to be reduced by the factor 100 (which corresponds to the recommended application rate):

Seeds of triticale were treated with a combination of the insecticide Gasur® (per

seed: 0.0175 mg imidacloprid, Bayer CropScience AG, Leverkusen, Germany) and the fungicide Efa® (per seed: 0.0006 mg triazoxide, 0.000225 mg tebuconazole, 0.00225 mg fluoxastrobin, 0.0015 mg prothioconazole; Bayer CropScience AG, Leverkusen, Germany).

This error has no influence on results and discussion, respectively, because the results (i.e. concentration of pesticides in guttation drops) are not correlated with the application rate.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s13592-011-0049-1>.

Corresponding author:

J.E. Reetz, reetz@uni-hohenheim.de