

Landscape effects on wild *Bombus terrestris* (Hymenoptera: Apidae) queens visiting highbush blueberry fields in south-central Chile

Lorena VIELI^{1,2}, Frank W. DAVIS², Bruce E. KENDALL², Miguel ALTIERI³

¹Instituto de Medio Ambiente, Universidad de la Frontera, Av. Francisco Salazar, 01145, Temuco, Chile

²Bren School of Environmental Science and Management, University of California, Santa Barbara, CA 93110-5131, USA

³Department of Environmental Science Policy and Management, University of California, Berkeley, CA 94720, USA

Received 11 March 2015 – Revised 15 October 2015 – Accepted 4 December 2015

Abstract – In this study pollinators visiting highbush blueberry fields set in landscapes with differing land use pattern in south-central Chile were investigated. Effects of spatial buffers from 0.5 to 8 km around each blueberry field on the abundance of the main wild pollinator, *Bombus terrestris* queens, were tested. Wild *B. terrestris* abundances were positively affected by natural forest area and negatively affected by high-food resource area, and these effects were strongest at a buffer radius of 1 and 3.5 km, respectively. Possibly, continuous food resources provided by natural forest areas favor colony establishment and growth, and/or increase overwintering survival of bumblebee queens. Also, pollinator dependent crop area can generate a “transient dilution effect” by decreasing the density of bumblebees in simultaneously flowering crops. Management strategies might increase crop pollination services by considering the importance of nesting and overwintering habitat quality/amount and area of simultaneously flowering crops requiring insect pollination.

Vaccinium corymbosum / pollinators / bumblebee / natural forest areas / flowering crops

1. INTRODUCTION

Insect pollination is important to commercial blueberry production, and farmers usually rent honeybee (*Apis mellifera*) hives and, when possible, also buy *Bombus terrestris* L. colonies and place them in the field during the flowering season. Bumblebees are very effective pollinators for this crop compared to other bees (Javorek et al. 2002; Desjardins and de Oliveira 2006). They can work relatively early in the season when most blueberries flower, at cool temperatures unfavorable to honeybee pollination (Corbet et al. 1993).

According to a review by Garibaldi et al. (2011) the proximity of natural areas can increase bumblebee density in crop areas since it can provide floral resources and/or undisturbed nesting/overwintering habitat. Floral resources provided by crop and non-crop areas can also increase bumblebee densities (Westphal et al. 2003; Carvell et al. 2006), but the temporal dynamics of flowering crops alters bumblebee densities as well. Previous studies reported a “transient dilution effect” in which bumblebee densities decrease with increasing area of oilseed rape fields during flowering, both in this crop area and in nearby grassland areas (Holzschuh et al. 2011). Only after the flowering events of oilseed rape do bumblebee densities increase in nearby areas (Westphal et al. 2003; but see Diekötter et al. 2010). Based on these observations, it is possible that bumblebee density in blueberry crops is

Corresponding author: L. Vieli,
lorena.vieli@ufrontera.cl
Handling Editor: Peter Rosenkranz

positively influenced by natural areas and negatively by simultaneously flowering crops in the surrounding landscape. Research questions of this study were as follows: (1) Are there any effects of surrounding natural areas and flowering crop areas on wild bumblebee abundance associated with blueberry fields? (2) What is the relevant spatial scale for these effects? Increasing the understanding of landscape effects on important wild pollinators can provide cues on possible management strategies to increase pollination services for blueberry production.

2. METHODS

2.1. Field samples

Eight blueberry farms in the central valley of the Region La Araucanía, Chile, were studied (Figure 1). Samples were collected from “Briggita” cultivar of highbush blueberry (*Vaccinium corymbosum* L.). Distance between farms ranged from 7.4 to 97.8 km (mean \pm SE of 42.6 ± 21.5 km). Blueberry fields varied in size from 0.5 to 120 ha (mean \pm SE of 45.3 ± 14.4 ha). Commercial honeybee colonies were employed in six farms while four of those also had *B. terrestris* colonies. None of the surveyed farms applied pesticides during flowering. Weeds were suppressed in and around all of the fields.

Sampling of pollinators was performed on two different dates within the flowering season, between October 13 and November 5, 2011, between 1100 to 1700 on days with favorable weather conditions (temperature > 14 °C, dry, wind speed < 5 m/s). All sampled fields had the same plant density (1×3 m). Flower density was assumed to be relatively constant as blueberry variety was the same and plant age was relatively homogeneous (5–7 years). Depending on the field size, four, five or six sites per farm were sampled; in one case, only one site was sampled because of the limited field size (0.5 ha). Sampled sites were placed at different locations within the fields including field edges and field centers, in order to capture possible variation of pollinator densities. Each sampling event ($n = 68$) consisted of separate counts of four successive rows in which a person walked through ten consecutive plants in a row for 5 min, recording all insects that visited flowers. All row counts were summed to produce a site-level estimate of abundance. Sampling was

performed when the proportion of open flowers on ten randomly chosen branches was ≥ 0.2 .

Bumblebees and honeybees were visually identified to the species level and other pollinators were recorded as other hymenopterans, syrphids, or other. During sampling wild bumblebee workers were not yet active and thus all sampled workers were assumed to be from commercial colonies placed by farmers. At the time of sampling no workers were observed in the area, except for those provided by commercial colonies.

2.2. Land cover

Natural forest areas and high-food-resources areas (i.e., blueberry or rapeseed crops flowering during the sampling period) surrounding the blueberry fields were mapped in a radius of 3.50 km from the center of the fields, based on orthorectified aerial photos acquired between 2008 and 2010 (2-m spatial resolution) depending on the location, and high resolution imagery available from Google Earth (<http://earth.google.com>). The software ArcGIS 9.3 (ESRI Inc., Redland, CA) was used for this purpose. Natural forest areas correspond to unmanaged secondary forests and do not include exotic pine and eucalyptus plantations. High-food resource areas were mapped using the imagery and identified via visual inspection as food resource if flowering in the mapped area at the time of the surveys. The influence of landscape context on pollinator abundance was analyzed at different spatial scales using circular neighborhoods centered on the central point of sampled sites within the farm with radii of 0.5, 1.0, 1.5, 2.0, 2.5, 3.0, and 3.5 km.

To explore larger landscape effects, circular areas of 5- and 8-km radii were used from on a land cover map of 15-m spatial resolution. This map was generated by classification of an ASTER satellite image from 2008 (Altamirano et al., unpublished data) and was used to measure only natural forest areas at these scales. One farm at these scales of analysis was excluded because its buffer overlapped with neighboring farms.

The proportion of natural forest and high-food resource areas varied across farms and both tended to decrease with increasing spatial scale (Table 1).

2.3. Statistical analysis

Analysis of landscape effects focused on the relationship between the proportion of forest area and high-




Figure 1. Sampled farm locations in south-central Chile.

Table I. Proportion of natural forest and high-food resource (flowering crops) areas (Mean[Min, Max]) for sampled farms at different radius distances. For simplicity some spatial scales are omitted.

Scale (m)	Natural forest	High-food resources
500	0.17 [0.02, 0.31]	0.32 [0.06, 0.73]
1500	0.13 [0.04, 0.24]	0.08 [0.01, 0.36]
2500	0.13 [0.04, 0.24]	0.04 [0.002, 0.14]
3500	0.12 [0.04, 0.29]	0.02 [0.001, 0.07]
5000	0.07 [0.02, 0.20]	–
8000	0.08 [0.03, 0.22]	–

food-resources area with the abundance of wild *B. terrestris* queens. This group was the most abundant among sampled pollinators.

Bumblebee abundances were averaged for each farm and across sampling dates. Linear regression models to predict farm-level bumblebee abundance were fitted as a function of the proportion of forest area and high-food resource area, accounting for unequal number of sites per farm weighting each observation by the number of sites of that farm. Models with two covariates (forest areas and high-food resources areas) were tested at each scale of analysis considering also all combinations of different spatial scales. In all models, \log_e - \log_e transformation of the data was used to improve linearity. Correlation between forest and high-food resource areas

at each spatial scale was tested using the Pearson correlation coefficient. R v.2.15.0 (R Development Core Team, 2012) was used for statistical analysis. The possible effect of commercial bumblebee and honeybee colonies was not evaluated. These colonies are placed in the field only during blueberry flowering and competition for food resources is unlikely given the over abundance of this resource.

3. RESULTS

Honeybees were the most abundant pollinator sampled, followed by naturally occurring *B. terrestris* queens and syrphids (Figure 2). Few individuals of *Bombus ruderatus* and *Bombus dahlbomii*, or other hymenopterans were found. In farms stocked with commercial bumblebee colonies, *B. terrestris* workers were also observed.

The abundance of wild *B. terrestris* queens was positively associated with the area of natural forest and negatively associated with areas of high-food resources. These associations were significant at various spatial scales (Figure 3) and peaked at 1- and 3.5-km radii for forest and high-food resources, respectively (Table II). No correlation was found between natural forest area and

high-food resource area at any spatial scale. However positive correlation of the proportion of natural forest and high-food resources areas between similar spatial scales (Pearson's $r > 0.8$) was found.

4. DISCUSSION

Among wild pollinators, *B. terrestris* queens were most abundant reflecting a successful spread of this exotic species in the study area. The only native bumblebee (*B. dahlbomii*) was almost absent which is consistent by previous reports on the decline of this species after the introduction of *B. ruderatus* and *B. terrestris* (Morales et al. 2013) in 1982–1983 and 1997–1998, respectively (Montalva et al. 2011; Smith-Ramírez et al. 2014).

The positive relationship between natural forest area and bumblebee abundance might reflect nesting suitability and/or hibernation habitat requirements of queen bumblebees. Undisturbed areas such as forests and forest edges can be suitable nesting habitat for bumblebees (Kells and Goulson 2003; Osborne et al. 2008). These areas might be a source of continuous pollen and nectar resources throughout the foraging stage of bumblebees. While food supply in early spring




Figure 2. Abundance of different pollinators in sampled sites. Bars of the boxplot represent the median, and the 25 and 75 % quartile boundaries, error bars represent range of values (excluding outliers) and points represent outliers.




Figure 3. R2 value for linear models to explain queen bumblebee abundances. Covariates of fitted models are fraction of natural forest and high-food-resources in circular areas of different radius distances (represented by x and y axis respectively). An asterisk indicates that both covariates were significant at $\alpha < 0.05$.

can favor bumblebee colony establishment and initial growth (Westphal et al. 2009), the reproductive success of the colony seems to be determined by late-season food availability provided by surrounding natural areas (Persson and Smith 2013). Late-season food supply is crucial for hibernating queens since Beekman et al. (1998) found that body weight at the start of diapause positively affects its success, while environmental temperature has no effect. Thus, forests can favor the number of queens produced per colony and/or favor hibernation success by providing food supply for queens before they start diapause. Further studies should focus on the relevance of food provision by natural forests and the type and

quality of nesting/overwintering sites for bumblebees in south-central Chile.

The results of this study also suggest that the presence of nearby flowering crops used by these pollinators can decrease the abundance of queen bumblebees in blueberry fields. This negative effect of high-food resource areas could correspond to a “transient dilution effect” in which the bumblebee density decreases as they are spread across larger areas of flowering crops and is consistent with previous studies that document such effect on oilseed rape (Holzschuh et al. 2011). Simultaneously flowering areas probably compete for highly mobile pollinators such as bumblebees when they are available within the foraging range

Table II. Results from the best fitted regression model of queen *B. terrestris* abundances, weighted by the number of sites per farm sampled. Covariates and response variable were \log_e transformed. For each covariate (fraction of land cover type) the radius of circular areas used is given in parenthesis.

Variable	Estimate	t	P	R2
Intercept	-0.15	-0.33	0.76	0.89
Natural forest (1000 m)	0.51	3.93	0.011	
High-food resources (3500 m)	-0.24	-3.77	0.013	

of the individuals. From the perspective of crop pollination services provision, this effect is important since the pollination service can decrease not only due to lack of neighboring natural areas but also nearby flowering crop area.

ACKNOWLEDGMENTS

We are grateful to all the farms that gave us permission to sample in their fields. We also thank Francisco Contreras who helped in collecting field data and Prof. R. Rebolledo for assistance in species identification. This research was supported by a Fulbright/Conicyt Chile Scholarship.

Effets du paysage sur les reines du bourdon sauvage *Bombus terrestris* (Hymenoptera: Apidae) visitant les champs de myrtille arbustive dans le sud du Chili central

***Vaccinium corymbosum* / pollinisateur/ zone de forêt naturelle/ cultures au stade floraison**

Landschaftseinflüsse auf wildlebende *Bombus terrestris* Königinnen beim Besuch von Heidelbeeren in Südchile

***Vaccinium corymbosum* / Bestäuber/ Hummeln/ natürliche Waldareale/ blühende Nutzpflanzen**

REFERENCES

- Beekman, M., Stratum, P.V., Lingeman, R. (1998) Diapause survival and post-diapause performance in bumblebee queens (*Bombus terrestris*). *Ent. Exp. Appl.* **89**, 207–214
- Carvell, C., Meek, W.R., Pywell, R.F., Goulson, D., Nowakowski, M. (2006) Comparing the efficacy of agri-environment schemes to enhance bumble bee abundance and diversity on arable field margins. *J. Appl. Ecol.* **44** (1), 29–40
- Corbet, S.A., Fussell, M., Ake, R., Fraser, A., Gunson, C., Savage, A., Smith, K. (1993) Temperature and the pollinating activity of social bees. *Ecol. Entomol.* **18**, 17–30
- Desjardins, E.C., de Oliveira, D. (2006) Commercial bumble bee *Bombus impatiens* (Hymenoptera: Apidae) as a pollinator in lowbush blueberry (Ericaceae: Ericaceae) fields. *J. Econ. Entomol.* **99** (2), 443–449
- Diekötter, T., Kadoya, T., Peter, F., Wolters, V., Jauker, F. (2010) Oilseed rape crops distort plant-pollinator interactions. *J. Appl. Ecol.* **47** (1), 209–214
- Garibaldi, L.A., Steffan-Dewenter, I., Kremen, C., Morales, J.M., Bommarco, R., Cunningham, S.A., et al. (2011) Stability of pollination services decreases with isolation from natural areas despite honey bee visits. *Ecol. Lett.* **14** (10), 1062–1072
- Holzschuh, A., Dormann, C.F., Tschamtkke, T., Steffan-Dewenter, I. (2011) Expansion of mass-flowering crops leads to transient pollinator dilution and reduced wild plant pollination. *Proc. R. Soc. B* **278**, 3444–3451
- Javorek, S.K., Mackenzie, K.E., Vander Kloet, S.P. (2002) Comparative pollination effectiveness among bees (Hymenoptera: Apoidea) on lowbush blueberry (Ericaceae: *Vaccinium angustifolium*). *Ann. Entomol. Soc. Am.* **95** (3), 345–351
- Kells, A.R., Goulson, D. (2003) Preferred nesting sites of bumblebee queens (Hymenoptera: Apidae) in agroecosystems in the UK. *Biol. Conserv.* **109** (2), 165–174
- Montalva, J., Dudley, L., Arroyo, M.K., Retamales, H., Abrahamovich, A.H. (2011) Geographic distribution and associated flora of native and introduced bumble bees (*Bombus* spp.) in Chile. *J. Apic. Res.* **50** (1), 11–21
- Morales, C.L., Arbetman, M.P., Cameron, S.A., Aizen, M.A. (2013) Rapid ecological replacement of a native bumble bee by invasive species. *Front. Ecol. Environ.* **11** (10), 529–534
- Osborne, J.L., Martin, A.P., Carreck, N.L., Swain, J.L., Knight, M.E., Goulson, D., Hale, R.J., Sanderson, R.A. (2008) Bumblebee flight distances in relation to the forage landscape. *J. Anim. Ecol.* **77** (2), 406–415
- Persson, A.S., Smith, H.G. (2013) Seasonal persistence of bumblebee populations is affected by landscape context. *Agric. Ecosyst. Environ.* **165**, 201–209
- R Development Core Team (2012) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Smith-Ramírez, C., Ramos-Jiliberto, R., Valdovinos, F.S., Martínez, P., Castillo, J., Armesto, J.J. (2014) Decadal trends in the pollinator assemblage of *Eucryphia cordifolia* in Chilean rainforests. *Oecologia* **176** (1), 157–169
- Westphal, C., Steffan-Dewenter, I., Tschamtkke, T. (2003) Mass flowering crops enhance pollinator densities at a landscape scale. *Ecol. Lett.* **6** (11), 961–965
- Westphal, C., Steffan-Dewenter, I., Tschamtkke, T. (2009) Mass flowering oilseed rape improves early colony growth but not sexual reproduction of bumblebees. *J. Appl. Ecol.* **46** (1), 187–193