Erratum to: Winter turnip rape as a soil N scavenging catch crop in a cool humid climate

Antti Tuulos • Markku Yli-Halla • Frederick Stoddard •
Pirjo Mäkelä

Published online: 27 June 2014
© INRA and Springer-Verlag France 2014

Erratum to: Agron Sustain Dev
Doi 10.1007/s13593-014-0229-2

In Table 1, the amount of NO_3^- -N in subsoil before growing season 2010–2011 is 15.9 kg/ha for all treatments instead of 4.6 kg/ha.

The author affiliation for Antti Tuulos, Frederick Stoddard and Pirjo Mäkelä is Department of Agricultural Sciences instead of Department of Food and Environmental Sciences.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s13593-014-0229-2>.

A. Tuulos (✉) • F. Stoddard • P. Mäkelä
Department of Agricultural Sciences, University of Helsinki,
P.O. Box 27, 00014 Helsinki, Finland
e-mail: antti.tuulos@helsinki.fi

M. Yli-Halla
Department of Food and Environmental Sciences, University of
Helsinki, P.O. Box 27, 00014 Helsinki, Finland

New version of Table 1.

Table 1 Amount of soil mineral nitrogen ($\text{NH}_4^+\text{-N}$, ammonium-nitrogen; $\text{NO}_3^-\text{-N}$, nitrate-nitrogen) species (kg N/ha) in topsoil (0–20 cm) and subsoil (30–50 cm) under different combinations of barley and winter turnip rape (WTR) and management practices. Data shown are means ($n=4$), except before growing season, in which case the results represent the mean ($n=4$) of the entire experimental area

Year	Soil layer	Crop and management	Before growing season		After barley harvest		Before winter		After winter	
			$\text{NH}_4^+\text{-N}$	$\text{NO}_3^-\text{-N}$	$\text{NH}_4^+\text{-N}$	$\text{NO}_3^-\text{-N}$	$\text{NH}_4^+\text{-N}$	$\text{NO}_3^-\text{-N}$	$\text{NH}_4^+\text{-N}$	$\text{NO}_3^-\text{-N}$
2009–2010	Topsoil	WTR, sown with barley	17.4	28.5	14.3 a	7.3 a	19.6 a	7.7 a	22.9 a	13.9 a
		WTR, sown after barley	17.4	28.5	13.5 a	10.2 a	19.3 ab	8.4 a	17.8 a	10.9 a
		Barley, left to stubble	17.4	28.5	17.4 a	11.1 a	18.1 ab	9.9 a	16.1 a	12.8 a
		Barley, ploughed	17.4	28.5	18.1 a	12.5 a	12.2 b	15.6 b	17.5 a	12.6 a
		Standard error of the mean	0.6	1.9	1.6	1.7	1.2	1.0	3.5	1.2
	Subsoil	WTR, sown with barley	NA	NA	11.1 a	5.5 a	7.3 a	3.7 a	9.8 a	12.7 a
		WTR, sown after barley	NA	NA	8.4 a	6.2 a	7.1 a	3.1 a	10.2 a	11.4 a
		Barley, left to stubble	NA	NA	8.9 a	7.9 a	6.8 a	7.1 a	8.1 a	13.7 a
		Barley, ploughed	NA	NA	9.9 a	7.1 a	8.0 a	18.5 b	9.5 a	11.9 a
		Standard error of the mean	NA	NA	1.8	1.5	0.5	2.3	1.2	2.7
2010–2011	Topsoil	WTR, sown with barley	5.5	35.5	5.5 a	27.0 a	5.4 a	18.6 a	3.7 a	24.5 a
		WTR, sown after barley	5.5	35.5	13.0 a	40.5 a	7.6 a	28.7 a	3.6 a	23.5 a
		Barley, left to stubble	5.5	35.5	4.2 a	37.0 a	6.0 a	32.5 ab	5.1 a	24.0 a
		Barley, ploughed	5.5	35.5	4.4 a	35.5 a	4.3 a	44.5 b	3.2 a	23.5 a
		Standard error of the mean	1.8	0.5	2.7	6.2	1.9	3.4	1.0	1.5
	Subsoil	WTR, sown with barley	4.6	15.9	6.2 a	13.0 a	3.8 a	8.3 a	3.4 a	18.6 a
		WTR, sown after barley	4.6	15.9	5.8 a	13.2 a	3.1 a	14.9 ab	5.6 a	24.1 a
		Barley, left to stubble	4.6	15.9	6.0 a	15.6 a	3.7 a	21.1 b	3.0 a	18.7 a
		Barley, ploughed	4.6	15.9	5.3 a	16.0 a	5.3 a	21.5 b	4.8 a	21.2 a
		Standard error of the mean	0.5	1.8	1.2	1.4	1.0	2.2	1.4	2.2

Within a year and soil layer, means followed by the same letter are not significantly different at $P=0.05$ by Tukey's test.