

# Graph convergence and generalized Yosida approximation operator with an application

Rais Ahmad<sup>1</sup> · Mohd. Ishtyak<sup>1</sup> · Mijanur Rahaman<sup>1</sup> · Iqbal Ahmad<sup>1</sup>

Received: 13 July 2016 / Accepted: 3 March 2017 / Published online: 30 March 2017  
© The Author(s) 2017. This article is an open access publication

**Abstract** In this paper, we introduce a Yosida inclusion problem as well as a generalized Yosida approximation operator. Using the graph convergence of  $H(\cdot, \cdot)$ -accretive operator and resolvent operator convergence discussed in Li and Huang (Appl Math Comput 217:9053–9061, 2011), we establish the convergence for generalized Yosida approximation operator. As an application, we solve a Yosida inclusion problem in  $q$ -uniformly smooth Banach spaces. An example is constructed, and through MATLAB programming, we show some graphics for the convergence of generalized Yosida approximation operator.

**Keywords** Graph convergence · Resolvent operator · Smooth Banach space · Yosida approximation operator · Yosida inclusion

**Mathematics Subject Classification** 47H09 · 49J40 · 65K15

---

✉ Mijanur Rahaman  
mrahaman96@yahoo.com

Rais Ahmad  
raisain\_123@rediffmail.com

Mohd. Ishtyak  
ishtyakalig@gmail.com

Iqbal Ahmad  
iqbalahmad120@gmail.com

<sup>1</sup> Department of Mathematics, Aligarh Muslim University, Aligarh 202002, India

## Introduction

A reasonable attention has been shown by many researchers for the study of variational inclusions (inequalities) and their generalized forms, which occupies a leading and significant role to connect research between analysis, geometry, biology, elasticity, optimization, image processing, biomedical and mathematical sciences, etc. A broad range of problems with which we encounter in physics, economics, management sciences, and operations research can be formulated as an inclusion problem  $0 \in T(x)$ , for a given set-valued mapping  $T$  on a Hilbert space  $H$ . Thus, the problem of finding a zero of  $T$ , i.e., a point  $x \in H$ , such that  $0 \in T(x)$  is a fundamental problem in many areas of applied sciences.

On the other hand, it is well known that monotone operators on Hilbert spaces can be regularized into single-valued Lipschitzian monotone operators via a process known as the Yosida approximation. This Yosida approximation operators are instrumental to approximate the solutions of general variational inclusion problems using non-expansive resolvent operators. Recently, many authors [2, 3, 5, 6, 8–10] have applied Yosida approximation operators and their generalized forms to solve some variational inclusion problems. Zou and Huang [14], Ahmad et al. [1] introduced and studied the graph convergence of  $H(\cdot, \cdot)$ -accretive operators and  $H(\cdot, \cdot)$ -co-accretive operators, respectively, for solving variational inclusion problems and their system. For more details, we refer to [4, 11, 12, 15].

This paper deals with the introduction of a generalized Yosida approximation operator with some of its properties. Under the concept of graph convergence of  $H(\cdot, \cdot)$ -accretive operators, we prove the convergence of generalized Yosida approximation operator. Finally, we solve a Yosida

inclusion problem in  $q$ -uniformly Banach spaces. A MATLAB programming related to graph convergence of generalized Yosida approximation operator is discussed with a consolidated example. Our results are applicable and new in this direction and refinement of results of Li and Huang [7].

### Preliminaries

Let  $X$  be a real Banach Space with its dual space  $X^*$ . We denote the duality pairing between  $X$  and  $X^*$  by  $\langle \cdot, \cdot \rangle$ , and  $2^X$  is the family of all nonempty subsets of  $X$ .

The generalized duality mapping  $F_q : X \rightarrow 2^{X^*}$  is defined by

$$F_q(x) = \left\{ f^* \in X^* : \langle x, f^* \rangle = \|x\|^q, \|f^*\| = \|x\|^{q-1} \right\}, \quad \forall x \in X,$$

where  $q > 1$  is a constant. For  $q = 2$ ,  $F_q$  coincides with the normalized duality mapping. If  $X$  is a Hilbert space,  $F_2$  becomes the identity mapping on  $X$ . It is to be noted that if  $X$  is uniformly smooth, then  $F_q$  is single-valued. Throughout the paper, we assume that  $X$  is a real Banach space and  $F_q$  is single-valued.

The function  $\rho_X : [0, \infty) \rightarrow [0, \infty)$  is called modulus of smoothness of  $X$ , such that

$$\rho_X(t) = \left\{ \frac{\|x+y\| + \|x-y\|}{2} - 1 : \|x\| \leq 1, \|y\| \leq t \right\}.$$

A Banach space  $X$  is called

1. uniformly smooth if  $\lim_{t \rightarrow 0} \frac{\rho_X(t)}{t} = 0$ ;
2.  $q$ -uniformly smooth if there exists a constant  $c > 0$ , such that

$$\rho_X(t) \leq ct^q, \quad q > 1.$$

While encountered with the characteristic inequalities, Xu [13] proved the following important Lemma in  $q$ -uniformly smooth Banach spaces.

**Lemma 1** *Let  $X$  be a real uniformly smooth Banach space. Then,  $X$  is  $q$ -uniformly smooth if and only if there exists a constant  $c_q > 0$ , such that for all  $x, y \in X$ ,*

$$\|x+y\|^q \leq \|x\|^q + q\langle y, F_q(x) \rangle + c_q\|y\|^q.$$

The following definitions and concepts are essential to achieve the aim of this paper.

**Definition 1** [14] Let  $A, B : X \rightarrow X$  and  $H : X \times X \rightarrow X$  be the single-valued mappings.

1.  $A$  is said to be accretive, if

$$\langle A(x) - A(y), F_q(x - y) \rangle \geq 0, \quad \forall x, y \in X;$$

2.  $A$  is said to be strictly accretive, if  $A$  is accretive and  $\langle A(x) - A(y), F_q(x - y) \rangle = 0$ , if and only if  $x = y$ ;
3.  $A$  is said to be  $\delta_A$ -strongly accretive, if there exists a constant  $\delta_A > 0$ , such that  $\langle Ax - Ay, F_q(x - y) \rangle \geq \delta_A\|x - y\|^q$ ;
4.  $A$  is said to be  $\gamma_A$ -Lipschitz continuous, if there exists a constant  $\gamma_A > 0$ , such that  $\|Ax - Ay\| \leq \gamma_A\|x - y\|, \quad \forall x, y \in X$ ;
5.  $H(A, \cdot)$  is said to be  $\alpha$ -strongly accretive with respect to  $A$ , if there exists a constant  $\alpha > 0$ , such that  $\langle H(Ax, \cdot) - H(Ay, \cdot), F_q(x - y) \rangle \geq \alpha\|x - y\|^q, \quad \forall x, y \in X$ ;
6.  $H(\cdot, B)$  is said to be  $\beta$ -relaxed accretive with respect to  $B$ , if there exists a constant  $\beta > 0$ , such that  $\langle H(\cdot, Bx) - H(\cdot, By), F_q(x - y) \rangle \geq -\beta\|x - y\|^q, \quad \forall x, y \in X$ ;
7.  $H(A, \cdot)$  is said to be  $\sigma$ -Lipschitz continuous with respect to  $A$ , if there exists a constant  $\sigma > 0$ , such that  $\|H(Ax, \cdot) - H(Ay, \cdot)\| \leq \sigma\|x - y\|, \quad \forall x, y \in X$ .

Similarly, we can define the Lipschitz continuity of  $H$  with respect to  $B$ .

**Definition 2** [14] Let  $H : X \rightarrow X$  be a single-valued mapping and  $M : X \rightarrow 2^X$  be a set-valued mapping. The mapping  $M$  is said to be

1. accretive, if  $\langle u - v, F_q(x - y) \rangle \geq 0, \quad \forall x, y \in X, u \in M(x), v \in M(y)$ ;
2.  $m$ -accretive, if  $M$  is accretive and  $(I + \lambda M)(X) = X$ , for all  $\lambda > 0$ , where  $I$  is the identity operator on  $X$ ;
3.  $H$ -accretive, if  $M$  is accretive and  $(H + \lambda M)(X) = X$ , for all  $\lambda > 0$ .

**Definition 3** [14] Let  $A, B : X \rightarrow X, H : X \times X \rightarrow X$  be the single-valued mappings and  $M : X \rightarrow 2^X$  be a set-valued mapping. The mapping  $M$  is said to be  $H(\cdot, \cdot)$ -accretive with respect to  $A$  and  $B$ , if  $M$  is accretive and  $[H(A, B) + \lambda M](X) = X$ , for every  $\lambda > 0$ .

**Lemma 2** [14] *Let  $H(A, B)$  be  $\alpha$ -strongly accretive with respect to  $A$ ,  $\beta$ -relaxed accretive with respect to  $B$  and  $\alpha > \beta$ . Let  $M$  be an  $H(\cdot, \cdot)$ -accretive operator with respect to  $A$  and  $B$ . Then, the operator  $[H(A, B) + \lambda M]^{-1}$  is single-*

valued and is called the resolvent operator, i.e.,  $R_{M,\lambda}^{H(\cdot,\cdot)} : X \rightarrow X$ , such that

$$R_{M,\lambda}^{H(\cdot,\cdot)}(u) = [H(A,B) + \lambda M]^{-1}(u), \quad \forall u \in X, \quad \lambda > 0. \quad (1)$$

Furthermore, the resolvent operator defined by Eq. (1) is  $\frac{1}{(\alpha-\beta)}$ -Lipschitz continuous.

**Lemma 3** [3] Let  $\{a_n\}$  and  $\{b_n\}$  be two non-negative real sequences satisfying

$$a_{n+1} \leq ka_n + b_n,$$

with  $0 < k < 1$  and  $b_n \rightarrow 0$ . Then,  $\lim_{n \rightarrow \infty} a_n = 0$ .

### Generalized Yosida approximation operator and its convergence

We define the generalized Yosida approximation operator using the resolvent operator defined by Eq. (1), that is

$$R_{M,\lambda}^{H(\cdot,\cdot)}(u) = [H(A,B) + \lambda M]^{-1}(u), \quad \forall x \in X, \quad \lambda > 0.$$

**Definition 4** The generalized Yosida approximation operator denoted by  $J_{M,\lambda}^{H(\cdot,\cdot)}$  is defined as

$$J_{M,\lambda}^{H(\cdot,\cdot)}(u) = \frac{1}{\lambda} [I - R_{M,\lambda}^{H(\cdot,\cdot)}](u), \quad \forall u \in X \quad \text{and} \quad \lambda > 0, \quad (2)$$

where  $I$  is the identity mapping on  $X$ .

**Lemma 4** The generalized Yosida approximation operator defined by Eq. (2) is

- $\theta_1$ -Lipschitz continuous, where  $\theta_1 = \frac{[\alpha-\beta+1]}{\lambda(\alpha-\beta)}, \alpha > \beta$ .
- $\theta_2$ -strongly monotone, where  $\theta_2 = \frac{[(\alpha-\beta)-1]}{\lambda(\alpha-\beta)}, \alpha > \beta$ .

*Proof*

- Let  $u, v \in X$  and  $\lambda > 0$ . Using Lemma 2, we have

$$\begin{aligned} & \left\| J_{M,\lambda}^{H(\cdot,\cdot)}(u) - J_{M,\lambda}^{H(\cdot,\cdot)}(v) \right\| \\ &= \frac{1}{\lambda} \left\| [I(u) - R_{M,\lambda}^{H(\cdot,\cdot)}(u)] - [I(v) - R_{M,\lambda}^{H(\cdot,\cdot)}(v)] \right\| \\ &\leq \frac{1}{\lambda} \left[ \|u - v\| + \left\| R_{M,\lambda}^{H(\cdot,\cdot)}(u) - R_{M,\lambda}^{H(\cdot,\cdot)}(v) \right\| \right] \\ &\leq \frac{1}{\lambda} \left[ \|u - v\| + \frac{1}{[\alpha - \beta]} \|u - v\| \right] \\ &= \frac{1}{\lambda} \left[ \frac{\alpha - \beta + 1}{\alpha - \beta} \right] \|u - v\|, \end{aligned}$$

i.e.,

$$\left\| J_{M,\lambda}^{H(\cdot,\cdot)}(u) - J_{M,\lambda}^{H(\cdot,\cdot)}(v) \right\| \leq \theta_1 \|u - v\|, \quad (3)$$

where  $\theta_1 = \frac{[\alpha-\beta+1]}{\lambda(\alpha-\beta)}, \alpha > \beta$ .

- For any  $u, v \in X$ , and  $\lambda > 0$  and using Lemma 2, we have

$$\begin{aligned} & \left\langle J_{M,\lambda}^{H(\cdot,\cdot)}(u) - J_{M,\lambda}^{H(\cdot,\cdot)}(v), F_q(u - v) \right\rangle \\ &= \frac{1}{\lambda} \left\langle [I(u) - R_{M,\lambda}^{H(\cdot,\cdot)}(u)] - [I(v) - R_{M,\lambda}^{H(\cdot,\cdot)}(v)], F_q(u - v) \right\rangle \\ &= \frac{1}{\lambda} \left[ \left\langle u - v, F_q(u - v) \right\rangle - \left\langle R_{M,\lambda}^{H(\cdot,\cdot)}(u) - R_{M,\lambda}^{H(\cdot,\cdot)}(v), F_q(u - v) \right\rangle \right] \\ &\geq \frac{1}{\lambda} \left[ \|u - v\|^q - \left\| R_{M,\lambda}^{H(\cdot,\cdot)}(u) - R_{M,\lambda}^{H(\cdot,\cdot)}(v) \right\| \|u - v\|^{q-1} \right] \\ &\geq \frac{1}{\lambda} \left[ \|u - v\|^q - \frac{1}{[\alpha - \beta]} \|u - v\| \|u - v\|^{q-1} \right] \\ &= \frac{1}{\lambda} \left[ \|u - v\|^q - \frac{1}{[\alpha - \beta]} \|u - v\|^q \right] \\ &= \frac{[(\alpha - \beta) - 1]}{\lambda(\alpha - \beta)} \|u - v\|^q. \end{aligned}$$

i.e.,

$$\left\langle J_{M,\lambda}^{H(\cdot,\cdot)}(u) - J_{M,\lambda}^{H(\cdot,\cdot)}(v), F_q(u - v) \right\rangle \geq \theta_2 \|u - v\|^q, \quad \forall u, v \in X, \lambda > 0$$

and  $\theta_2 = \frac{[(\alpha-\beta)-1]}{\lambda(\alpha-\beta)}, \alpha > \beta$ .

□

*Note 1* It is interesting to note that resolvent operator defined by Eq. (1) and generalized Yosida approximation operator defined by Eq. (2) are connected by the following relation:

$$\lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x) \in [\lambda M + H(A, B) - I] \left( R_{M,\lambda}^{H(\cdot,\cdot)}(x) \right).$$

Let  $M : X \rightarrow 2^X$  be a set-valued mapping. The graph of the mapping  $M$  is defined by

$$graph(M) = \{(x, y) \in X \times Y : y \in M(x)\}$$

**Definition 5** [7] Let  $A, B : X \rightarrow X$  and  $H : X \times X \rightarrow X$  be the single-valued mappings. Let  $M_n, M : X \rightarrow 2^X$  be  $H(\cdot, \cdot)$ -accretive operators for  $n = 0, 1, 2, \dots$ . The sequence  $\{M_n\}$  is said to be graph convergence to  $M$ , denoted by  $M_n \xrightarrow{G} M$ , if for every  $(x, y) \in graph(M)$ , there exists a sequence  $(x_n, y_n) \in graph(M_n)$ , such that

$$x_n \rightarrow x, \quad y_n \rightarrow y \quad \text{as } n \rightarrow \infty.$$

**Theorem 1** [7] *Let  $M_n, M : X \rightarrow 2^X$  be  $H(\cdot, \cdot)$ -accretive operators for  $n = 0, 1, 2, \dots$ . Assume that  $H : X \times X \rightarrow X$  is a single-valued mapping, such that*

1.  $H(A, B)$  is  $\alpha$ -strongly accretive with respect to  $A$  and  $\beta$ -relaxed accretive with respect to  $B$ ,  $\alpha > \beta$ ;
2.  $H(A, B)$  is  $\gamma_1$ -Lipschitz continuous with respect to  $A$  and  $\gamma_2$ -Lipschitz continuous with respect to  $B$ .

Then,  $M_n \xrightarrow{G} M$  if and only if

$$R_{M_n, \lambda}^{H(\cdot, \cdot)}(u) \rightarrow R_{M, \lambda}^{H(\cdot, \cdot)}(u), \quad \forall u \in X, \quad \lambda > 0,$$

where  $R_{M_n, \lambda}^{H(\cdot, \cdot)} = [H(A, B) + \lambda M_n]^{-1}$  and  $R_{M, \lambda}^{H(\cdot, \cdot)} = [H(A, B) + \lambda M]^{-1}$ .

Now, we prove the convergence of generalized Yosida approximation operator in the light of graph convergence of  $H(\cdot, \cdot)$ -accretive operator without using the convergence of resolvent operator defined by Eq. (1).

**Theorem 2** *Let  $M_n, M : X \rightarrow 2^X$  be  $H(\cdot, \cdot)$ -accretive operators for  $n = 0, 1, 2, \dots$ , and  $H : X \times X \rightarrow X$  be a single-valued mapping, such that conditions (1) and (2) of Theorem 1 hold.*

Then  $M_n \xrightarrow{G} M$  if and only if

$$J_{M_n, \lambda}^{H(\cdot, \cdot)}(x) \rightarrow J_{M, \lambda}^{H(\cdot, \cdot)}(x), \quad \forall x \in X, \quad \lambda > 0,$$

where

$$J_{M_n, \lambda}^{H(\cdot, \cdot)}(x) = \frac{1}{\lambda} [I - R_{M_n, \lambda}^{H(\cdot, \cdot)}](x),$$

$$J_{M, \lambda}^{H(\cdot, \cdot)}(x) = \frac{1}{\lambda} [I - R_{M, \lambda}^{H(\cdot, \cdot)}](x), \quad \forall x \in X,$$

and  $R_{M_n, \lambda}^{H(\cdot, \cdot)}$  and  $R_{M, \lambda}^{H(\cdot, \cdot)}$  are defined in Theorem 1.

*Proof* Necessary part: Suppose that  $M_n \xrightarrow{G} M$ . For any given  $x \in X$ , let

$$z_n = J_{M_n, \lambda}^{H(\cdot, \cdot)}(x) \quad \text{and} \quad z = J_{M, \lambda}^{H(\cdot, \cdot)}(x).$$

Then,

$$z = J_{M, \lambda}^{H(\cdot, \cdot)}(x) = \frac{1}{\lambda} [I - R_{M, \lambda}^{H(\cdot, \cdot)}](x),$$

implies that

$$(x - \lambda z) = R_{M, \lambda}^{H(\cdot, \cdot)}(x) = [H(A, B) + \lambda M]^{-1}(x),$$

i.e.,

$$H(A, B)(x - \lambda z) + \lambda M(x - \lambda z) = x.$$

It follows that

$$\frac{1}{\lambda} [x - H(A, B)(x - \lambda z)] \in M(x - \lambda z).$$

That is

$$\left( x - \lambda z, \frac{1}{\lambda} [x - H(A, B)(x - \lambda z)] \right) \in \text{graph}(M).$$

By Definition 4, there exists a sequence  $(w_n, y_n) \in \text{graph}(M_n)$ , such that

$$w_n \rightarrow (x - \lambda z), \quad y_n \rightarrow \frac{1}{\lambda} [x - H(A, B)(x - \lambda z)]. \quad (4)$$

Since  $y_n \in M_n(w_n)$ , we have

$$H(Aw_n, Bw_n) + \lambda y_n \in [H(A, B) + \lambda M_n](w_n),$$

and so,

$$\begin{aligned} w_n &= [H(A, B) + \lambda M_n]^{-1} [H(Aw_n, Bw_n) + \lambda y_n], \\ &= R_{M_n, \lambda}^{H(\cdot, \cdot)} [H(Aw_n, Bw_n) + \lambda y_n], \\ &= [I - \lambda J_{M_n, \lambda}^{H(\cdot, \cdot)}]^{-1} [H(Aw_n, Bw_n) + \lambda y_n], \end{aligned}$$

which implies that

$$\frac{1}{\lambda} w_n = \frac{1}{\lambda} H(Aw_n, Bw_n) + y_n - J_{M_n, \lambda}^{H(\cdot, \cdot)} [H(Aw_n, Bw_n) + \lambda y_n]. \quad (5)$$

Using (1) of Lemma 4 and Eq. (5), we have

$$\begin{aligned} \|z_n - z\| &= \|J_{M_n, \lambda}^{H(\cdot, \cdot)}(x) - z\| \\ &= \left\| J_{M_n, \lambda}^{H(\cdot, \cdot)}(x) + \frac{1}{\lambda} w_n - \frac{1}{\lambda} w_n - z \right\| \\ &= \left\| J_{M_n, \lambda}^{H(\cdot, \cdot)}(x) + \frac{1}{\lambda} H(Aw_n, Bw_n) + y_n - J_{M_n, \lambda}^{H(\cdot, \cdot)} [H(Aw_n, Bw_n) + \lambda y_n] - \frac{1}{\lambda} w_n - z \right\| \\ &\leq \left\| J_{M_n, \lambda}^{H(\cdot, \cdot)}(x) - J_{M_n, \lambda}^{H(\cdot, \cdot)} [H(Aw_n, Bw_n) + \lambda y_n] \right\| \\ &\quad + \left\| \frac{1}{\lambda} H(Aw_n, Bw_n) + y_n - \frac{1}{\lambda} w_n - z \right\| \\ &\leq \theta_1 \|x - H(Aw_n, Bw_n) - \lambda y_n\| \\ &\quad + \left\| \frac{1}{\lambda} H(Aw_n, Bw_n) + y_n - \frac{1}{\lambda} x \right\| + \left\| \frac{1}{\lambda} w_n - \frac{1}{\lambda} x + z \right\| \\ &= \left( \theta_1 - \frac{1}{\lambda} \right) \|x - H(Aw_n, Bw_n) - \lambda y_n\| + \frac{1}{\lambda} \|w_n - x + \lambda z\| \\ &= \left( \theta_1 - \frac{1}{\lambda} \right) \|x - H(Aw_n, Bw_n) + H(A, B)(x - \lambda z) \\ &\quad - H(A, B)(x - \lambda z) - \lambda y_n\| + \frac{1}{\lambda} \|w_n - x + \lambda z\| \\ &\leq \left( \theta_1 - \frac{1}{\lambda} \right) \|x - H(A, B)(x - \lambda z) - \lambda y_n\| \\ &\quad + \left( \theta_1 - \frac{1}{\lambda} \right) \|H(A, B)(x - \lambda z) - H(Aw_n, Bw_n)\| \\ &\quad + \frac{1}{\lambda} \|w_n - x + \lambda z\|. \end{aligned} \quad (6)$$

Since  $H$  is  $\gamma_1$ -Lipschitz continuous with respect to  $A$  and  $\gamma_2$ -Lipschitz continuous with respect to  $B$ , we have

$$\begin{aligned} & \left\| H(A, B)(x - \lambda z) - H(A, B)w_n \right\| \\ &= \left\| H(A(x - \lambda z), B(x - \lambda z)) - H(A(x - \lambda z), Bw_n) \right. \\ & \quad \left. + H(A(x - \lambda z), Bw_n) - H(Aw_n, Bw_n) \right\| \\ &\leq \left\| H(A(x - \lambda z), B(x - \lambda z)) - H(A(x - \lambda z), Bw_n) \right\| \\ & \quad + \left\| H(A(x - \lambda z), Bw_n) - H(Aw_n, Bw_n) \right\| \\ &\leq \gamma_2 \|x - \lambda z - w_n\| + \gamma_1 \|x - \lambda z - w_n\| \\ &= (\gamma_1 + \gamma_2) \|x - \lambda z - w_n\|. \end{aligned} \tag{7}$$

Using Eqs. (7), (6) becomes

$$\begin{aligned} \|z_n - z\| &\leq \left( \theta_1 - \frac{1}{\lambda} \right) \|x - H(A, B)(x - \lambda z) - \lambda y_n\| \\ & \quad + \left[ \left( \theta_1 - \frac{1}{\lambda} \right) (\gamma_1 + \gamma_2) + \frac{1}{\lambda} \right] \|w_n - x + \lambda z\|. \end{aligned}$$

By Eq. (4), we have

$$w_n \rightarrow (x - \lambda z), \quad y_n \rightarrow \frac{1}{\lambda} [x - H(A, B)(x - \lambda z)],$$

i.e.,

$$\|w_n - x + \lambda z\| \rightarrow 0, \quad \frac{1}{\lambda} \|x - H(A, B)(x - \lambda z) - \lambda y_n\| \rightarrow 0,$$

and so

$$\|z_n - z\| \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

i.e.,

$$J_{M_n, \lambda}^{H(\cdot, \cdot)}(x) \rightarrow J_{M, \lambda}^{H(\cdot, \cdot)}(x).$$

**Sufficient Part:** Suppose that

$$J_{M_n, \lambda}^{H(\cdot, \cdot)}(x) \rightarrow J_{M, \lambda}^{H(\cdot, \cdot)}(x), \quad \forall x \in X, \quad \lambda > 0.$$

For any  $(x, y) \in \text{graph}(M)$ , we have  $y \in M(x)$ , and hence

$$H(Ax, Bx) + \lambda y \in [H(A, B) + \lambda M](x).$$

Therefore,

$$x = \left[ I - \lambda J_{M, \lambda}^{H(\cdot, \cdot)} \right] (H(Ax, Bx) + \lambda y).$$

Let  $x_n = \left[ I - \lambda J_{M_n, \lambda}^{H(\cdot, \cdot)} \right] (H(Ax, Bx) + \lambda y)$ . This implies that

$$\frac{1}{\lambda} [H(Ax, Bx) - H(Ax_n, Bx_n) + \lambda y] \in M_n(x_n).$$

Let  $y'_n = \frac{1}{\lambda} [H(Ax, Bx) - H(Ax_n, Bx_n) + \lambda y]$  and using the same arguments as for Eq. (7), we have

$$\begin{aligned} \|y'_n - y\| &= \left\| \frac{1}{\lambda} [H(Ax, Bx) - H(Ax_n, Bx_n) + \lambda y] - y \right\| \\ &= \frac{1}{\lambda} \|H(Ax, Bx) - H(Ax_n, Bx_n)\| \\ &= \frac{1}{\lambda} \|H(Ax, Bx) - H(Ax_n, Bx) \\ & \quad + H(Ax_n, Bx) - H(Ax_n, Bx_n)\| \\ &\leq \frac{1}{\lambda} \|H(Ax, Bx) - H(Ax_n, Bx)\| \\ & \quad + \frac{1}{\lambda} \|H(Ax_n, Bx) - H(Ax_n, Bx_n)\| \\ &\leq \left( \frac{\gamma_1 + \gamma_2}{\lambda} \right) \|x_n - x\|. \end{aligned} \tag{8}$$

Using above arguments, we have

$$\begin{aligned} \|x_n - x\| &= \left\| \left( I - \lambda J_{M_n, \lambda}^{H(\cdot, \cdot)} \right) [H(Ax, Bx) + \lambda y] - \left( I - \lambda J_{M, \lambda}^{H(\cdot, \cdot)} \right) \right. \\ & \quad \left. [H(Ax, Bx) + \lambda y] \right\| \\ &= \left\| \left[ \left( I - \lambda J_{M_n, \lambda}^{H(\cdot, \cdot)} \right) - \left( I - \lambda J_{M, \lambda}^{H(\cdot, \cdot)} \right) \right] [H(Ax, Bx) + \lambda y] \right\|. \end{aligned} \tag{9}$$

Since  $J_{M_n, \lambda}^{H(\cdot, \cdot)}(x) \rightarrow J_{M, \lambda}^{H(\cdot, \cdot)}(x)$ , we have from (9) that

$$\|x_n - x\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Thus, from (8), it follows that  $y'_n \rightarrow y$  as  $n \rightarrow \infty$ , i.e.,

$$M_n \xrightarrow{G} M.$$

This completes the proof.  $\square$

Combining Theorems 1 and 2, we have the following remark.

*Remark 1* The convergence of the resolvent operator  $R_{M_n, \lambda}^{H(\cdot, \cdot)}(x) \rightarrow R_{M, \lambda}^{H(\cdot, \cdot)}(x)$ , and the convergence of the generalized Yosida approximation operator  $J_{M_n, \lambda}^{H(\cdot, \cdot)}(x) \rightarrow J_{M, \lambda}^{H(\cdot, \cdot)}(x)$  are equivalent if and only if the operator  $M_n \xrightarrow{G} M$ .

*Proof* Suppose that  $M_n \xrightarrow{G} M$  and  $R_{M_n, \lambda}^{H(\cdot, \cdot)}(x) \rightarrow R_{M, \lambda}^{H(\cdot, \cdot)}(x)$ . Then

$$\begin{aligned} R_{M_n, \lambda}^{H(\cdot, \cdot)}(x) &\rightarrow R_{M, \lambda}^{H(\cdot, \cdot)}(x), \quad \forall x \in X \\ &\Rightarrow \left[ I - R_{M_n, \lambda}^{H(\cdot, \cdot)} \right] (x) \rightarrow \left[ I - R_{M, \lambda}^{H(\cdot, \cdot)} \right] (x) \\ &\Rightarrow \frac{1}{\lambda} \left[ I - R_{M_n, \lambda}^{H(\cdot, \cdot)} \right] (x) \rightarrow \frac{1}{\lambda} \left[ I - R_{M, \lambda}^{H(\cdot, \cdot)} \right] (x) \\ &\Rightarrow J_{M_n, \lambda}^{H(\cdot, \cdot)}(x) \rightarrow J_{M, \lambda}^{H(\cdot, \cdot)}(x), \quad \forall x \in X. \end{aligned}$$

On similar way, we can show that  $J_{M_n, \lambda}^{H(\cdot, \cdot)}(x) \rightarrow J_{M, \lambda}^{H(\cdot, \cdot)}(x)$  implies that  $R_{M_n, \lambda}^{H(\cdot, \cdot)}(x) \rightarrow R_{M, \lambda}^{H(\cdot, \cdot)}(x)$ .  $\square$

We construct the following consolidated example which shows that the mapping  $M$  is  $H(\cdot, \cdot)$ -accretive with respect to  $A$  and  $B$ ,  $M_n \xrightarrow{G} M$  and  $J_{M_n, \lambda}^{H(\cdot, \cdot)} \rightarrow J_{M, \lambda}^{H(\cdot, \cdot)}$ . Through MATLAB programming, we show some graphics for the convergence of generalized Yosida approximation operator.

*Example 1* Let  $X = \mathbb{R}$ ;  $A, B : \mathbb{R} \rightarrow \mathbb{R}$  and  $H : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  be the mappings defined by

$$A(x) = \frac{x^3}{8}, \quad B(x) = \frac{x}{2},$$

and

$$H(A(x), B(x)) = A(x) - B(x), \quad \forall x \in \mathbb{R},$$

with the condition  $x^2 + y^2 + xy \geq 1$ . Suppose  $M_n, M : \mathbb{R} \rightarrow 2^{\mathbb{R}}$  are the set-valued mappings defined by

$$M_n(x) = \frac{x}{2} + \frac{1}{n^2},$$

and

$$M(x) = \frac{x}{2}.$$

Then, for any fixed  $u \in \mathbb{R}$ , we have

$$\begin{aligned} \langle H(Ax, u) - H(Ay, u), x - y \rangle &= \langle Ax - Ay, x - y \rangle \\ &= \frac{1}{8}(x - y)^2(x^2 + y^2 + xy) \\ &\geq \frac{1}{8}(x - y)^2 = \frac{1}{8}\|x - y\|^2. \end{aligned}$$

Hence,  $H(A, B)$  is  $\frac{1}{8}$ -strongly accretive with respect to  $A$ . In addition

$$\begin{aligned} \langle H(u, Bx) - H(u, By), x - y \rangle &= -\langle Bx - By, x - y \rangle \\ &= -\frac{1}{2}(x - y)^2 \geq -\frac{3}{2}(x - y)^2. \end{aligned}$$

Hence,  $H(A, B)$  is  $\frac{3}{2}$ -relaxed accretive with respect to  $B$ .

One can easily verify that for  $\lambda = 1$ ,

$$[H(A, B) + \lambda M]^{-1}(\mathbb{R}) = \mathbb{R}.$$

Hence,  $M$  is  $H(\cdot, \cdot)$ -accretive with respect to  $A$  and  $B$ .

Now, we show that  $M_n \xrightarrow{G} M$ . For any  $(x, y) \in \text{graph}(M)$ , there exists a sequence  $(x_n, y_n) \in \text{graph}(M_n)$ , where let

$$x_n = \left(1 + \frac{1}{n}\right)x,$$

and

$$y_n = M_n(x_n) = \frac{x_n}{2} + \frac{1}{n^2}, \quad \forall n \in \mathbb{N}.$$

Since

$$\lim_n x_n = \lim_n \left[ \left(1 + \frac{1}{n}\right)x \right] = x,$$

we have,

$$x_n \rightarrow x \quad \text{as } n \rightarrow \infty.$$

In addition, by definition of graph, it follows that

$$\lim_n y_n = \lim_n \left( \frac{x_n}{2} + \frac{1}{n^2} \right) = \frac{1}{2}x = M(x) = y.$$

It follows that  $y_n \rightarrow y$  as  $n \rightarrow \infty$  and hence,  $M_n \xrightarrow{G} M$ .

Furthermore, we show that  $J_{M_n, \lambda}^{H(\cdot, \cdot)} \rightarrow J_{M, \lambda}^{H(\cdot, \cdot)}$  as  $M_n \xrightarrow{G} M$ .

Let for  $\lambda = 1$ , the resolvent operators are given by

$$R_{M_n, \lambda}^{H(\cdot, \cdot)}(x) = [H(A, B) + \lambda M_n]^{-1}(x) = 2\sqrt[3]{\left(x - \frac{1}{n^2}\right)},$$

and

$$R_{M, \lambda}^{H(\cdot, \cdot)}(x) = [H(A, B) + \lambda M]^{-1}(x) = 2\sqrt[3]{x},$$

and the generalized Yosida approximation operators are given by

$$J_{M_n, \lambda}^{H(\cdot, \cdot)}(x) = \frac{1}{\lambda} [I - R_{M_n, \lambda}^{H(\cdot, \cdot)}](x) = \left[ x - 2\sqrt[3]{\left(x - \frac{1}{n^2}\right)} \right],$$

and

$$J_{M, \lambda}^{H(\cdot, \cdot)}(x) = \frac{1}{\lambda} [I - R_{M, \lambda}^{H(\cdot, \cdot)}](x) = (x - 2\sqrt[3]{x}).$$

We evaluate

$$\left\| J_{M_n, \lambda}^{H(\cdot, \cdot)} - J_{M, \lambda}^{H(\cdot, \cdot)} \right\| = \left\| \left[ x - 2\sqrt[3]{\left(x - \frac{1}{n^2}\right)} \right] - (x - 2\sqrt[3]{x}) \right\|,$$


which shows that

$$\left\| J_{M_n, \lambda}^{H(\cdot, \cdot)} - J_{M, \lambda}^{H(\cdot, \cdot)} \right\| \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

i.e.,

$$J_{M_n, \lambda}^{H(\cdot, \cdot)} \rightarrow J_{M, \lambda}^{H(\cdot, \cdot)} \quad \text{as } M_n \xrightarrow{G} M.$$

Using the above example, the convergence of generalized Yosida approximation operator  $J_{M_n, \lambda}^{H(\cdot, \cdot)}$  to  $J_{M, \lambda}^{H(\cdot, \cdot)}$  is illustrated in the following figure for  $n = 1, 2, 5, 15$ .



### A Yosida inclusion problem and existence of solution

First, we state a Yosida inclusion problem and its equivalence with a fixed point problem.

Let  $X$  be  $q$ -uniformly smooth Banach space and let  $M : X \rightarrow 2^X$  be  $H(\cdot, \cdot)$ -accretive operator. We consider the following problem.

Find  $x \in X$ , such that

$$0 \in J_{M, \lambda}^{H(\cdot, \cdot)}(x) + M(x), \quad \forall x \in X, \quad \lambda > 0, \tag{10}$$

where  $J_{M, \lambda}^{H(\cdot, \cdot)}$  is the generalized Yosida approximation operator defined by Eq. (2). Problem (10) is called Yosida inclusion problem.

The fixed point formulation of the problem Eq. (10) is as follows:

$$x = R_{M, \lambda}^{H(\cdot, \cdot)} \left[ H(A, B)x - \lambda J_{M, \lambda}^{H(\cdot, \cdot)}(x) \right], \quad \forall x \in X, \quad \lambda > 0. \tag{11}$$

Using the definition of the resolvent operator  $R_{M, \lambda}^{H(\cdot, \cdot)}$  defined by Eq. (1), one can easily obtain the equivalence of Eqs. (10) and (11).

Based on Eq. (11), we construct the following iterative algorithm for solving Yosida inclusion problem Eq. (10).

**Algorithm 1** For any  $x_0 \in X$ , compute the sequence  $\{x_n\} \subset X$  by the following scheme:

$$x_{n+1} = R_{M_n, \lambda}^{H(\cdot, \cdot)} \left[ H(A, B)x_n - \lambda J_{M_n, \lambda}^{H(\cdot, \cdot)}(x_n) \right], \tag{12}$$

where  $\lambda > 0, \quad n = 0, 1, 2, \dots$

If  $J_{M_n, \lambda}^{H(\cdot, \cdot)} = T$ , where  $T : X \rightarrow X$  is a mapping, then the Yosida inclusion problem (10) and Algorithm 1 reduces to the variational inclusion problem (10) and Algorithm 1 of Li and Huang [7], respectively, and note that for

suitable choice of operators in the formulation of (12), one can obtain many existing problems and algorithms in literature.

**Theorem 3** Let  $X$  be a  $q$ -uniformly smooth Banach space and  $A, B : X \rightarrow X$  be the single-valued mappings. Let  $H : X \times X \rightarrow X$  be a single-valued mapping and  $M_n, M : X \rightarrow 2^X$  be the  $H(\cdot, \cdot)$ -accretive operators, such that  $M_n \xrightarrow{G} M$ . Assume that

1.  $H(A, B)$  is  $\alpha$ -strongly accretive with respect to  $A$  and  $\beta$ -relaxed accretive with respect to  $B$  and  $\alpha > \beta$ ;
2.  $H(A, B)$  is  $\gamma_1$ -Lipschitz continuous with respect to  $A$  and  $\gamma_2$ -Lipschitz continuous with respect to  $B$ ;
3.  $(\alpha - \beta) \geq \sqrt[q]{1 + c_q(\gamma_1 + \gamma_2)^q} - q(\alpha - \beta) + \sqrt[q]{1 - q\lambda\theta_2 + c_q\lambda^q\theta_1}$ ;
4.  $(\alpha - \beta) \geq [\gamma_1 + \gamma_2 + \lambda\theta_1]$ .

where  $\theta_1 = \frac{[(\alpha - \beta) + 1]}{\lambda(\alpha - \beta)}$ ,  $\theta_2 = \frac{[(\alpha - \beta) - 1]}{\lambda(\alpha - \beta)}$ ,  $\alpha > \beta$  and  $c_q$  is same as in Lemma 2.1. Then, the Yosida inclusion problem (10) has a unique solution and the iterative sequence  $\{x_n\}$  generated by Algorithm 1 converges strongly to  $x$ .

*Proof* Let the mapping  $F : X \rightarrow X$  be defined by

$$F(x) = R_{M, \lambda}^{H(\cdot, \cdot)} \left[ H(A, B)x - \lambda J_{M, \lambda}^{H(\cdot, \cdot)}(x) \right], \quad \forall x \in X, \quad \lambda > 0.$$

For any  $x, y \in X$  and using Lemma 2, we have

$$\begin{aligned} \|F(x) - F(y)\| &= \left\| R_{M, \lambda}^{H(\cdot, \cdot)} \left[ H(A, B)x - \lambda J_{M, \lambda}^{H(\cdot, \cdot)}(x) \right] - R_{M, \lambda}^{H(\cdot, \cdot)} \left[ H(A, B)y - \lambda J_{M, \lambda}^{H(\cdot, \cdot)}(y) \right] \right\| \\ &\leq \frac{1}{(\alpha - \beta)} \left\| H(A, B)x - \lambda J_{M, \lambda}^{H(\cdot, \cdot)}(x) - H(A, B)y + \lambda J_{M, \lambda}^{H(\cdot, \cdot)}(y) \right\| \\ &= \frac{1}{(\alpha - \beta)} \left\| H(A, B)x - H(A, B)y - (x - y) + (x - y) - \lambda J_{M, \lambda}^{H(\cdot, \cdot)}(x) + \lambda J_{M, \lambda}^{H(\cdot, \cdot)}(y) \right\| \\ &\leq \frac{1}{(\alpha - \beta)} \|H(A, B)x - H(A, B)y - (x - y)\| \\ &\quad + \frac{1}{(\alpha - \beta)} \left\| (x - y) - \lambda \left( J_{M, \lambda}^{H(\cdot, \cdot)}(x) - J_{M, \lambda}^{H(\cdot, \cdot)}(y) \right) \right\|. \end{aligned} \tag{13}$$

Using the same arguments as used in Li and Huang [7], we have

$$\begin{aligned} \|H(A, B)x - H(A, B)y - (x - y)\|^q &\leq [1 + c_q(\gamma_1 + \gamma_2)^q - q(\alpha - \beta)] \|x - y\|^q, \end{aligned}$$

and hence

$$\begin{aligned} \|H(A, B)x - H(A, B)y - (x - y)\| &\leq \sqrt[q]{1 + c_q(\gamma_1 + \gamma_2)^q - q(\alpha - \beta)} \|x - y\|. \end{aligned} \tag{14}$$

Using (1) and (2) of Lemma 4, we obtain

$$\begin{aligned} & \left\| (x - y) - \lambda \left( J_{M,\lambda}^{H(\cdot,\cdot)}(x) - J_{M,\lambda}^{H(\cdot,\cdot)}(y) \right) \right\|^q \\ & \leq \|x - y\|^q - q\lambda \left\langle J_{M,\lambda}^{H(\cdot,\cdot)}(x) - J_{M,\lambda}^{H(\cdot,\cdot)}(y), F_q(x - y) \right\rangle \\ & \quad + c_q \lambda^q \left\| J_{M,\lambda}^{H(\cdot,\cdot)}(x) - J_{M,\lambda}^{H(\cdot,\cdot)}(y) \right\|^q \\ & \leq \|x - y\|^q - q\lambda\theta_2 \|x - y\|^q + c_q \lambda^q \theta_1 \|x - y\|^q \\ & = (1 - q\lambda\theta_2 + c_q \lambda^q \theta_1) \|x - y\|^q, \end{aligned}$$

i.e.,

$$\begin{aligned} & \left\| (x - y) - \lambda \left( J_{M,\lambda}^{H(\cdot,\cdot)}(x) - J_{M,\lambda}^{H(\cdot,\cdot)}(y) \right) \right\| \\ & \leq \sqrt[q]{1 - q\lambda\theta_2 + c_q \lambda^q \theta_1} \|x - y\|^q. \end{aligned} \tag{15}$$

Using Eqs. (14), (15), (13) becomes

$$\begin{aligned} \|F(x) - F(y)\| & \leq \frac{1}{\alpha - \beta} \left[ \sqrt[q]{1 + c_q(\gamma_1 + \gamma_2)^q - q(\alpha - \beta)} \right. \\ & \quad \left. + \sqrt[q]{1 - q\lambda\theta_2 + c_q \lambda^q \theta_1} \right] \|x - y\|, \end{aligned}$$

i.e.,

$$\|F(x) - F(y)\| \leq k \|x - y\|, \tag{16}$$

where

$$\begin{aligned} k & = \frac{1}{\alpha - \beta} \left[ \sqrt[q]{1 + c_q(\gamma_1 + \gamma_2)^q - q(\alpha - \beta)} \right. \\ & \quad \left. + \sqrt[q]{1 - q\lambda\theta_2 + c_q \lambda^q \theta_1} \right]. \end{aligned}$$

By condition (3), it follows that  $0 < k < 1$  and so (16) implies that the mapping  $F$  has a unique fixed point  $x \in X$ . Thus,  $x$  is a unique solution of Yosida inclusion problem (10).

Next, we show that the sequence  $\{x_n\}$  generated by the Algorithm 1 strongly converges to  $x$ .

Using Eqs. (11) and (12), we obtain

$$\begin{aligned} \|x_{n+1} - x\| & = \left\| R_{M,\lambda}^{H(\cdot,\cdot)} \left[ H(A, B)x_n - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) \right] \right. \\ & \quad \left. - R_{M,\lambda}^{H(\cdot,\cdot)} \left[ H(A, B)x - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x) \right] \right\| \\ & = \left\| R_{M,\lambda}^{H(\cdot,\cdot)} \left[ H(Ax_n, Bx_n) - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) \right] \right. \\ & \quad \left. - R_{M,\lambda}^{H(\cdot,\cdot)} \left[ H(Ax_n, Bx_n) - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) \right] \right. \\ & \quad \left. + R_{M,\lambda}^{H(\cdot,\cdot)} \left[ H(Ax_n, Bx_n) - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) \right] \right. \\ & \quad \left. - R_{M,\lambda}^{H(\cdot,\cdot)} \left[ H(Ax, Bx) - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x) \right] \right\| \\ & \leq \left\| R_{M,\lambda}^{H(\cdot,\cdot)} \left[ H(Ax_n, Bx_n) - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) \right] \right. \\ & \quad \left. - R_{M,\lambda}^{H(\cdot,\cdot)} \left[ H(Ax_n, Bx_n) - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) \right] \right\| \\ & \quad + \left\| R_{M,\lambda}^{H(\cdot,\cdot)} \left[ H(Ax_n, Bx_n) - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) \right] \right. \\ & \quad \left. - R_{M,\lambda}^{H(\cdot,\cdot)} \left[ H(Ax, Bx) - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x) \right] \right\| \\ & \leq b_n + \frac{1}{\alpha - \beta} \left\| H(Ax_n, Bx_n) - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) \right. \\ & \quad \left. - \left[ H(Ax, Bx) - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x) \right] \right\|, \end{aligned} \tag{17}$$

where

$$\begin{aligned} b_n & = \left\| R_{M,\lambda}^{H(\cdot,\cdot)} \left[ H(Ax_n, Bx_n) - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) \right] \right. \\ & \quad \left. - R_{M,\lambda}^{H(\cdot,\cdot)} \left[ H(Ax_n, Bx_n) - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) \right] \right\|. \end{aligned}$$

Using Lipschitz continuity of  $H(A, B)$  in both the arguments and Lipschitz continuity of generalized Yosida approximation operator, we obtain

$$\begin{aligned} & \left\| H(Ax_n, Bx_n) - H(Ax, Bx) - \lambda \left[ J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) - J_{M,\lambda}^{H(\cdot,\cdot)}(x) \right] \right\| \\ & \leq \left\| H(Ax_n, Bx_n) - H(Ax_n, Bx) + H(Ax_n, Bx) - H(Ax, Bx) \right. \\ & \quad \left. - \lambda \left[ J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) - J_{M,\lambda}^{H(\cdot,\cdot)}(x) \right] - J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) + J_{M,\lambda}^{H(\cdot,\cdot)}(x) \right\| \\ & \leq \|H(Ax_n, Bx_n) - H(Ax_n, Bx)\| + \|H(Ax_n, Bx) - H(Ax, Bx)\| \\ & \quad + \lambda \left\| J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) - J_{M,\lambda}^{H(\cdot,\cdot)}(x) \right\| + \lambda \left\| J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) - J_{M,\lambda}^{H(\cdot,\cdot)}(x) \right\| \\ & \leq \gamma_2 \|x_n - x\| + \gamma_1 \|x_n - x\| + \lambda c_n + \lambda \theta_1 \|x_n - x\|, \end{aligned} \tag{18}$$

where  $c_n = \left\| J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) - J_{M,\lambda}^{H(\cdot,\cdot)}(x) \right\|$ .

Using Eq. (18), (17) becomes

$$\|x_{n+1} - x\| \leq b_n + \frac{1}{\alpha - \beta} [\gamma_1 + \gamma_2 + \lambda \theta_1] \|x_n - x\| + \lambda c_n,$$

where  $\theta_1 = \frac{[\alpha - \beta + 1]}{\lambda(\alpha - \beta)}$ .

By Theorems 1 and 2, we have

$$\begin{aligned} & R_{M,\lambda}^{H(\cdot,\cdot)} \left[ H(Ax_n, Bx_n) - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) \right] \rightarrow \\ & R_{M,\lambda}^{H(\cdot,\cdot)} \left[ H(Ax_n, Bx_n) - \lambda J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) \right] \end{aligned}$$

and hence,

$$J_{M,\lambda}^{H(\cdot,\cdot)}(x_n) \rightarrow J_{M,\lambda}^{H(\cdot,\cdot)}(x_n).$$

Thus,  $b_n \rightarrow 0$  and  $c_n \rightarrow 0$  as  $n \rightarrow \infty$ . It follows that

$$\|x_{n+1} - x\| \leq P(\theta) \|x_n - x\| + d_n,$$

where  $d_n = b_n + \lambda c_n$ , and  $P(\theta) = \frac{1}{(\alpha - \beta)} [\gamma_1 + \gamma_2 + \lambda \theta_1]$ . By condition (4), we have  $0 < P(\theta) < 1$  and  $d_n \rightarrow 0$  as  $b_n, c_n \rightarrow 0 (n \rightarrow \infty)$ . By Lemma 3, we have

$$\|x_{n+1} - x\| \rightarrow 0.$$

This completes the proof.  $\square$

**Remark 2** If we take  $J_{M,\lambda}^{H(\cdot,\cdot)} = T$ , where  $T : X \rightarrow X$  is a mapping and deleting condition (4) from Theorem 3, we can obtain Theorem 4.1 of Li and Huang [7].

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



## References

- Ahmad, R., Akram, M., Dilshad, M.: Graph Convergence for the  $H(\cdot, \cdot)$ -co-accretive mapping with an application. *Bull. Malays. Math. Sci. Soc.* **38**(4), 1481–1506 (2015)
- Agrawal, R.P., Verma, R.U.: Generalized system of  $(A, \eta)$ -maximal relaxed monotone variational inclusion problems based on generalized hybrid algorithms. *Commun. Nonlinear Sci. Numer. Simul.* **15**(2), 238–251 (2010)
- Cao, H.W.: Yosida approximation equations technique for system of generalized set-valued variational inclusions, *J. Inequal. Appl.* **2013**, 455 (2013)
- Fang, Y.P., Huang, N.J.:  $H$ -monotone operator and resolvent operator technique for variational inclusions. *Appl. Math. Comput.* **145**, 795–803 (2003)
- Ikehata, R., Okozawa, N.: Yosida approximation and Nonlinear Hybrid Equation. *Nonlinear Anal.* **15**(5), 479–495 (1990)
- Iwamiya, T., Okochi, H.: Monotonicity, resolvents and Yosida approximations of operators on Hilbert manifolds. *Nonlinear Anal.* **54**(2), 205–214 (2003)
- Li, X., Huang, N.J.: Graph convergence for the  $H(\cdot, \cdot)$ -accretive operator in Banach spaces with an application. *Appl. Math. Comput.* **217**, 9053–9061 (2011)
- Lan, H.Y.: Generalized Yosida approximations based on relatively  $A$ -maximal  $m$ -relaxed monotonicity frameworks. *Abstr. Appl. Anal.* **2013** (2013) (**article ID 157190**)
- Moudafi, A.: A Duality algorithm for solving general variational inclusions. *Adv. Model. Optim.* **13**(2), 213–220 (2011)
- Penot, J.-P., Ratsimahalo, R.: On the Yosida approximat on of operators. *Proc. R. Soc. Edinb. Math.* **131A**, 945–966 (2001)
- Verma, R.U.: General system of  $A$ -monotone nonlinear variational inclusion problems with applications. *J. Optim. Theory Appl.* **131**(1), 151–157 (2006)
- Verma, R.U.: General nonlinear variational inclusion problems involving  $A$ -monotone mapping. *Appl. Math. Lett.* **19**, 960–963 (2006)
- Xu, H.K.: Inequalities in Banach Spaces with applications. *Nonlinear Anal.* **16**(12), 1127–1138 (1991)
- Zou, Y.Z., Huang, N.J.:  $H(\cdot, \cdot)$ -accretive operator with an application for solving variational inclusions in Banach spaces. *Appl. Math. Comput.* **204**, 809–816 (2008)
- Zou, Y.Z., Huang, N.J.: A new system of Variational Inclusions involving  $(A, \eta)$ -accretive operator in Banach spaces. *Appl. Math. Comput.* **212**, 135–144 (2009)

