Hermite–Hadamard-type inequalities for generalized s -convex functions on real linear fractal set $\mathbb{R}^\alpha$ ($0 < \alpha < 1$)

Huixia Mo¹ · Xin Sui¹

Received: 27 August 2015 / Accepted: 4 February 2016 / Published online: 19 June 2017
© The Author(s) 2017. This article is an open access publication

Abstract In this paper, we establish the Hermite–Hadamard-type inequalities for the generalized s -convex functions in the second sense on real linear fractal set $\mathbb{R}^\alpha$ ($0 < \alpha < 1$).

Keywords Fractal set · Local fractional integral · Hermite–Hadamard-type inequality · Generalized s -convex function

Introduction

The convex function plays an important role in the class mathematical analysis course and other fields. In [1], Hudzik and Maligranda introduced two kinds of s -convex functions in the space of European space $\mathbb{R}$. In addition, many important inequalities are established for the s -convex functions in $\mathbb{R}$. For example, the Hermite–Hadamard's inequality is one of the best known results in the literature, see [2–4] and so on.

In recent years, the fractal theory has received significantly remarkable attention [5]. The calculus on fractal set can lead to better comprehension for the various real-world models from the engineering and science [6].

On the fractal set, Mo et al. [7, 8] introduced the definition of the generalized convex function and established Hermite–Hadamard-type inequality. In [9], the authors

introduced two kinds of generalized s -convex functions on fractal sets $\mathbb{R}^\alpha$ ($0 < \alpha < 1$).

The definitions of the generalized s -convex functions are as follows:

Definition 1.1 [9] Suppose that $\mathbb{R}_+ = [0, \infty)$. If the function $f : \mathbb{R}_+ \rightarrow \mathbb{R}^\alpha$ satisfies the following inequality:

$$f(\lambda_1 u + \lambda_2 v) \leq \lambda_1^{zs} f(u) + \lambda_2^{zs} f(v), \quad (1.1)$$

for all $u, v \in \mathbb{R}_+$ and all $\lambda_1, \lambda_2 \geq 0$ with $\lambda_1^s + \lambda_2^s = 1$ and $0 < s < 1$, then f is said to be a generalized s -convex function in the first sense. We denote this by $f \in GK_s^1$.

Definition 1.2 [9] Suppose that $\mathbb{R}_+ = [0, \infty)$. If the function $f : \mathbb{R}_+ \rightarrow \mathbb{R}^\alpha$ satisfies the following inequality:

$$f(\lambda_1 u + \lambda_2 v) \leq \lambda_1^{zs} f(u) + \lambda_2^{zs} f(v), \quad (1.2)$$

for all $u, v \in \mathbb{R}_+$ and all $\lambda_1, \lambda_2 \geq 0$ with $\lambda_1 + \lambda_2 = 1$ and $0 < s < 1$, then f is said to be a generalized s -convex function in the second sense. We denote this by $f \in GK_s^2$.

Note that the generalized s -convex function in both sense is generalized convex function [9] for $s = 1$.

Inspired by [2, 3, 8], in this paper, we will establish the Hermite–Hadamard-type inequalities for the generalized s -convex functions.

Preliminaries

Now, let us review the operations with real line number on fractal space. In addition, we will use the Gao–Yang–Kang's idea to describe the definitions of the local fractional derivative and local fractional integral [10–14].

Let a^α , b^α and c^α belong to the set $\mathbb{R}^\alpha$ ($0 < \alpha < 1$) of real line numbers, then

The research is supported by the NNSF of China (11601035, 11471050).

✉ Huixia Mo
huixiamo@bupt.edu.cn

¹ School of Science, Beijing University of Posts and Telecommunications, Beijing 100876, China

1. $a^\alpha b^\alpha$ and $a^\alpha + b^\alpha$ belong to the set $\mathbb{R}^\alpha$;
2. $a^\alpha + b^\alpha = (a + b)^\alpha = b^\alpha + a^\alpha = (b + a)^\alpha$;
3. $a^\alpha + (b^\alpha + c^\alpha) = (a^\alpha + b^\alpha) + c^\alpha$;
4. $a^\alpha b^\alpha = (ab)^\alpha = b^\alpha a^\alpha = (ba)^\alpha$;
5. $a^\alpha (b^\alpha c^\alpha) = (a^\alpha b^\alpha) c^\alpha$;
6. $a^\alpha (b^\alpha + c^\alpha) = a^\alpha b^\alpha + a^\alpha c^\alpha$;
7. $0^\alpha + a^\alpha = a^\alpha + 0^\alpha = a^\alpha$ and $1^\alpha \cdot a^\alpha = a^\alpha \cdot 1^\alpha = a^\alpha$.

Definition 2.1 ([10]) If the function $f : [a, b] \rightarrow \mathbb{R}^\alpha$ satisfies the inequality

$$|f(x) - f(y)| < c|x - y|^\alpha, \quad (x, y \in [a, b]),$$

for $c > 0$ and $\alpha(0 < \alpha \leq 1)$, then f is called a Hölder continuous function. In this case, we think that f is in the space $C_\alpha[a, b]$.

Definition 2.2 [10] Let $\Delta^\alpha(f(x) - f(x_0)) \cong \Gamma(1 + \alpha)(f(x) - f(x_0))$. Then, the local fractional derivative of f of order α at $x = x_0$ is defined by

$$f^{(\alpha)}(x_0) = \frac{d^\alpha f(x)}{dx^\alpha} \Big|_{x=x_0} = \lim_{x \rightarrow x_0} \frac{\Delta^\alpha(f(x) - f(x_0))}{(x - x_0)^\alpha}.$$

If there exists $f^{((k+1)\alpha)}(x) = \overbrace{D_x^\alpha \dots D_x^\alpha}^{k+1 \text{ times}} f(x)$ for any $x \in I \subseteq \mathbb{R}$, then we denoted $f \in D_{(k+1)\alpha}(I)$, where $k = 0, 1, 2, \dots$

Definition 2.3 [10] For $f \in C_\alpha[a, b]$, the local fractional integral of the function f is defined by

$$\begin{aligned} & {}_a I_b^{(\alpha)} f(x) \\ &= \frac{1}{\Gamma(1 + \alpha)} \int_a^b f(t) (dt)^\alpha \\ &= \frac{1}{\Gamma(1 + \alpha)} \lim_{\Delta t \rightarrow 0} \sum_{j=0}^N f(t_j) (\Delta t_j)^\alpha, \end{aligned}$$

where $\Delta t_j = t_{j+1} - t_j$, $\Delta t = \max\{\Delta t_1, \Delta t_2, \Delta t_j, \dots\}$ and $[t_j, t_{j+1}]$, $j = 0, \dots, N-1$, $t_0 = a$, $t_N = b$, is a partition of the interval $[a, b]$.

Lemma 2.1 [10] Let $f \in C_\alpha[g(a), g(b)]$ and $g \in C_1[a, b]$, then

$$g(a) I_{g(b)}^{(\alpha)} f(x) = {}_a I_b^{(\alpha)} f(g(s)) [g'(s)]^\alpha.$$

Lemma 2.2 [10]

1. Let $f(x) = g^{(\alpha)}(x) \in C_\alpha[a, b]$, then we have ${}_a I_b^{(\alpha)} f(x) = g(b) - g(a)$.
2. Let $f(x), g(x) \in D_\alpha[a, b]$ and $f^{(\alpha)}(x), g^{(\alpha)}(x) \in C_\alpha[a, b]$, then we have

$${}_a I_b^{(\alpha)} f(x) g^{(\alpha)}(x) = f(x) g(x) \Big|_a^b - {}_a I_b^{(\alpha)} f^{(\alpha)}(x) g(x).$$

Lemma 2.3 [10]

$$\frac{d^\alpha x^{k\alpha}}{dx^\alpha} = \frac{\Gamma(1 + k\alpha)}{\Gamma(1 + (k-1)\alpha)} x^{(k-1)\alpha}.$$

From the above formula and Lemma 2.2, we have

$$\frac{1}{\Gamma(1 + \alpha)} \int_a^b x^{k\alpha} (dx)^\alpha = \frac{\Gamma(1 + k\alpha)}{\Gamma(1 + (k+1)\alpha)} (b^{(k+1)\alpha} - a^{(k+1)\alpha}), k \in \mathbb{R}.$$

Lemma 2.4 [10] (Generalized Hölder's inequality) Let $f, g \in C_\alpha[a, b]$ and $p, q > 1$ with $1/p + 1/q = 1$. Then, it follows that

$$\begin{aligned} & \frac{1}{\Gamma(1 + \alpha)} \int_a^b |f(x)g(x)| (dx)^\alpha \leq \left(\frac{1}{\Gamma(1 + \alpha)} \int_a^b |f(x)|^p (dx)^\alpha \right)^{1/p} \\ & \times \left(\frac{1}{\Gamma(1 + \alpha)} \int_a^b |g(x)|^q (dx)^\alpha \right)^{1/q}. \end{aligned}$$

Main results

Theorem 3.1 Let $f : \mathbb{R}_+ \rightarrow \mathbb{R}^\alpha$ be a generalized s -convex function in the second sense for $0 < s < 1$ and $a, b \in [0, \infty)$ with $a < b$. Then, for $f \in C_\alpha[a, b]$, the following inequalities hold:

$$\frac{2^{(s-1)\alpha}}{\Gamma(1 + \alpha)} f\left(\frac{a+b}{2}\right) \leq \frac{{}_a I_b^{(\alpha)} f(x)}{(b-a)^\alpha} \leq \frac{\Gamma(1 + s\alpha)}{\Gamma(1 + (s+1)\alpha)} (f(a) + f(b)). \quad (3.1)$$

Proof Let $x = a + b - t$. Then, from Lemma 2.1, we have $\frac{a+b}{2} I_b^{(\alpha)} f(x) = {}_a I_{\frac{a+b}{2}}^{(\alpha)} f(a + b - t)$.

Since f is a generalized s -convex function in the second sense, then

$$\begin{aligned} {}_a I_b^{(\alpha)} f(x) &= {}_a I_{\frac{a+b}{2}}^{(\alpha)} [f(x) + f(a + b - x)] \\ &\geq 2^{s\alpha} {}_a I_{\frac{a+b}{2}}^{(\alpha)} f\left(\frac{a+b}{2}\right) \\ &= \frac{2^{(s-1)\alpha}}{\Gamma(1 + \alpha)} (b-a)^\alpha f\left(\frac{a+b}{2}\right). \end{aligned}$$

In the other hand, let $x = b - (b-a)t$, $0 \leq t \leq 1$, then we get

$$\begin{aligned} {}_a I_b^{(\alpha)} f(x) &= (b-a)^\alpha {}_0 I_1^{(\alpha)} f[ta + (1-t)b] \\ &\leq (b-a)^\alpha {}_0 I_1^{(\alpha)} [t^{s\alpha} f(a) + (1-t)^{s\alpha} f(b)] \\ &= (b-a)^\alpha [f(a) {}_0 I_1^{(\alpha)} t^{s\alpha} + f(b) {}_0 I_1^{(\alpha)} (1-t)^{s\alpha}]. \end{aligned}$$

From Lemma 2.3, it is easy to see that

$${}_0 I_1^{(\alpha)} t^{s\alpha} = \frac{\Gamma(1 + s\alpha)}{\Gamma(1 + (s+1)\alpha)},$$

and

$${}_0I_1^{(\alpha)}(1-t)^{\alpha s} = \frac{\Gamma(1+s\alpha)}{\Gamma(1+(s+1)\alpha)}.$$

Therefore

$${}_aI_b^{(\alpha)}f(x) \leq (b-a)^{\alpha} \frac{\Gamma(1+s\alpha)}{\Gamma(1+(s+1)\alpha)} (f(a) + f(b)).$$

Combining the above estimates, we obtain

$$\begin{aligned} \frac{2^{(s-1)\alpha}}{\Gamma(1+\alpha)} f\left(\frac{a+b}{2}\right) &\leq \frac{{}_aI_b^{(\alpha)}f(x)}{(b-a)^{\alpha}} \\ &\leq \frac{\Gamma(1+s\alpha)}{\Gamma(1+(s+1)\alpha)} (f(a) + f(b)). \end{aligned}$$

□

Theorem 3.2 Let $I \subset \mathbb{R}$ be an interval, and I^0 be the interior of I . Suppose that $f : I \rightarrow \mathbb{R}^{\alpha}$ is a differentiable function on I^0 such that $f^{(\alpha)} \in C_{\alpha}[a, b]$, where $a, b \in I^0$ with $a < b$. If $|f^{(\alpha)}|^q$ is a generalized s -convex function in the second sense on $[a, b]$ for some fixed $s \in (0, 1)$ and $q \geq 1$, then

$$\begin{aligned} &\left| \frac{f(a) + f(b)}{2^{\alpha}} - \frac{\Gamma(1+\alpha)}{(b-a)^{\alpha}} {}_aI_b^{(\alpha)}f(x) \right| \\ &\leq \frac{(b-a)^{\alpha}}{2^{\alpha}} \left(\frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \right)^{\frac{q-1}{q}} \\ &\quad \left[\frac{\Gamma(1+s\alpha)}{\Gamma(1+(s+1)\alpha)} + \frac{\Gamma(1+\alpha)\Gamma(1+s\alpha)}{\Gamma(1+(s+2)\alpha)} \left(\left(\frac{1}{2} \right)^{\alpha s} - 2^{\alpha} \right) \right]^{\frac{1}{q}} \\ &\quad \times \left[|f^{(\alpha)}(a)|^q + |f^{(\alpha)}(b)|^q \right]^{\frac{1}{q}}. \end{aligned}$$

To show Theorem 3.2 is right, we need the following Lemma.

Lemma 3.1 ([8]) Let $f : I \rightarrow \mathbb{R}^{\alpha}$, $I \subset [0, \infty)$. If $f \in D_{\alpha}(I^0)$ and $f^{(\alpha)} \in C_{\alpha}[a, b]$ for $a, b \in I^0$ with $a < b$, then the following equality holds:

$$\begin{aligned} &\frac{f(a) + f(b)}{2^{\alpha}} - \frac{\Gamma(1+\alpha)}{(b-a)^{\alpha}} {}_aI_b^{(\alpha)}f(x) \\ &= \frac{(b-a)^{\alpha}}{2^{\alpha}} \frac{1}{\Gamma(1+\alpha)} \int_0^1 (1-2t)^{\alpha} f^{(\alpha)}(ta + (1-t)b) (dt)^{\alpha}. \end{aligned}$$

Now, let us give the proof of Theorem 3.2.

Proof From Lemma 3.1, it is obvious that

$$\begin{aligned} &\left| \frac{f(a) + f(b)}{2^{\alpha}} - \frac{\Gamma(1+\alpha)}{(b-a)^{\alpha}} {}_aI_b^{(\alpha)}f(x) \right| \\ &\leq \frac{(b-a)^{\alpha}}{2^{\alpha}} \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^{\alpha} |f^{(\alpha)}(ta + (1-t)b)| (dt)^{\alpha}. \end{aligned} \quad (3.2)$$

Let us estimate

$$\frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^{\alpha} |f^{(\alpha)}(ta + (1-t)b)| (dt)^{\alpha},$$

for $q = 1$ and $q > 1$.

Case I. $q = 1$.

Since $|f^{(\alpha)}|$ is generalized s -convex on $[a, b]$ in the second sense, we can know that for any $t \in [0, 1]$

$$|f^{(\alpha)}(ta + (1-t)b)| \leq t^{\alpha s} |f^{(\alpha)}(a)| + (1-t)^{\alpha s} |f^{(\alpha)}(b)|.$$

Then, we have

$$\begin{aligned} &\frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^{\alpha} |f^{(\alpha)}(ta + (1-t)b)| (dt)^{\alpha} \\ &\leq \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^{\alpha} \left[t^{\alpha s} |f^{(\alpha)}(a)| \right. \\ &\quad \left. + (1-t)^{\alpha s} |f^{(\alpha)}(b)| \right] (dt)^{\alpha} \\ &= \left[|f^{(\alpha)}(a)| \frac{1}{\Gamma(1+\alpha)} \int_0^1 t^{\alpha s} |1-2t|^{\alpha} (dt)^{\alpha} + |f^{(\alpha)}(b)| \right. \\ &\quad \left. \frac{1}{\Gamma(1+\alpha)} \int_0^1 (1-t)^{\alpha s} |1-2t|^{\alpha} (dt)^{\alpha} \right]. \end{aligned} \quad (3.3)$$

From Lemmas 2.2 and 2.3, it is easy to see that

$$\begin{aligned} &\frac{1}{\Gamma(1+\alpha)} \int_0^1 t^{\alpha s} |1-2t|^{\alpha} (dt)^{\alpha} \\ &= \frac{1}{\Gamma(1+\alpha)} \left[\int_0^{\frac{1}{2}} t^{\alpha s} (1-2t)^{\alpha} (dt)^{\alpha} + \int_{\frac{1}{2}}^1 t^{\alpha s} (2t-1)^{\alpha} (dt)^{\alpha} \right] \\ &= \left[\frac{\Gamma(1+s\alpha)}{\Gamma(1+(s+1)\alpha)} + \frac{\Gamma(1+\alpha)\Gamma(1+s\alpha)}{\Gamma(1+(s+2)\alpha)} \left(\left(\frac{1}{2} \right)^{\alpha s} - 2^{\alpha} \right) \right]. \end{aligned} \quad (3.4)$$

In addition, let $1-t = x$, then by Lemma 2.1 and (3.4), we have

$$\begin{aligned} &\frac{1}{\Gamma(1+\alpha)} \int_0^1 (1-t)^{\alpha s} |1-2t|^{\alpha} (dt)^{\alpha} \\ &= \frac{1}{\Gamma(1+\alpha)} \int_0^1 x^{\alpha s} |1-2x|^{\alpha} (dx)^{\alpha} \\ &= \left[\frac{\Gamma(1+s\alpha)}{\Gamma(1+(s+1)\alpha)} + \frac{\Gamma(1+\alpha)\Gamma(1+s\alpha)}{\Gamma(1+(s+2)\alpha)} \left(\left(\frac{1}{2} \right)^{\alpha s} - 2^{\alpha} \right) \right]. \end{aligned} \quad (3.5)$$

Thus, substituting (3.4) and (3.5) into (3.3), we have

$$\begin{aligned} & \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha |f^{(\alpha)}(ta+(1-t)b)| (dt)^\alpha \\ & \leq \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \left[\frac{\Gamma(1+s\alpha)}{\Gamma(1+(s+1)\alpha)} + \frac{\Gamma(1+\alpha)\Gamma(1+s\alpha)}{\Gamma(1+(s+2)\alpha)} \right. \\ & \quad \times \left. \left(\left(\frac{1}{2} \right)^{\alpha s} - 2^\alpha \right) \right] \times \left[|f^{(\alpha)}(a)|^q + |f^{(\alpha)}(b)|^q \right]. \end{aligned} \quad (3.6)$$

Thus, from (3.2), we obtain

$$\begin{aligned} & \left| \frac{f(a)+f(b)}{2^\alpha} - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^{(\alpha)} f(x) \right| \\ & \leq \frac{(b-a)^\alpha}{2^\alpha} \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \left[\frac{\Gamma(1+s\alpha)}{\Gamma(1+(s+1)\alpha)} \right. \\ & \quad + \frac{\Gamma(1+\alpha)\Gamma(1+s\alpha)}{\Gamma(1+(s+2)\alpha)} \left(\left(\frac{1}{2} \right)^{\alpha s} - 2^\alpha \right) \left. \right] \\ & \quad \times \left[|f^{(\alpha)}(a)|^q + |f^{(\alpha)}(b)|^q \right]. \end{aligned}$$

Case II. $q > 1$.

Using the generalized Hölder's inequality (Lemma 2.4), we obtain

$$\begin{aligned} & \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha |f^{(\alpha)}(ta+(1-t)b)| (dt)^\alpha \\ & = \frac{1}{\Gamma(1+\alpha)} \\ & \quad \int_0^1 |1-2t|^{\alpha \frac{q-1}{q}} |1-2t|^{\alpha \frac{1}{q}} |f^{(\alpha)}(ta+(1-t)b)| (dt)^\alpha \\ & \leq \left(\frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha (dt)^\alpha \right)^{\frac{q-1}{q}} \\ & \quad \times \left(\frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha |f^{(\alpha)}(ta+(1-t)b)|^q (dt)^\alpha \right)^{\frac{1}{q}}. \end{aligned} \quad (3.7)$$

It is obvious that

$$\begin{aligned} & \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha (dt)^\alpha \\ & = \frac{1}{\Gamma(1+\alpha)} \int_0^{\frac{1}{2}} (1-2t)^\alpha (dt)^\alpha \\ & \quad + \frac{1}{\Gamma(1+\alpha)} \int_{\frac{1}{2}}^1 (2t-1)^\alpha (dt)^\alpha = \frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)}. \end{aligned} \quad (3.8)$$

Moreover, since $|f^{(\alpha)}|^q$ is generalized s convex in the second sense on $[a, b]$, then

$$\begin{aligned} & \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha |f^{(\alpha)}(ta+(1-t)b)|^q (dt)^\alpha \\ & \leq \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha \\ & \quad \left(t^{\alpha s} |f^{(\alpha)}(a)|^q + (1-t)^{\alpha s} |f^{(\alpha)}(b)|^q \right) (dt)^\alpha \\ & = |f^{(\alpha)}(a)|^q \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha t^{\alpha s} (dt)^\alpha \\ & \quad + |f^{(\alpha)}(b)|^q \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha (1-t)^{\alpha s} (dt)^\alpha. \end{aligned}$$

From (3.3) and (3.4), it is easy to see that

$$\begin{aligned} & \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha t^{\alpha s} (dt)^\alpha \\ & = \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha (1-t)^{\alpha s} (dt)^\alpha \\ & = \left[\frac{\Gamma(1+s\alpha)}{\Gamma(1+(s+1)\alpha)} + \frac{\Gamma(1+\alpha)\Gamma(1+s\alpha)}{\Gamma(1+(s+2)\alpha)} \left(\left(\frac{1}{2} \right)^{\alpha s} - 2^\alpha \right) \right]. \end{aligned}$$

Therefore

$$\begin{aligned} & \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha |f^{(\alpha)}(ta+(1-t)b)|^q (dt)^\alpha \\ & = \left[\frac{\Gamma(1+s\alpha)}{\Gamma(1+(s+1)\alpha)} + \frac{\Gamma(1+\alpha)\Gamma(1+s\alpha)}{\Gamma(1+(s+2)\alpha)} \left(\left(\frac{1}{2} \right)^{\alpha s} - 2^\alpha \right) \right]^{\frac{1}{q}} \\ & \quad \left[|f^{(\alpha)}(a)|^q + |f^{(\alpha)}(b)|^q \right]^{\frac{1}{q}}. \end{aligned} \quad (3.9)$$

Thus, substituting (3.8) and (3.9) into (3.7), we have

$$\begin{aligned} & \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha |f^{(\alpha)}(ta+(1-t)b)| (dt)^\alpha \\ & \leq \left(\frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \right)^{\frac{q-1}{q}} \\ & \quad \left[\frac{\Gamma(1+s\alpha)}{\Gamma(1+(s+1)\alpha)} + \frac{\Gamma(1+\alpha)\Gamma(1+s\alpha)}{\Gamma(1+(s+2)\alpha)} \left(\left(\frac{1}{2} \right)^{\alpha s} - 2^\alpha \right) \right]^{\frac{1}{q}} \\ & \quad \left[|f^{(\alpha)}(a)|^q + |f^{(\alpha)}(b)|^q \right]^{\frac{1}{q}}. \end{aligned}$$

Therefore, from (3.2), it follows that

$$\begin{aligned}
& \left| \frac{f(a)+f(b)}{2^\alpha} - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^{(\alpha)} f(x) \right| \\
& \leq \frac{(b-a)^\alpha}{2^\alpha} \left(\frac{\Gamma(1+\alpha)}{\Gamma(1+2\alpha)} \right)^{\frac{q-1}{q}} \\
& \left[\frac{\Gamma(1+s\alpha)}{\Gamma(1+(s+1)\alpha)} + \frac{\Gamma(1+\alpha)\Gamma(1+s\alpha)}{\Gamma(1+(s+2)\alpha)} \left(\left(\frac{1}{2} \right)^{\alpha s} - 2^\alpha \right) \right]^{\frac{1}{q}} \\
& \quad \times \left[|f^{(\alpha)}(a)|^q + |f^{(\alpha)}(b)|^q \right]^{\frac{1}{q}}.
\end{aligned}$$

Thus, we complete the proof of Theorem 3.2. $\square$

Theorem 3.3 Suppose that $f: I \rightarrow \mathbb{R}^\alpha$, $I \subset [0, \infty)$ is a differentiable function on I^0 , such that $f^{(\alpha)} \in C_\alpha[a, b]$, where $a, b \in I$ with $a < b$. If $|f^{(\alpha)}|^q$ is a generalized s -convex function in the second sense on $[a, b]$ for some fixed $s \in (0, 1)$ and $q > 1$, then

$$\begin{aligned}
& \left| \frac{f(a)+f(b)}{2^\alpha} - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^{(\alpha)} f(x) \right| \\
& \leq \frac{(b-a)^\alpha}{2^\alpha} \left[\frac{\Gamma(1+\frac{q}{q-1}\alpha)}{2^\alpha \Gamma(1+\frac{2q-1}{q-1}\alpha)} \right]^{\frac{q-1}{q}} \\
& \left(\frac{\Gamma(1+s\alpha)}{2^\alpha \Gamma(1+(s+1)\alpha)} \right)^{\frac{1}{q}} \\
& \quad \times \left[\left(|f^{(\alpha)}(a)|^q + \left| f^{(\alpha)}\left(\frac{a+b}{2}\right) \right|^q \right)^{\frac{1}{q}} \right. \\
& \quad \left. + \left(\left| f^{(\alpha)}\left(\frac{a+b}{2}\right) \right|^q + |f^{(\alpha)}(b)|^q \right)^{\frac{1}{q}} \right].
\end{aligned}$$

Proof From Lemma 3.1, we have

$$\begin{aligned}
& \left| \frac{f(a)+f(b)}{2^\alpha} - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^{(\alpha)} f(x) \right| \\
& \leq \frac{(b-a)^\alpha}{2^\alpha} \frac{1}{\Gamma(1+\alpha)} \int_0^1 |1-2t|^\alpha |f^{(\alpha)}(ta+(1-t)b)| (dt)^\alpha \\
& \leq \frac{(b-a)^\alpha}{2^\alpha} \left[\frac{1}{\Gamma(1+\alpha)} \int_0^{\frac{1}{2}} (1-2t) |f^{(\alpha)}(ta+(1-t)b)| (dt)^\alpha \right. \\
& \quad \left. + \frac{1}{\Gamma(1+\alpha)} \int_{\frac{1}{2}}^1 (2t-1) |f^{(\alpha)}(ta+(1-t)b)| (dt)^\alpha \right]. \quad (3.10)
\end{aligned}$$

Let us estimate

$$\frac{1}{\Gamma(1+\alpha)} \int_0^{\frac{1}{2}} (1-2t) |f^{(\alpha)}(ta+(1-t)b)| (dt)^\alpha$$

and

$$\frac{1}{\Gamma(1+\alpha)} \int_{\frac{1}{2}}^1 (2t-1) |f^{(\alpha)}(ta+(1-t)b)| (dt)^\alpha,$$

respectively.

Using the generalized Hölder's inequality (Lemma 2.4), we obtain

$$\begin{aligned}
& \frac{1}{\Gamma(1+\alpha)} \int_0^{\frac{1}{2}} (1-2t) |f^{(\alpha)}(ta+(1-t)b)| (dt)^\alpha \\
& \leq \left(\frac{1}{\Gamma(1+\alpha)} \int_0^{\frac{1}{2}} (1-2t)^{\frac{q}{q-1}} (dt)^\alpha \right)^{\frac{q-1}{q}} \\
& \quad \left(\frac{1}{\Gamma(1+\alpha)} \int_0^{\frac{1}{2}} |f^{(\alpha)}(ta+(1-t)b)|^{q\alpha} (dt)^\alpha \right)^{\frac{1}{q}}. \quad (3.11)
\end{aligned}$$

It is easy to see that

$$\begin{aligned}
& \frac{1}{\Gamma(1+\alpha)} \int_0^{\frac{1}{2}} (1-2t)^{\frac{q}{q-1}} (dt)^\alpha = \\
& \frac{1}{\Gamma(1+\alpha)} \int_{\frac{1}{2}}^1 (1-2t)^{\frac{q}{q-1}} (dt)^\alpha = \frac{\Gamma(1+\frac{q}{q-1}\alpha)}{2^\alpha \Gamma(1+\frac{2q-1}{q-1}\alpha)}. \quad (3.12)
\end{aligned}$$

Let $|f^{(\alpha)}(ta+(1-t)b)|^q = U(t)$. It is easy to see that $U(t)$ is a generalized s -convex function in the second sense. Thus, from the right-hand side of (3.1), it follows that

$$\begin{aligned}
& \frac{1}{\Gamma(1+\alpha)} \int_0^{\frac{1}{2}} |f^{(\alpha)}(ta+(1-t)b)|^q (dt)^\alpha \\
& = \frac{1}{\Gamma(1+\alpha)} \int_0^{\frac{1}{2}} U(t) (dt)^\alpha \\
& \leq \left(\frac{1}{2} - 0 \right)^\alpha \frac{\Gamma(1+s\alpha)}{\Gamma(1+(s+1)\alpha)} \left(U(0) + U\left(\frac{1}{2}\right) \right) \\
& = \frac{\Gamma(1+s\alpha)}{2^\alpha \Gamma(1+(s+1)\alpha)} \left(\left| f^{(\alpha)}\left(\frac{a+b}{2}\right) \right|^q + |f^{(\alpha)}(b)|^q \right). \quad (3.13)
\end{aligned}$$

Thus, substituting (3.12) and (3.13) into (3.11), we get

$$\begin{aligned}
& \frac{1}{\Gamma(1+\alpha)} \int_0^{\frac{1}{2}} (1-2t) |f^{(\alpha)}(ta+(1-t)b)| (dt)^\alpha \\
& \leq \left(\frac{\Gamma(1+\frac{q}{q-1}\alpha)}{2^\alpha \Gamma(1+\frac{2q-1}{q-1}\alpha)} \right)^{\frac{q-1}{q}} \left(\frac{\Gamma(1+s\alpha)}{2^\alpha \Gamma(1+(s+1)\alpha)} \right)^{\frac{1}{q}} \\
& \quad \left(|f^{(\alpha)}(a)|^q + \left| f^{(\alpha)}\left(\frac{a+b}{2}\right) \right|^q \right)^{\frac{1}{q}}. \quad (3.14)
\end{aligned}$$

Moreover

$$\begin{aligned} & \frac{1}{\Gamma(1+\alpha)} \int_{\frac{1}{2}}^1 (2t-1)^{\frac{\alpha}{q-1}} (dt)^\alpha \\ &= \frac{1}{\Gamma(1+\alpha)} \int_{\frac{1}{2}}^1 (1-2t)^{\frac{\alpha}{q-1}} (dt)^\alpha = \frac{\Gamma(1+\frac{q}{q-1}\alpha)}{2^\alpha \Gamma(1+\frac{2q-1}{q-1}\alpha)}. \end{aligned}$$

In addition, similar to the estimate of (3.13), we have

$$\begin{aligned} & \frac{1}{\Gamma(1+\alpha)} \int_{\frac{1}{2}}^1 |f^{(\alpha)}(ta + (1-t)b)|^q (dt)^\alpha \\ & \leq \frac{\Gamma(1+s\alpha)}{2^\alpha \Gamma(1+(s+1)\alpha)} \left(\left| f^{(\alpha)}\left(\frac{a+b}{2}\right) \right|^q + |f^{(\alpha)}(b)|^q \right). \end{aligned}$$

Therefore, it is analogous to the estimate of (3.11), we have

$$\begin{aligned} & \frac{1}{\Gamma(1+\alpha)} \int_0^{\frac{1}{2}} (1-2t) |f^{(\alpha)}(ta + (1-t)b)| (dt)^\alpha \\ & \leq \left(\frac{1}{\Gamma(1+\alpha)} \int_{\frac{1}{2}}^1 (2t-1)^{\frac{\alpha}{q-1}} (dt)^\alpha \right)^{\frac{q-1}{q}} \\ & \quad \left(\frac{1}{\Gamma(1+\alpha)} \int_{\frac{1}{2}}^1 |f^{(\alpha)}(ta + (1-t)b)|^{q\alpha} (dt)^\alpha \right)^{\frac{1}{q}} \\ & \leq \left(\frac{\Gamma(1+\frac{q}{q-1}\alpha)}{2^\alpha \Gamma(1+\frac{2q-1}{q-1}\alpha)} \right)^{\frac{q-1}{q}} \left(\frac{\Gamma(1+s\alpha)}{2^\alpha \Gamma(1+(s+1)\alpha)} \right)^{\frac{1}{q}} \\ & \quad \left(\left| f^{(\alpha)}\left(\frac{a+b}{2}\right) \right|^q + |f^{(\alpha)}(b)|^q \right)^{\frac{1}{q}}. \end{aligned} \quad (3.15)$$

Thus, combining (3.10), (3.14), and (3.15), we obtain

$$\begin{aligned} & \left| \frac{f(a)+f(b)}{2^\alpha} - \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^{(\alpha)} f(x) \right| \\ & \leq \frac{(b-a)^\alpha}{2^\alpha} \left[\frac{\Gamma(1+\frac{q}{q-1}\alpha)}{2^\alpha \Gamma(1+\frac{2q-1}{q-1}\alpha)} \right]^{\frac{q-1}{q}} \left(\frac{\Gamma(1+s\alpha)}{2^\alpha \Gamma(1+(s+1)\alpha)} \right)^{\frac{1}{q}} \\ & \quad \times \left[\left(|f^{(\alpha)}(a)|^q + \left| f^{(\alpha)}\left(\frac{a+b}{2}\right) \right|^q \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\left| f^{(\alpha)}\left(\frac{a+b}{2}\right) \right|^q + |f^{(\alpha)}(b)|^q \right)^{\frac{1}{q}} \right]. \end{aligned}$$

□

Therefore, we complete the proof of Theorem 3.3.

Acknowledgements The authors would like to express their gratitude to the Editors and referees for some very valuable suggestion.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Hudzik, H., Maligranda, L.: Some remarks on s -convex functions. *Aequationes Math.* **48**(1), 100–111 (1994)
- Kirmaci, U.S., Bakula, M.K., Özdemir, M.E., Pečarić, J.: Hadamard-type inequalities for s -convex functions. *Appl. Math. Comput.* **193**(1), 26–35 (2007)
- Avci, M., Kavurmaci, H., Ozdemir, M.E.: New inequalities of Hermite-Hadamard type via s -convex functions in the second sense with applications. *Appl. Math. Comput.* **217**, 5171–5176 (2011)
- Shuang, Y., Yin, H.P., Qi, F.: Hermite-Hadamard type integral inequalities for geometric-arithmetically s -convex functions. *Analysis*. **33**(2), 197–208 (2013)
- Kilbas, A.A., Srivastava, H.M., Trujillo, J.J.: *Theory and Applications of Fractional Differential Equations*. North-Holland Mathematical Studies. Elsevier (North-Holland) Science Publishers, Amsterdam, London and New York (2006)
- Gao, F., Zhong, W.P., Shen, X.M.: Applications of Yang-Fourier transform to local fractional equations with local fractional derivative and local fractional integral. *Adv. Mater. Res.* **461**, 306–310 (2012)
- Mo, H., Sui, X., Yu, D.: Generalized convex functions and some inequalities on fractal sets. *Abstr. Appl. Anal.* **2014**, Article ID 636751 (2014)
- Mo, H.: Generalized Hermite-Hadamard Type Inequalities Involving Local Fractional Integrals. [arxiv: 1410.1062](https://arxiv.org/abs/1410.1062)
- Mo, H., Sui, X.: Generalized s -convex function on fractal sets. *Abstr. Appl. Anal.* **2014**, Article ID 254737 (2014)
- Yang, X.J.: *Advanced Local Fractional Calculus and Its Applications*. World Science Publisher, New York, NY, USA (2012)
- Wie, W., Srivastava, H.M., Zhang, Y., Wang, L., Shen, P., Zhang, J.: A local fractional integral inequality on fractal space analogous to andersons inequality. *Abstr. Appl. Anal.* **2014**, Article ID 797561 (2014)
- Yang, X.J., Baleanu, D., Machado, J.A.T.: Mathematical aspects of heisenberg uncertainty principle within local fractional fourier analysis. *Bound. Value Probl.* **2013**, 131–146 (2013)
- Zhao, Y., Cheng, D.F., Yang, X.J.: Approximation solutions for local fractional Schrödinger Equation in the on-dimensional cantor system. *Adv. Math. Phys.* **2013**, Article ID 291386 (2013)
- Yang, A.M., Chen, Z.S., Srivastava, H.M., Yang, X.J.: Application of the local fractional series expansion method and the variational iteration method to the Helmholtz equation involving local fractional derivative operators. *Abstr. Appl. Anal.* **2013**, Article ID 259125 (2013)