



A simple and flexible modification of Grünwald–Letnikov fractional derivative in image processing

H. Jalalinejad¹ · A. Tavakoli² · F. Zarmehi³

Received: 24 May 2018 / Accepted: 24 August 2018
© The Author(s) 2018

Abstract

In image processing, edge detection and image enhancement can make use of fractional differentiation operators, especially the Grünwald–Letnikov derivative. In this paper, we present a modified Grünwald–Letnikov derivative to enhance more and detect better the edges of an image. Our proposed fractional derivative is very flexible and can be easily performed. We present some examples to justify our suggested approach.

Keywords Grünwald–Letnikov fractional derivative · Image enhancement · Edge detection

Introduction

The fractional differential equation has a long history of more than 300 years. Many mathematicians such as Euler, Laplace, Abel, Liouville, Riemann, Grünwald, Letnikov and Riez have worked in this field of mathematics. In 1974, first conference on fractional calculus and its application was held [1]. In Podlubny [2] wrote a book that provides the basic theory of fractional differentiation, equations and methods of their solution. Models based on partial differential equations and calculus of variations are also generalized for fractional derivatives. For instance, fractional-order partial differential equation-based formulation are applicable for multi-scale nonlocal contrast enhancement with texture preserving [3] and iterative learning control with high-order internal models [4]. In image processing,

fractional calculus is exploited in image denoising using the diffusion equation [5–8] and in image segmentation with active contours using the fractional derivative within energy functional [9]. Mathieu et al. [10] applied the fractional differentiation for edge detection. Also, they discussed on the texture enhancement of multi-scale fractional mask.

Zhang et al. [11] have proposed fractional differential mask based on the definition of Riemann–Liouville. For fractional order of 1 to 2, they enhanced the texture and edges in multi-scale by controlling the fractional order. For denoising an image, Pu et al. applied fractional calculus based on the definition of Riemann–Liouville [12]. Also, Gao et al. in [13] applied an improved fractional differential operator based on a piecewise quaternion for image enhancement. Furthermore, in [14], the generalized fractional image denoising algorithm based on Srivastava–owa fractional differential operator is introduced for image denoising. The Grünwald–Letnikov derivative is also used for image enhancement in [15, 16]. In Gao et al. [17] by development of the real fractional derivative and its applications in the signal processing extended the quaternion fractional differential (QFD) based on Grünwald–Letnikov and applied it to edge detection of color image. He et al. in [18] proposed a model based on the Grünwald–Letnikov fractional differential operator that improves denoising operator mask. The total coefficient of this mask is not equal to zero, which means that its response value is not zero in flat areas of the image. The total coefficient of

✉ A. Tavakoli
a.tavakoli@umz.ac.ir

H. Jalalinejad
hoda.jalali@iauk.ac.ir

F. Zarmehi
f.zarmehi@mail.vru.ac.ir

¹ Department of Mathematics, Kerman Branch, Islamic Azad University, Kerman, Iran

² Mathematics Department, University of Mazandaran, Babolsar, Iran

³ Mathematics Department, Vali-e-Asr University of Rafsanjan, P. O. Box 518, Rafsanjan, Iran

this mask is not equal to zero, which means that its response value is not zero in flat areas of the image. In 2017, Jalab et al. proposed a new contrast enhancement technique for medical images based on image entropy. Their method enhances edges accurately while preserving smooth textures [19]. We aim to redefine the Grünwald–Letnikov derivative, in order to better show the rate of changes of the derivative in image processing. In this paper, we highlight the defects of Grünwald–Letnikov derivative in image processing and based on them, we present a new definition of Grünwald–Letnikov derivative that is very flexible.

Preliminaries

In this section, we introduce some basic concepts which are essential to our discussions in the next sections. Let us now recall that the n th-order derivative of function f is defined by:

$$f^{(n)}(x) = \frac{d^n f}{dx^n} = \lim_{h \rightarrow 0} \frac{1}{h^n} \sum_{r=0}^n (-1)^r \binom{n}{r} f(x - rh).$$

Accordingly, the Grünwald–Letnikov fractional derivative for one variable function f is defined as follows [2]:

$$D_{G-L}^\alpha f(x) = \lim_{h \rightarrow 0} \frac{1}{h^\alpha} \sum_{r=0}^{\lfloor \frac{x-a}{h} \rfloor} (-1)^r \binom{\alpha}{r} f(x - rh),$$

where

$$\binom{\alpha}{r} = \frac{\Gamma(\alpha + 1)}{\Gamma(r + 1)\Gamma(\alpha - r + 1)},$$

and Γ is the gamma function.

Usually, an image can be defined as a two-dimensional function $f(x, y)$ where x and y are spatial coordinates. The value of $f(x, y)$ is called the color intensity of image at point (x, y) . In the field of image processing, the Grünwald–Letnikov derivative in two dimensions in the x -direction can be defined as follows [15, 20]:

$$D_{G-L}^\alpha f_x(x, y) = f(x, y) - \alpha f(x - 1, y) + \frac{\alpha(\alpha - 1)}{2} f(x - 2, y). \tag{1}$$

Similarly, the Grünwald–Letnikov derivative is defined in y -direction. Hence, the Grünwald–Letnikov fractional derivative can be defined by

$$D_{G-L}^\alpha f(x, y) = \sqrt{(D_{G-L}^\alpha f_x(x, y))^2 + (D_{G-L}^\alpha f_y(x, y))^2}, \tag{2}$$

or

$$D_{G-L}^\alpha f(x, y) \approx |D_{G-L}^\alpha f_x(x, y)| + |D_{G-L}^\alpha f_y(x, y)|. \tag{3}$$

The similarities and the differences of regular derivative and Grünwald–Letnikov fractional derivative can be summarized as follows:

1. For the region of an image I whose color intensities are the same, the gradient of I is zero inside of the region(not on the edge points), but it is nonzero for Grünwald–Letnikov derivative. Furthermore, the more the intensity is closer to white (255), the larger the Grünwald–Letnikov derivative.
2. In edge pixels that gradient is positive (negative), the Grünwald–Letnikov derivative is also positive (negative). However, the (absolute) value of Grünwald–Letnikov derivative is usually larger than that of regular gradient.

By presenting some examples, we show that the definition of Grünwald–Letnikov derivative will arise some disorderliness in the application of derivative in image processing. In the following examples for simplicity, we consider $0 < \alpha \leq 1$ and study the Grünwald–Letnikov derivative in x -direction.

Example 1 Let $f(x - 1, y) = f(x - 2, y) = f(x, y) = 250$. By (1) we get

$$\begin{aligned} D_{G-L}^\alpha f_x(x, y) &= 250 - \alpha 250 + \frac{\alpha(\alpha - 1)}{2} 250 \\ &= (1 - \alpha)(2 - \alpha)125, \end{aligned}$$

that implies

$$0 < D_{G-L}^\alpha f_x(x, y) < 250.$$

In the special case $\alpha = 1/2$, we have $D_{G-L}^\alpha f_x(x, y) = 93.75$.

Example 2 Let $f(x - 2, y) = f(x - 1, y) = f(x, y) = 1$. We have

$$D_{G-L}^\alpha f_x(x, y) = 1 - 1\alpha + \frac{\alpha(\alpha - 1)}{2} = (1 - \alpha)(2 - \alpha)/2.$$

Again, for $0 < \alpha \leq 1$, we have

$$0 \leq D_{G-L}^\alpha f_x(x, y) < 1.$$

In Examples 1 and 2, the value of $f(x, y)$ is constant in the x -neighborhood of $f(x, y)$, hence, we expect no change or a few change of (fractional) derivative of f in x -direction. However, we see the value of $D_{G-L}^\alpha f_x(x, y)$ severely depends on the intensity of f rather than the difference of f and their x -neighborhoods.

Example 3 Let $f(x - 2, y) = f(x - 1, y) = 1$ and $f(x, y) = 250$. By computing the Grünwald–Letnikov derivative, we obtain

$$D_{G-L}^{\alpha} f_x(x, y) = 250 - \alpha + \frac{\alpha(\alpha - 1)}{2} = 250 - \alpha(3 - \alpha)/2.$$

For $0 < \alpha \leq 1$, we have

$$249 \leq D_{G-L}^{\alpha} f_x(x, y) < 250,$$

and for $\alpha = 1/2$, we have $D_{G-L}^{\alpha} f_x(x, y) = 249.3750$.

Example 4 Let $f(x - 2, y) = f(x - 1, y) = 250$ and $f(x, y) = 1$. We get

$$D_{G-L}^{\alpha} f_x(x, y) = 1 - 250\alpha + \frac{\alpha(\alpha - 1)250}{2} = 1 - 125\alpha(3 - \alpha),$$

that for $0 < \alpha \leq 1$ we have

$$-249 \leq D_{G-L}^{\alpha} f_x(x, y) < 1.$$

For $\alpha = 1/2$, we have $D_{G-L}^{\alpha} f_x(x, y) = -155.2500$.

In Examples 3 and 4, we observe that the difference of $f(x, y)$ and its x-neighborhoods are the same; however, the Grünwald–Letnikov derivatives of $f(x, y)$ in x-direction are very different. The above examples show that Grünwald–Letnikov derivative is sensitive to the intensity of the pixels rather than the difference of the intensities.

According to these examples, the definition of Grünwald–Letnikov derivative should be modified in order to better represent the rate of changes of the derivative.

Modified Grünwald–Letnikov derivative

In this section, we express a modified definition of Grünwald–Letnikov derivative. To this end, we first take

$$M(x, y) = \frac{1}{s^n} \min\{f(x, y), f(x - 1, y), f(x - 2, y)\},$$

where $s \geq 255$ is an integer number and $0 \leq n \leq 1$ is a real number. The equation of the line passing through of two points $(0, M(x, y))$ and $(s, 0)$ is

$$Y(x, y) = M(x, y) \left(\frac{s - X(x, y)}{s} \right).$$

By substituting

$$X(x, y) = |f(x, y) - \alpha f(x - 1, y) + \frac{\alpha(\alpha - 1)}{2} f(x - 2, y)|,$$

in the above equation, the value of $Y(x, y)$ is obtained. Now, we define the modified Grünwald–Letnikov derivative in x-direction as follows:

$${}_m D_{G-L}^{\alpha} f_x(x, y) = \frac{f(x, y) - \alpha f(x - 1, y) + \frac{\alpha(\alpha - 1)}{2} f(x - 2, y)}{Y(x, y) + 1}. \tag{4}$$

In Eq. 4, the value 1 is added to $Y(x, y)$ to avoid of vanishing the denominator. The coefficient $\frac{1}{Y(x, y)+1}$ is thought of as modifier parameters of Grünwald–Letnikov derivative. Moreover, it is important to note that for $0 < n \leq 1$,

$$\lim_{s \rightarrow \infty} Y(x, y) = 0.$$

This yields the following lemma;

Lemma 1 *The modified Grünwald–Letnikov derivative defined by (4) will be the same Grünwald–Letnikov as defined by (1), if $s \rightarrow +\infty$.*

By (4), we get

$${}_m D_{G-L}^{\alpha} f_x(x, y) = \frac{s^{n+1} A}{\theta(s - |A|) + s^{n+1}}, \tag{5}$$

where $\theta = \min\{f(x, y), f(x - 1, y), f(x - 2, y)\}$ and $A = D_{G-L}^{\alpha} f_x(x, y)$.

Furthermore, by (5), it is seen that if $s = |A|$, then the regular and modified Grünwald–Letnikov derivatives will be the same. By considering the parameters s and n , we have two degree of freedom. In fact, the modified Grünwald–Letnikov derivative generally has a behavior between the regular derivative and Grünwald–Letnikov fractional derivative. Analogously, one can define the modified Grünwald–Letnikov derivative in y-direction. Hence, the modified Grünwald–Letnikov fractional derivative can be defined by

$${}_m D_{G-L}^{\alpha} f(x, y) = \sqrt{({}_m D_{G-L}^{\alpha} f_x(x, y))^2 + ({}_m D_{G-L}^{\alpha} f_y(x, y))^2}, \tag{6}$$

or

$${}_m D_{G-L}^{\alpha} f(x, y) \approx |{}_m D_{G-L}^{\alpha} f_x(x, y)| + |{}_m D_{G-L}^{\alpha} f_y(x, y)|. \tag{7}$$

Now, we compute the modified Grünwald–Letnikov derivative for the preceding examples. By (5), for Example 1, we have

$$0 \leq {}_m D_{G-L}^{\alpha} f_x(x, y) < \frac{250s^{n+1}}{250(s - 250) + s^{n+1}},$$

in which $0 < \alpha \leq 1$. The special case $\alpha = 1/2, s = 255$ and $n = 1$ yields ${}_m D_{G-L}^{\alpha} f_x(x, y) = 57.8725$. For Example 2,

$$0 \leq {}_m D_{G-L}^{\alpha} f_x(x, y) < \frac{s^{n+1}}{(s - 1) + s^{n+1}} < 1,$$

in which $0 < \alpha \leq 1$. For Example 3,

$$0 < \frac{s^{n+1}}{(s - 249) + s^{n+1}} \leq {}_m D_{G-L}^{\alpha} f_x(x, y) < \frac{s^{n+1}}{(s - 250) + s^{n+1}} < 1,$$

in which $0 < \alpha \leq 1$. Finally, for Example (4), we have

$$\frac{-249s^{n+1}}{(s-249) + s^{n+1}} \leq {}_m D_{G-L}^\alpha f_x(x, y) < \frac{s^{n+1}}{(s-1) + s^{n+1}} < 1,$$

in which $0 < \alpha \leq 1$. The special case $\alpha = 1/2, s = 255$ and $n = 1$ yields

$${}_m D_{G-L}^\alpha f_x(x, y) = -155.01,$$

that is approximately equal to the value of usual Grünwald–Letnikov derivative.

We observe that the multiplier $\frac{1}{Y(x,y)+1}$ in the modified Grünwald–Letnikov derivative moderates the value of the derivative.

Numerical examples

In this section, we aim to demonstrate that the modified Grünwald–Letnikov fractional derivative can be efficiently applied for edge detection and image enhancement. We, moreover, present a comparison between the modified and original Grünwald–Letnikov derivatives for two prototype image.

Example 5 (Edge detection). Consider Fig. 1a as an original image. Figure 1b shows its Grünwald–Letnikov derivative defined by (3) and Fig. 1c shows its modified Grünwald–Letnikov derivative defined by (7). In both Fig. 1b, c, we put $\alpha = 0.5$. Also, for modified Grünwald–Letnikov derivative, $s = 255$ and $n = 0.5$ is selected. As it is seen the modified Grünwald–Letnikov derivative shows only the edges of the main figure while Grünwald–Letnikov derivative shows the whole of figure with low




Fig. 1 a Is an original image; b shows its Grünwald–Letnikov derivative and c shows the modified Grünwald–Letnikov derivative



Fig. 2 The original image of an infant

intensity. Based on Lemma 1, as s tends to infinity, the Grünwald–Letnikov derivative and its modified will be the same.

Example 6 (Image enhancement). Figure 2 shows a gray-scale image of an infant. Figures 3 and 4 show the enhanced images of Fig. 2 by Grünwald–Letnikov derivative and modified Grünwald–Letnikov derivative with $\alpha = 0.2, 0.4, 0.6$ and $\alpha = 0.8$, respectively. We considered $s = 255$ and $n = 0.5$ for enhancing by modified Grünwald–Letnikov derivative. As it is seen, the modified Grünwald–




Fig. 3 The Grünwald–Letnikov derivative of Fig. 2 with different values of α




Fig. 4 The modified Grünwald–Letnikov derivative of Fig. 2 with different values of α

Letnikov derivative gives a better quality in comparison with the usual Grünwald–Letnikov derivative.

Example 7 Figure 5a shows an original image of Lena, and Fig. 5b shows a regular derivative of it. It is computed as $\sqrt{(\frac{\partial u}{\partial x})^2 + (\frac{\partial u}{\partial y})^2}$ where u is the image of Lena. Figures 6 and 7 show the effect of Grünwald–Letnikov derivative and its modified for $\alpha = 0.2, 0.4, 0.6$ and $\alpha = 0.8$, respectively. For modified $G - L$ derivative, we considered $s = 255$ and $n = 0.5$. It is clear that the modified $G - L$ derivatives tends to regular $G - L$ derivatives, as s tends to infinity.



Fig. 5 a original image of “Lena” and it’s regular derivative



Fig. 6 The Grünwald–Letnikov derivative of image “Lena” with different values of α




Fig. 7 The modified Grünwald–Letnikov derivative of image “Lena” with different values of α

Conclusion

In order to better show the rate of change of derivative in image processing, we need to redefine the Grünwald–Letnikov fractional derivative. We highlight the defects of the

Grünwald–Letnikov derivative in image processing, next, we present a new definition of Grünwald–Letnikov fractional derivative that is very flexible. The proposed modified Grünwald–Letnikov can be efficiently employed in different areas of image processing such as image enhancement, edge detection and medical diagnostic.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Ross, B. (ed.): *Fractional Calculus and Its Applications*, Lecture Notes in Mathematics, vol. 457. Springer, Berlin (1975)
- Podlubny, I.: *Fractional Differential Equations*. Academic Press, New York (1999)
- Pu, Y.F., Siarry, P., Chatterjee, A., Wang, Z.N., Yi, Z., Liu, Y., Zhou, J., Wang, Y.: A fractional-order variational framework for retinex: fractional-order partial differential equation-based formulation for multi-scale nonlocal contrast enhancement with texture preserving. *IEEE Trans. Image Process.* **27**(3), 1214–1229 (2018)
- Liu, Sh, Wang, J.: Analysis of iterative learning control with high-order internal models for fractional differential equations. *J. Vib. Control* **24**(6), 1145–1161 (2018)
- Abrirami, A., Prakesh, P., Thangavel, K.: Fractional diffusion equation-based image denoising model using CN-GL scheme. *Int. J. Comput. Math.* **95**(6–7), 1222–1239 (2018)
- Bai, J., Feng, X.-C.: Fractional-order anisotropic diffusion for image denoising. *IEEE Trans. Image Process.* **16**(10), 2492–2502 (2007)
- Verma, A.K., Saini, B.S.: Forward-backward processing technique for image denoising using FDZP 2D filter. *J. Appl. Res. Technol.* **15**, 538–592 (2018)
- Zhang, W., Li, J., Yang, Y.: A fractional diffusion-wave equation with non-local regularization for image denoising. *Signal Process.* **103**, 6–15 (2014)
- Ren, Z.: Adaptive active contour model driven by fractional order fitting energy. *Signal Process.* **117**, 138–150 (2015)
- Mathieu, B., Melchior, P., Oustaloup, A., Ceyral, Ch.: Fractional differentiation for edge detection. *Signal Process.* **83**(11), 2421–2432 (2003)
- Zhang, Y., Pu, Y., Zhou, J.: Construction of fractional differential masks based on Riemann–Liouville definition. *J. Comput. Inf. Syst.* **6**(10), 3191–3199 (2010)
- Hu, J., Pu, Y., Zhou, J.: A novel image denoising algorithm based on Riemann–Liouville definition. *J. Comput.* **6**(7), 1332–1338 (2011)
- Gao, C.B., Zhou, J.L., Zheng, X.Q., Lang, F.N.: Image enhancement based on improved fractional differentiation. *J. Comput. Inf. Syst.* **7**(1), 257–264 (2011)
- Jalab, H.A., Ibrahim, R.W.: Fractional masks based on generalized fractional differential operator for image denoising, world academy of science, engineering and technology. *Int. J. Comput. Inf. Sci. Eng.* **7**(2), 124–129 (2013)
- Pu, Y.F., Zhou, J.L., Yuan, X.: Fractional differential mask: a fractional differential-based approach for multiscale texture enhancement. *IEEE Trans. Image Process.* **19**(2), 491–511 (2010)
- Pu, Y.: Fractional calculus approach to texture of digital image. In: *IEEE Proceeding of the 8th International Conference on Signal Processing*, pp. 1002–1006 (2006)
- Gao, C.B., Zhou, J.L., Hu, J.R., Lang, F.N.: Edge detection of colour image based on quaternion fractional differential. *IET Image Process.* **5**(3), 261–272 (2011)
- He, N., Wang, J.B., Zhang, L.L., Lu, K.: An improved fractional-order differentiation model for image denoising. *Signal Process.* **112**, 180–188 (2015)
- Jalab, H.A., Ibrahim, R.W., Jalab, D.H., Jalab, A.A., Hasan, A.M.: Medical image enhancement based on statistical distributions in fractional calculus. In: *Computing Conference* (2017)
- Huading, J., Pu, Y.: Fractional calculus method for enhancing digital image of bank slip. *Proc. Congr. Image Signal Process.* **3**, 326–330 (2008)
- Gonzalez, R.C., Woods, R.E.: *Digital Image Processing*, 3rd edn. Prentice Hall, New Jersey (2007)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.