GABA_A Receptor Genetics and Clinical Pharmacology

Paul A. Davies

Published online: 10 January 2014
© Springer Science + Business Media New York 2014

Abstract GABA_A receptors (GABA_ARs) are Cl[−]-selective ligand-gated ion channels that mediate both fast synaptic inhibition and persistent tonic inhibition in the CNS. The receptor is composed of subunits that affect biophysical and pharmacological properties of the channel. A growing number of studies have examined copy number variations, mutations, and polymorphisms in genes encoding GABA_AR subunits and found that changes in subunit function and expression are linked to a variety of neurological disorders. GABA_ARs are targets for a wide range of pharmaceutical agents including benzodiazepines, barbiturates, general anesthetics, and neurosteroids. It remains to be fully determined how genetic variations in GABA_A subunits affect clinical pharmacology. Conversely, there is growing interest in the use of GABAergic drugs to treat disorders with an etiology arising from genetic variations in GABA_A subunits.

Keywords GABA · Benzodiazepines · Neurosteroids · General anesthetics · Phasic inhibition · Tonic inhibition

Introduction

The GABA_A receptor (GABA_AR) belongs to a superfamily of ligand-gated ion channels which includes the nicotinic acetylcholine (nACh) receptor, glycine receptor, serotonin type III (5-HT₃) receptor, and the zinc activated channel [1]. Members of this superfamily are pentameric heterooligomers, the subunits of which have a common structure

encompassing a large extracellular N-terminal domain that contains the neurotransmitter binding pocket, four transmembrane domains (TM1–4), and a major intracellular domain between TM3 and TM4 [1]. The five subunits form a ring-like structure with a central, ion-conducting pore with TM2 lining the pore.

The 19 subunits of the GABA_AR family fall into several subunit classes, with each subunit having different temporal and spatial expression throughout the CNS. The subunit composition of GABA_ARs can alter the biophysical and pharmacological characteristics of the receptor. The genes for the subunits form chromosomal clusters: chromosome 4p12 (α2, α4, β1, and γ1); chromosome 5q31–q35 (α1, α6, β2, γ2, and π); chromosome 6q13–q16.3 (ρ1, and ρ2); chromosome 15q11.2–q12 (α5, β3, and γ3); and chromosome Xq28 (α3, ε, and θ). The genes for ρ3, and δ are on chromosomes 3q11.2 and 1p36.3, respectively. The clustering of the subunit genes is believed to help coordinate gene expression [2].

Benzodiazepine-sensitive receptor subtypes are composed of α1–3, βn, and γn subunits [3–5]. These types of GABA_ARs are typically located at the synapse and are activated by brief (1–2 ms) pulses of GABA from presynaptic GABA vesicle release. The released GABA is rapidly removed by GABA transporters (GAT1–3) located in the membranes of neurons and glia, thus limiting GABA's actions in the synapse. The GABA-mediated current is transient as a result of this rapid rise and fall in synaptic GABA levels, and inhibition is termed phasic inhibition. Extrasynaptic GABA_ARs are commonly composed of α4–6, βn, and either γ2 or δ subunits, although in some regions of the brain other subunit combinations, including heteromeric αβ receptors, maybe found [6, 7]. Little is known about the synaptic or extrasynaptic location of γ1 or γ3 subunits. These extrasynaptic GABA_ARs are activated by ambient GABA to

P. A. Davies (✉)
Department of Neuroscience, Tufts University School of
Medicine, 136 Harrison Ave, Boston, MA 02111, USA
e-mail: paul.davies@tufts.edu

produce a persistent or tonic current (discussed by Houston et al. [8]). Extrasynaptic receptors are characterized by a high GABA potency, enabling them to detect the low ambient GABA concentration in the extracellular space [6, 7, 9–11]. Although the amplitude of tonic current is smaller than that of synaptic current, the persistent nature of tonic current enables greater charge transfer over time. This tonic inhibition significantly affects neuronal and circuit excitability [12]. Ambient GABA originates from several sources, including synaptic spillover, release from glia, or release from neurogliaform cells [13, 14]. In addition to activation of GABA_ARs from ambient GABA, some GABA_ARs have a higher probability to transition from a closed to an open state in the absence of GABA. A recent study demonstrated spontaneously (constitutively) open channels in dentate gyrus granule cells that contribute to tonic current [15]. An interesting complication of constitutively open channels is the low efficacy of block with bicuculline or gabazine [15, 16], antagonists typically used to measure tonic current amplitude, and so the constitutively open channel component would not be detected. However, the Cys-loop receptor channel blocker picrotoxin does block constitutively open channels and would be more appropriate for measurement of total tonic current from GABA-mediated channel activity and constitutively open GABA_A channel activity.

The ion-conducting pore of GABA_ARs is selective for chloride anions (Cl[−]), which flow down their electrochemical gradient. In adult mammalian neurons this flow tends to be inward, hence hyperpolarizing the neuron, taking the membrane potential further away from its threshold potential. The low intracellular Cl[−] concentration in an adult neuron is created by efficient potassium chloride cotransporters (KCC-2). A common misconception is that KCC-2 function and surface expression are static in adult neurons, and hence, GABA_A-mediated responses are always hyperpolarizing. Recent results have revealed that neuronal excitation can rapidly alter the phosphorylated state of KCC-2 and hence the rate of ion transport and the cell surface stability of KCC-2, leading to a depolarizing GABA_A-mediated response [17, 18]. Hence, KCC-2 function can be plastic, depending on the activity of a circuit, and the altered Cl[−] homeostasis would have a further effect on neuronal excitability. In several neurological disorders, for example epilepsy, ischemia, and chronic pain, disruption of KCC-2 function has been described which would lead to a perturbed Cl[−] homeostasis and GABA-mediated inhibition [19].

Genetics

GABA_ARs, as mentioned above, are critical for setting the inhibitory tone and for affecting the precise timing of

neuronal firing in networks. It is of no surprise that abnormalities in the expression and function of GABA_ARs will have a pathological phenotype [20]. Many copy-number variations (CNV), mutations, and polymorphisms in genes encoding GABA_AR subunits have been described. The biophysical and pharmacological properties of GABA_ARs are substantially affected by the type of subunit which is expressed in the receptor. Any alterations in the expression of subunits with CNVs and changes in subunit structure as a result of mutations and polymorphisms could affect inhibitory signaling and pharmacological responses, complicating therapy and responses to drugs. These changes can also have far-reaching consequences on other proteins; the physiological state of the neuron could be changed such that other signal pathways, downstream of the Cl[−] flux, change the expression of other genes [21].

Both phasic and tonic inhibition are sensitive to changes and have been described for several disease states. Alterations of the efficacy of tonic inhibition have been strongly implicated in premenstrual disorder, post-partum depression, post-partum seizures, catalepsy, and temporal lobe epilepsy [22–24]. Deficits in tonic inhibition are also found in Fragile X syndrome, suggesting that such inhibition is important for cognitive function [25, 26]. The changes to tonic inhibition in these conditions can be brought about by several factors (on their own or in combination), for example changes in functional receptors, changes in the concentration of agonist, or changes in concentration of neurosteroids that greatly affect tonic inhibition.

In recent years it has emerged that de-novo mutations are crucially involved in epilepsies and neurodevelopmental disorders including autism, intellectual disability, and schizophrenia. Mutations in genes encoding for GABA_AR subunits are a common occurrence in these disorders. A recent study found a de-novo mutation in the gene for the GABA_AR β3 subunit that had significant occurrence as a single-gene cause of epileptic encephalopathies [27]. The study also observed de-novo mutations in the gene for the GABA_AR α1 subunit. Neurodevelopmental disorders occurred among patients with a CNV in chromosome 4q12-13 resulting in the duplication of the genes for GABA_A subunits α2, α4, β1, and γ1 [28]. Similarly, maternal duplications of chromosome 15q11-13, which includes genes for the α5, β3, and γ3 subunits, have been linked to autism spectrum disorders [29].

Idiopathic epilepsy syndromes make up ~50 % of epilepsies and several GABA_A subunit mutations (usually inherited but also de-novo mutations) have been associated with these epilepsies (reviewed in depth by Macdonald et al. [30]). A mutation in the N-terminal of the β3 subunit (G32R) is associated with childhood absence epilepsy [31]. The mutation leads to an increase in the glycosylation of the β3 subunit, which alters the subunit assembly of

mature GABA_ARs and reduces the gating of the receptor to an open state. For subunit assembly, the mutation results in a strong salt bridge forming between $\beta 3$ subunits. Interestingly, the authors suggested that the mutation would result in a change from $\alpha\beta\gamma$ to more receptors composed of binary $\alpha 1\beta 3$ and homomeric $\beta 3$ subunits. The resulting cells expressing these receptors would be likely to change their electrical activity and pharmacological properties, as homomeric $\beta 3$ receptors are not activated by GABA but can be directly activated by neurosteroids and propofol [32].

Expression of GABA_A subunits may also be affected by mutations in other genes. Fragile X mental retardation protein (FMRP) is an RNA binding protein responsible for transport and translational regulation of several proteins [33]. Fragile X syndrome occurs with a CGG repeat mutation in the *Fmr1* gene that encodes FMRP [34], leading to reduced expression of FMRP in Fragile X and other autism spectrum disorders [35]. One target of FMRP is the GABA_A δ subunit mRNA [36]. With the *Fmr1* gene mutation, reduced δ subunit expression and reduced tonic inhibition result in an imbalance of the inhibitory-to-excitatory ratio. Interestingly, FMRP is also downregulated in schizophrenia, bipolar disorder, and major depression [37, 38], neurological conditions in which changes in GABA_AR expression and function are known to occur [20, 37, 39].

Clinical Pharmacology

Because GABA is the major inhibitory neurotransmitter in the adult CNS, GABA_ARs are crucially important in the excitability of neurons. Many clinically relevant compounds have been shown to interact with GABA_ARs and in so doing potentiate the GABA-evoked Cl[−] current. Such compounds include benzodiazepines [40], the anesthetic barbiturates [41], and propofol [42]. Other relevant agents that positively modulate the receptor include the anesthetic synthetic steroid, alphaxalone [43], and the progesterone and deoxycorticosterone metabolites, allopregnanolone and THDOC [44]. These compounds are used clinically as anxiolytics, anticonvulsants, and sedative and/or hypnotic and/or anesthetic agents.

Benzodiazepines

Benzodiazepines are among the most widely used pharmaceutical agents. The high frequency of clinical use is because of their anxiolytic, anticonvulsant, amnestic, muscle relaxant, and sedative–hypnotic properties [3, 45, 46]. Unfortunately, benzodiazepines also cause unwanted

side effects, for example excessive (or unwanted) sedation, tolerance, dependence, and abuse. The benzodiazepine binding site on GABA_ARs is well documented. Benzodiazepine-sensitive GABA_ARs are formed from an α subunit ($\alpha 1$, $\alpha 2$, $\alpha 3$, or $\alpha 5$) with a β subunit and a γ subunit in 2:2:1 stoichiometry. The benzodiazepine binding site is at the subunit interface between the α and γ subunits [47].

Whereas early benzodiazepines were not selective for any particular α subunit, newer benzodiazepine compounds have been produced that do have selectivity for the α subunit, providing insight into the subunit specificity of benzodiazepine behavioral profiles [3]. In addition, transgenic mice that have had their benzodiazepine-sensitive α subunit mutated into a benzodiazepine-insensitive one have provided the unique ability to test the behavioral properties of benzodiazepines. As a result of using these transgenic techniques it is now known that benzodiazepine-enhancing $\alpha 1$ subunits containing GABA_ARs mediate sedative effects whereas anxiolysis and myorelaxant actions are mediated through the $\alpha 2$ and $\alpha 3$ -containing GABA_ARs. Benzodiazepines' ability to cause amnesia is mediated through GABA_ARs containing the $\alpha 5$ subunit [48, 49].

Benzodiazepines such as diazepam and lorazepam are the first treatment for status epilepticus (SE). Refractory SE occurs when SE continues despite benzodiazepine treatment; hypnotic agents, for example volatile anesthetics, barbiturates, and propofol, are then used [19, 50]. It has been suggested that internalized synaptic GABA_ARs cause this resistance to benzodiazepine during SE, but recently we have demonstrated, by use of an in-vitro model, that benzodiazepine resistance can occur because of impaired KCC-2 function leading to accumulation of intracellular Cl[−] and depolarizing GABA_AR-mediated responses. Interesting even though GABA_AR-mediated responses were depolarizing, propofol was still able to inhibit action potentials because it predominantly enhanced the tonic current, electrically shunting membrane conductance [51•].

General Anesthetics

General anesthetics have widely heterogeneous structures yet during the early 1990s it was demonstrated that most anesthetics at clinically relevant concentrations enhance the effects of GABA on GABA_ARs. At higher concentrations, some anesthetics, particularly the i.v. anesthetics propofol, etomidate, and the barbiturates, could directly activate GABA_ARs in the absence of GABA. This direct activation by anesthetics distinguished them from benzodiazepines which are not GABA_Amimetic.

After the early observations of GABA_AR modulation by general anesthetics, much effort was devoted to investigation of which subunits, and in particular which residues,

were important to anesthetic properties. Site-directed mutagenesis studies produced several residues of interest, particularly in the TM regions of the α and β subunits, for both volatile and i.v. anesthetics [52]. The mutagenesis studies naturally led to the generation of transgenic animals to test the in-vivo effects of these residues.

After the discovery that a residue in the $\beta 2$ and $\beta 3$ subunits was important for modulation of GABA-evoked currents by i.v. anesthetics, transgenic mice were generated carrying $\beta 2(N265S)$ and $\beta 3(N265 M)$ mutations that render the GABA_AR carrying them insensitive to the enhancing actions of etomidate and propofol [53, 54]. Investigation of the anesthetic endpoints of etomidate and propofol in these transgenic mice revealed that loss of consciousness can be mediated purely through $\beta 3$ -containing GABA_ARs, whereas the $\beta 2$ subunit mediates the sedative properties of anesthetics.

Mutated receptor subunits have provided us with much information, but one problem with this technique is the mutation itself can cause conformational changes that alter the interaction with the anesthetic, or prevent the conformational change needed for the allosteric enhancement when the anesthetic has bound to the receptor. Photo-affinity labeling uses analogues of general anesthetics that photo-react when exposed to light of a specific wavelength and, if the anesthetic has already bound to the receptor, will label the residue in contact with the anesthetic. By using this method, azietomidate photo-affinity labeling identified a methionine in TM1 of the $\alpha 1$ subunit (M236) and a methionine in TM3 of the $\beta 3$ subunit (M286) [55]. Use of homology models revealed $\alpha 1$ -M236 and $\beta 3$ -M286 face each other, thus forming a binding site between the two subunits. Recently, a propofol analogue was used to photo-label a site in the $\beta 3$ subunit of both $\alpha 1\beta 3$ heteromeric and $\beta 3$ homomeric receptors. Histidine 267 lies near the extracellular end of TM2 [56]. Another recent study examined a photoreactive barbiturate which labeled $\alpha 1$ subunit TM3 at A296 and Y294, $\gamma 2$ subunit TM3 at S301, and $\beta 3$ subunit TM1 at methionine 227. These sites are located on the extracellular side of the channel at subunit interfaces [57]. It seems, therefore, that the structurally diverse anesthetics bind at different sites within the channel to allosterically modulate the channel.

Neurosteroids

Neurosteroids are synthesized de novo in the brain from cholesterol or from steroid hormone precursors. Increasing the concentration of neurosteroids in the CNS causes anxiolysis, sedation, anticonvulsant action, and anesthesia and reduces depressive-like behavior [58]. Dysregulation of neurosteroid signaling is associated with neurological

and psychiatric disorders which include traumatic brain injury, post-traumatic disorder, premenstrual dysphoric disorder, panic disorder, depression, schizophrenia, and bipolar disorder [59]. Neurosteroids exert most of their actions by allosterically enhancing the activity of GABA on GABA_ARs. At low nanomolar concentrations they potentiate submaximum GABA-dependent currents; at a single channel level neurosteroids increase the open time and frequency of opening of the channel [60]. At micromolar concentrations, they directly activate GABA_ARs in the absence of GABA [61, 62]. This direct activation property has largely been ignored because larger, non-physiological, concentrations are needed to see this direct gating effect. However, it has been reported that neurosteroids can slowly directly activate GABA_ARs at high nanomolar concentrations via accumulation in the membrane where they are able to access an activation site [63].

Neurosteroids have been shown to bind GABA_ARs at an allosteric site distinct from that of GABA, benzodiazepines, or barbiturates [46, 64, 65]. Hosie et al., identified residues located within the transmembrane domain of GABA_AR α and β subunits that are critical for the direct activation ($\alpha 1$ –6; threonine 236, $\beta 1$ –3; tyrosine 284) and allosteric potentiation ($\alpha 1$ –6 asparagine 407, and $\alpha 1$ –6 glutamine 241) by neurosteroids [66–68]. Accordingly, mutation of glutamine 241 (Q241) within the $\alpha 1$ –6 subunits, prevents allosteric potentiation of GABA_AR composed of $\alpha\beta\gamma$ and $\alpha\beta\delta$ subunits by neurosteroids [66, 67]. In photo-affinity labeling experiments using a photo-labeling neurosteroid analog and homomeric $\beta 3$ receptors, the only residue labeled was phenylalanine 301 in TM3 of the $\beta 3$ subunit [69].

Neurosteroids are increasingly being thought of as therapeutic agents for numerous conditions arising from anxiety, traumatic brain injury, epilepsy, and neurodevelopment [70]. Ganaxolone, a synthetic analog of allopregnanolone, a metabolite of progesterone, is currently showing great promise in clinical trials for use in pediatric and adult cases of seizure disorders [71].

Conclusion

Mutations, CNVs, and polymorphisms of GABA_AR subunits have been linked to numerous neurodevelopment disorders and epilepsies. It remains unknown how these modifications to the genes of subunits change the properties of the mature protein and change pharmacological responses to the numerous compounds that target GABA_ARs. Altered GABA_AR function or expression is an obvious therapeutic target to counteract these changes but we need to better understand the changes that occur at a cellular, circuit, and organism level.

Acknowledgments Paul A. Davies has received financial support through NIH-NIMH Grant # MH097446.

Compliance with Ethics Guidelines

Conflict of Interest Paul A. Davies declares that he has no conflict of interest.

Human and Animal Rights and Informed Consent This article does not contain any studies with human or animal subjects performed by the author.

References

Papers of particular interest, published recently, have been highlighted as:

- Of importance

1. Barnard EA, Darlison MG, Fujita N, Glencorse TA, Levitan ES, Reale V, Schofield PR, Seeburg PH, Squire MD, Stephenson FA. Molecular biology of the GABAA receptor. *Adv Exp Med Biol*. 1988;236:31–45.
2. Joyce CJ. In silico comparative genomic analysis of GABAA receptor transcriptional regulation. *BMC Genom*. 2007;8:203.
3. Rudolph U, Mohler H. GABA-based therapeutic approaches: GABAA receptor subtype functions. *Curr Opin Pharmacol*. 2006;6:18–23.
4. Luscher B, Keller CA. Regulation of GABAA receptor trafficking, channel activity, and functional plasticity of inhibitory synapses. *Pharmacol Ther*. 2004;102:195–221.
5. Jacob TC, Moss SJ, Jurd R. GABA(A) receptor trafficking and its role in the dynamic modulation of neuronal inhibition. *Nat Rev Neurosci*. 2008;9:331–43.
6. Mortensen M, Smart TG. Extrasynaptic alphabeta subunit GABAA receptors on rat hippocampal pyramidal neurons. *J Physiol*. 2006;577:841–56.
7. Mortensen M, Patel B, Smart TG. GABA potency at GABA(A) receptors found in synaptic and extrasynaptic zones. *Front Cell Neurosci*. 2011;6:1.
8. Houston CM, McGee TP, Mackenzie G, Troyano-Cuturi K, Rodriguez PM, Kutsarova E, Diamanti E, Hosie AM, Franks NP, Brickley SG. Are extrasynaptic GABAA receptors important targets for sedative/hypnotic drugs? *J Neurosci*. 2012;32:3887–97.
9. Chandra D, Jia F, Liang J, Peng Z, Suryanarayanan A, Werner DF, Spigelman I, Houser CR, Olsen RW, Harrison NL, et al. GABAA receptor alpha 4 subunits mediate extrasynaptic inhibition in thalamus and dentate gyrus and the action of gaboxadol. *Proc Natl Acad Sci USA*. 2006;103:15230–5.
10. Glykys J, Mann EO, Mody I. Which GABAA receptor subunits are necessary for tonic inhibition in the hippocampus? *J Neurosci*. 2008;28:1421–6.
11. Glykys J, Mody I. Activation of GABAA receptors: views from outside the synaptic cleft. *Neuron*. 2007;56:763–70.
12. Brickley SG, Mody I. Extrasynaptic GABA(A) receptors: their function in the CNS and implications for disease. *Neuron*. 2012;73:23–34.
13. Lee S, Yoon BE, Berglund K, Oh SJ, Park H, Shin HS, Augustine GJ, Lee CJ. Channel-mediated tonic GABA release from glia. *Science*. 2010;330:790–6.
14. Olah S, Fule M, Komlosi G, Varga C, Baldi R, Barzo P, Tamas G. Regulation of cortical microcircuits by unitary GABA-mediated volume transmission. *Nature*. 2009;461:1278–81.
15. • Wlodarczyk AI, Sylantyev S, Herd MB, Kersante F, Lambert JJ, Rusakov DA, Linthorst AC, Semyanov A, Belelli D, Pavlov I, et al. GABA-independent GABAA receptor openings maintain tonic currents. *J Neurosci* 2013;33:3905–4. *The authors describe in dentate gyrus granule cells a tonic current that is insensitive to block by gabazine but is blocked by picrotoxin. Because the GABA competitive antagonist did not block tonic current the authors proposed that the current was coming from a GABA independent spontaneous current.*
16. Maksay G, Thompson SA, Wafford KA. The pharmacology of spontaneously open alpha 1 beta 3 epsilon GABA A receptor-ionophores. *Neuropharmacology*. 2003;44:994–1002.
17. Lee HH, Deeb TZ, Walker JA, Davies PA, Moss SJ. NMDA receptor activity downregulates KCC2 resulting in depolarizing GABAA receptor-mediated currents. *Nat Neurosci*. 2011;14:736–43.
18. Deeb TZ, Lee HH, Walker JA, Davies PA, Moss SJ. Hyperpolarizing GABAergic transmission depends on KCC2 function and membrane potential. *Channels (Austin)*. 2011;5:475–81.
19. Deeb TZ, Maguire J, Moss SJ. Possible alterations in GABAA receptor signaling that underlie benzodiazepine-resistant seizures. *Epilepsia*. 2012;53(Suppl 9):79–88.
20. Luscher B, Shen Q, Sahir N. The GABAergic deficit hypothesis of major depressive disorder. *Mol Psychiatry*. 2011;16:383–406.
21. Petryshen TL, Middleton FA, Tahl AR, Rockwell GN, Purcell S, Aldinger KA, Kirby A, Morley CP, McGann L, Gentile KL, et al. Genetic investigation of chromosome 5q GABAA receptor subunit genes in schizophrenia. *Mol Psychiatry*. 2005;10:1074–88.
22. Maguire J, Mody I. Neurosteroid synthesis-mediated regulation of GABA(A) receptors: relevance to the ovarian cycle and stress. *J Neurosci*. 2007;27:2155–62.
23. Maguire JL, Stell BM, Rafizadeh M, Mody I. Ovarian cycle-linked changes in GABA(A) receptors mediating tonic inhibition alter seizure susceptibility and anxiety. *Nat Neurosci*. 2005;8:797–804.
24. Maguire J, Mody I. GABA(A)R plasticity during pregnancy: relevance to postpartum depression. *Neuron*. 2008;59:207–13.
25. Curia G, Papouin T, Seguela P, Avoli M. Downregulation of tonic GABAergic inhibition in a mouse model of fragile X syndrome. *Cereb Cortex*. 2009;19:1515–20.
26. D'Hulst C, Kooy RF. The GABAA receptor: a novel target for treatment of fragile X? *Trends Neurosci*. 2007;30:425–31.
27. • Epi KC, Epilepsy Phenome/Genome P, Allen AS, Berkovic SF, Cossette P, Delanty N, Dlugos D, Eichler EE, Epstein MP, Glauser T, et al. De novo mutations in epileptic encephalopathies. *Nature*. 2013;501:217–21. *This study implicated a mutation in the gene encoding for the beta 3 subunit for epileptic encephalopathies. It is unknown what these mutations do to channel function but this study raises the possibility of personalized therapy and drug development directed to GABA_ARs containing the beta 3 subunit.*
28. Polan MB, Pastore MT, Steingass K, Hashimoto S, Thrush DL, Pyatt R, Reshmi S, Gastier-Foster JM, Astbury C, McBride KL. Neurodevelopmental disorders among individuals with duplication of 4p13 to 4p12 containing a GABA receptor subunit gene cluster. *Eur J Hum Genet*. 2014;22:105–9.
29. Hogart A, Wu D, LaSalle JM, Schanen NC. The comorbidity of autism with the genomic disorders of chromosome 15q11.2-q13. *Neurobiol Dis*. 2010;38:181–91.
30. Macdonald RL, Kang JQ, Gallagher MJ. GABAA receptor subunit mutations and genetic epilepsies. In: Noebels JL, Noebels JL, Avoli M, Rogawski MA, Olsen RW, Delgado-Escueta AV, editors. *Jasper's basic mechanisms of the epilepsies*. 4th ed. Oxford: Oxford University Press; 2012.
31. • Gurba KN, Hernandez CC, Hu N, Macdonald RL. GABRB3 mutation, G32R, associated with childhood absence epilepsy alters alpha1beta3gamma2L gamma-aminobutyric acid type A (GABAA) receptor expression and channel gating. *J Biol Chem*. 2012;287:12083–97. *The beta 3 subunit G32R mutation altered salt*

- bridges at subunit interfaces, and the authors suggested and increase in the likelihood of homomeric beta3 receptors. It is an interesting prospect that homomeric beta3 receptors can form in vivo.
32. Davies PA, Kirkness EF, Hales TG. Modulation by general anaesthetics of rat GABAA receptors comprised of alpha 1 beta 3 and beta 3 subunits expressed in human embryonic kidney 293 cells. *Br J Pharmacol*. 1997;120:899–909.
 33. Bassell GJ, Warren ST. Fragile X syndrome: loss of local mRNA regulation alters synaptic development and function. *Neuron*. 2008;60:201–14.
 34. Verkerk AJ, Pieretti M, Sutcliffe JS, Fu YH, Kuhl DP, Pizzuti A, Reiner O, Richards S, Victoria MF, Zhang FP, et al. Identification of a gene (FMR-1) containing a CGG repeat coincident with a breakpoint cluster region exhibiting length variation in fragile X syndrome. *Cell*. 1991;65:905–14.
 35. Fatemi SH, Folsom TD, Kneeland RE, Liesch SB. Metabotropic glutamate receptor 5 upregulation in children with autism is associated with underexpression of both Fragile X mental retardation protein and GABAA receptor beta 3 in adults with autism. *Anat Rec (Hoboken)*. 2011;294:1635–45.
 36. Miyashiro KY, Beckel-Mitchener A, Purk TP, Becker KG, Barret T, Liu L, Carbonetto S, Weiler IJ, Greenough WT, Eberwine J. RNA cargoes associating with FMRP reveal deficits in cellular functioning in Fmr1 null mice. *Neuron*. 2003;37:417–31.
 37. Fatemi SH, Folsom TD, Rooney RJ, Thuras PD. mRNA and protein expression for novel GABAA receptors theta and rho2 are altered in schizophrenia and mood disorders; relevance to FMRP-mGluR5 signaling pathway. *Transl Psychiatry*. 2013;3:e271.
 38. Fatemi SH, Kneeland RE, Liesch SB, Folsom TD. Fragile X mental retardation protein levels are decreased in major psychiatric disorders. *Schizophr Res*. 2010;124:246–7.
 39. Fatemi SH, Folsom TD, Rooney RJ, Thuras PD. Expression of GABAA alpha2-, beta1- and varepsilon-receptors are altered significantly in the lateral cerebellum of subjects with schizophrenia, major depression and bipolar disorder. *Transl Psychiatry*. 2013;3:e303.
 40. Pritchett DB, Luddens H, Seeburg PH. Type I and type II GABAA-benzodiazepine receptors produced in transfected cells. *Science*. 1989;245:1389–92.
 41. Macdonald RL, Weddell MG, Gross RA. Benzodiazepine, beta-carboline, and barbiturate actions on GABA responses. *Adv Biochem Psychopharmacol*. 1986;41:67–78.
 42. Hales TG, Lambert JJ. The actions of propofol on inhibitory amino acid receptors of bovine adrenomedullary chromaffin cells and rodent central neurones. *Br J Pharmacol*. 1991;104:619–28.
 43. Cottrell GA, Lambert JJ, Peters JA. Modulation of GABAA receptor activity by alphaxalone. *Br J Pharmacol*. 1987;90:491–500.
 44. Peters JA, Kirkness EF, Callachan H, Lambert JJ, Turner AJ. Modulation of the GABAA receptor by depressant barbiturates and pregnane steroids. *Br J Pharmacol*. 1988;94:1257–69.
 45. Crestani F, Low K, Keist R, Mandelli M, Mohler H, Rudolph U. Molecular targets for the myorelaxant action of diazepam. *Mol Pharmacol*. 2001;59:442–5.
 46. Rudolph U, Mohler H. Analysis of GABAA receptor function and dissection of the pharmacology of benzodiazepines and general anesthetics through mouse genetics. *Annu Rev Pharmacol Toxicol*. 2004;44:475–98.
 47. Fritschy JM, Mohler H. GABAA-receptor heterogeneity in the adult rat brain: differential regional and cellular distribution of seven major subunits. *J Comp Neurol*. 1995;359:154–94.
 48. Rudolph U, Crestani F, Mohler H. GABA(A) receptor subtypes: dissecting their pharmacological functions. *Trends Pharmacol Sci*. 2001;22:188–94.
 49. Crestani F, Keist R, Fritschy JM, Benke D, Vogt K, Prut L, Bluthmann H, Mohler H, Rudolph U. Trace fear conditioning involves hippocampal alpha5 GABA(A) receptors. *Proc Natl Acad Sci USA*. 2002;99:8980–5.
 50. Ferlisi M, Shorvon S. The outcome of therapies in refractory and super-refractory convulsive status epilepticus and recommendations for therapy. *Brain*. 2012;135:2314–28.
 51. • Deeb TZ, Nakamura Y, Frost GD, Davies PA, Moss SJ. Disrupted Cl(−) homeostasis contributes to reductions in the inhibitory efficacy of diazepam during hyperexcited states. *Eur J Neurosci* 2013;38:2453–67. *We describe how the efficacy of diazepam is reduced due to the accumulation of intracellular Cl[−] as a result of a decrease in KCC-2 function. However, propofol is still effective at inhibiting neuronal activity by shunting the membrane conductance. This provides a target for further investigation into therapies for refractory status epilepticus.*
 52. Garcia PS, Kolesky SE, Jenkins A. General anesthetic actions on GABA(A) receptors. *Curr Neuropharmacol*. 2010;8:2–9.
 53. Jurd R, Arras M, Lambert S, Drexler B, Siegwart R, Crestani F, Zaugg M, Vogt KE, Ledermann B, Antkowiak B, et al. General anesthetic actions in vivo strongly attenuated by a point mutation in the GABA(A) receptor beta3 subunit. *FASEB J*. 2003;17:250–2.
 54. Reynolds DS, Rosahl TW, Cirone J, O'Meara GF, Haythornthwaite A, Newman RJ, Myers J, Sur C, Howell O, Rutter AR, et al. Sedation and anesthesia mediated by distinct GABA(A) receptor isoforms. *J Neurosci*. 2003;23:8608–17.
 55. Li GD, Chiara DC, Sawyer GW, Husain SS, Olsen RW, Cohen JB. Identification of a GABAA receptor anesthetic binding site at subunit interfaces by photolabeling with an etomidate analog. *J Neurosci*. 2006;26:11599–605.
 56. Yip GM, Chen ZW, Edge CJ, Smith EH, Dickinson R, Hohenester E, Townsend RR, Fuchs K, Sieghart W, Evers AS, et al. A propofol binding site on mammalian GABAA receptors identified by photolabeling. *Nat Chem Biol*. 2013;9:715–20.
 57. Chiara DC, Jayakar SS, Zhou X, Zhang X, Savechenkov PY, Bruzik KS, Miller KW, Cohen JB. Specificity of intersubunit general anesthetic-binding sites in the transmembrane domain of the human alpha1beta3gamma2 gamma-aminobutyric acid type A (GABAA) receptor. *J Biol Chem*. 2013;288:19343–57.
 58. Belelli D, Lambert JJ. Neurosteroids: endogenous regulators of the GABA(A) receptor. *Nat Rev Neurosci*. 2005;6:565–75.
 59. Belelli D, Harrison NL, Maguire J, Macdonald RL, Walker MC, Cope DW. Extrasynaptic GABAA receptors: form, pharmacology, and function. *J Neurosci*. 2009;29:12757–63.
 60. Twyman RE, Macdonald RL. Neurosteroid regulation of GABAA receptor single-channel kinetic properties of mouse spinal cord neurons in culture. *J Physiol*. 1992;456:215–45.
 61. Lambert JJ, Belelli D, Peden DR, Vardy AW, Peters JA. Neurosteroid modulation of GABAA receptors. *Prog Neurobiol*. 2003;71:67–80.
 62. Callachan H, Cottrell GA, Hather NY, Lambert JJ, Nooney JM, Peters JA. Modulation of the GABAA receptor by progesterone metabolites. *Proc R Soc Lond B*. 1987;231:359–69.
 63. Shu HJ, Eisenman LN, Jinadasa D, Covey DF, Zorumski CF, Mennerick S. Slow actions of neuroactive steroids at GABAA receptors. *J Neurosci*. 2004;24:6667–75.
 64. Olsen RW, Li GD. GABA(A) receptors as molecular targets of general anesthetics: identification of binding sites provides clues to allosteric modulation. *Can J Anaesth*. 2012. doi:10.1007/s12630-010-9429-7.
 65. Olsen RW, Chang CS, Li G, Hancher HJ, Wallner M. Fishing for allosteric sites on GABA(A) receptors. *Biochem Pharmacol*. 2004;68:1675–84.
 66. Hosie AM, Clarke L, da Silva H, Smart TG. Conserved site for neurosteroid modulation of GABA A receptors. *Neuropharmacology*. 2009;56:149–54.

67. Hosie AM, Wilkins ME, da Silva HM, Smart TG. Endogenous neurosteroids regulate GABAA receptors through two discrete transmembrane sites. *Nature*. 2006;444:486–9.
68. Hosie AM, Wilkins ME, Smart TG. Neurosteroid binding sites on GABA(A) receptors. *Pharmacol Ther*. 2007;116:7–19.
69. Chen ZW, Manion B, Townsend RR, Reichert DE, Covey DF, Steinbach JH, Sieghart W, Fuchs K, Evers AS. Neurosteroid analog photolabeling of a site in the third transmembrane domain of the beta3 subunit of the GABA(A) receptor. *Mol Pharmacol*. 2012;82:408–19.
70. Carver CM, Reddy DS. Neurosteroid interactions with synaptic and extrasynaptic GABAA receptors: regulation of subunit plasticity, phasic and tonic inhibition, and neuronal network excitability. *Psychopharmacology*. 2013;230:151–88.
71. Bialer M, Johannessen SI, Levy RH, Perucca E, Tomson T, White HS. Progress report on new antiepileptic drugs: a summary of the Eleventh Eilat Conference (EILAT XI). *Epilepsy Res*. 2013;103:2–30.