Retraction Note

Jasbir Singh

Published online: 9 April 2014
© Springer International Publishing Switzerland 2014

Retraction to:

DOI 10.2165/00044011-200323060-00004,
DOI 10.2165/00044011-200424120-00006,
DOI 10.2165/00044011-200525080-00006,
DOI 10.2165/00044011-200626040-00004,
DOI 10.2165/00044011-200626110-00003

The following articles have been retracted by the journal after an investigation found the data to be fabricated:

Fujii Y, Tanaka H. Prevention of nausea and vomiting with ramosetron after total hip replacement. Clin Drug Investig. 2003;23:405–9, DOI 10.2165/00044011-200323060-00004

Fujii Y, Nakayama M. Reduction of Propofol-Induced Pain through Pretreatment with Lidocaine and/or Flurbiprofen. Clin Drug Investig. 2004;24:749–53, DOI 10.2165/00044011-200424120-00006

Fujii Y, Nakayama M. Efficacy of Lignocaine plus Ketamine at Different Doses in the Prevention of Pain Due to Propofol Injection. Clin Drug Investig. 2005;25:537–42, DOI 10.2165/00044011-200525080-00006

Fujii Y, Tanaka H. Efficacy of granisetron for the treatment of postoperative nausea and vomiting in women undergoing breast surgery: a randomised, double-blind, placebo-controlled trial. Clin Drug Investig. 2006;26:203–8, DOI 10.2165/00044011-200626040-00004

Fujii Y, Shiga Y. Age-related differences in metoclopramide requirement for pain on injection of propofol. Clin Drug Investig. 2006;26:639–44, DOI 10.2165/00044011-200626110-00003

Jasbir Singh
Editor

The online version of the original article can be found under doi:10.2165/00044011-200525080-00006, doi:10.2165/00044011-200424120-00006, doi:10.2165/00044011-200525080-00006, doi:10.2165/00044011-200626040-00004, doi:10.2165/00044011-200626110-00003.

J. Singh (✉)
Adis, 41 Centorian Drive, Private Bag 65901,
Mairangi Bay, North Shore, 0754 Auckland, New Zealand
e-mail: cdi@adis.com