RETRACTION NOTE

Retraction Note to: Combination Antiemetic Regimens for Prevention of Postoperative Nausea and Vomiting

Focus on High-Risk Patients

Yoshitaka Fujii¹

Published online: 26 February 2018
© Springer International Publishing AG, part of Springer Nature 2018

Retraction Note to:
Clin Drug Invest 2002; 22 (9): 561–574
<https://doi.org/10.2165/00044011-200222090-00001>

The Editor-in-Chief has retracted the original article because a number of studies [1–14] included in this review (cited as references 25–29, 31, 32, 47, 48, 50–54) have been retracted, specifically those included in Tables III, IV, Sects. 3 and 4.2. This has rendered the content of the review scientifically unreliable.

Author Yoshitaka Fujii has not responded to any correspondence from the Editor-in-Chief about this retraction.

References

1. Fujii Y, Tanaka H, Toyooka H. Prophylactic antiemetic therapy with granisetron-dexamethasone combination in women undergoing breast surgery. *Acta Anaesthesiol Scand*. 1998;42:1038–42.
2. Fujii Y, Toyooka H, Tanaka H. Prophylactic antiemetic therapy with a combination of granisetron and dexamethasone in patients undergoing middle ear surgery. *Br J Anaesth*. 1998;81:754–6.
3. Fujii Y, Saitoh Y, Tanaka H, et al. Granisetron/dexamethasone combination for preventing postoperative nausea and vomiting after laparoscopic cholecystectomy. *Eur J Anaesthesiol*. 2000;17:64–8.
4. Fujii Y, Tanaka H, Kobayashi N. Granisetron/dexamethasone combination for the prevention of postoperative nausea and vomiting after thyroidectomy. *Anaesth Intensive Care*. 2000;28:266–9.
5. Fujii Y, Tanaka H, Toyooka H. Retracted Article: Granisetron plus dexamethasone provide more improved prevention of postoperative emesis than granisetron alone in children. *Can J Anaesth*. 1996;43:1229–32.
6. Fujii Y, Tanaka H, Toyooka H. The effects of dexamethasone on antiemetics in female patients undergoing gynecologic surgery. *Anesth Analg*. 1997;85:913–7.
7. Fujii Y, Saitoh Y, Tanaka H, et al. Granisetron/dexamethasone combination for reducing nausea and vomiting during spinal anesthesia for cesarean section: Retracted. *Anaesth Analg*. 1999;88:1346–50.
8. Fujii Y, Toyooka H, Tanaka H. Prevention of postoperative nausea and vomiting with a combination of granisetron and droperidol. *Anesth Analg*. 1998;86:613–6.
9. Fujii Y, Saitoh Y, Tanaka H, et al. RETRACTED ARTICLE: Prophylactic antiemetic therapy with granisetron-droperidol combination in patients undergoing laparoscopic cholecystectomy. *Can J Anaesth*. 1998;45:541–4.
10. Fujii Y, Tanaka H, Toyooka H. Granisetron-droperidol combination for the prevention of postoperative nausea and vomiting in female patients undergoing breast surgery. *Br J Anaesth*. 1998;81:387–9.
11. Fujii Y, Saitoh Y, Tanaka H, et al. Retracted: Combination of granisetron and droperidol in the prevention of nausea and vomiting after middle ear surgery. *J Clin Anesth*. 1999;11:108–12.
12. Fujii Y, Saitoh Y, Tanaka H, et al. Prevention of post-operative nausea and vomiting with combined granisetron and droperidol in women undergoing thyroidectomy. *Eur J Anaesthesiol*. 1999;16:688–91.
13. Fujii Y, Saitoh Y, Tanaka H, et al. Retracted: Combination of granisetron and droperidol for the prevention of vomiting after paediatric strabismus surgery. *Paediatr Anaesth*. 1999;9:329–33.
14. Fujii Y, Toyooka H, Tanaka H. A granisetron-droperidol combination prevents postoperative vomiting in children: Retracted. *Anesth Analg*. 1998;87:761–5.

The original article can be found online at <https://doi.org/10.2165/00044011-200222090-00001>.

✉ Yoshitaka Fujii
yfujii@igaku.md.tsukuba.ac.jp

¹ Department of Anaesthesiology, University of Tsukuba
Institute of Clinical Medicine, Tsukuba City, Ibaraki, Japan