Foreword

J. Craig Nelson

Published online: 25 May 2013
© Springer International Publishing Switzerland 2013

Depression is a common disorder worldwide. In the USA, the lifetime prevalence of major depressive disorder is 16.6 % [1]. Most individuals will have a family member or close friend who has had depression if they have not experienced depression themselves. In addition, Kessler et al. [1] found that the incidence of new cases appears to be rising in younger cohorts. If this trend continues, the prevalence of depression will increase.

In Asia reported prevalence rates are lower. In this issue, Drs Sato and Yeh discuss this subject, noting that high rates of suicide in Korea and Japan suggest that depression is underreported in Asia. This could result from stigma associated with admitting to a mental disorder or simply underrecognition of the symptoms of the disorder. In particular, somatic symptoms of depression are common but may not be recognized as symptoms of depression.

Although a variety of treatments is available for depression, most patients do not recover with initial treatment. The largest study to examine this question, the Sequenced Treatment Alternatives to Relieve Depression (STAR*D) study found that only 36 % of patients achieved remission after 12 weeks of treatment with citalopram [2]. Two-thirds of the patients failed to recover, thus the question of what to do next is regularly confronted. Drs Patkar and Pae provide a very comprehensive review of guidelines for the treatment of major depression from around the world. One challenge for the guidelines is keeping up with emerging evidence. For example, the American Psychiatric Association guideline for the treatment of major depression, while published in 2010, is

already out of date with respect to the assessment of the level of evidence for some treatment strategies [3]. Nevertheless, guidelines have flourished partly because of the paucity of evidence regarding a variety of practical questions. As a consequence, clinical consensus continues to be relied on.

One of the strategies for patients who have failed to remit with initial treatment is to add another agent. Drs Chang, Sato and Han review the evidence reported over the past three decades for adjunctive treatment with a variety of agents other than atypical antipsychotic drugs. These agents include lithium, thyroid, modafinil, buspirone, pindolol, methylphenidate, omega 3, folic acid and anticonvulsant drugs. During the past decade a large body of evidence supporting the use of atypical antipsychotic agents has emerged [4]. Two of these agents—aripiprazole and quetiapine XR—have been approved by the US Food and Drug Administration for use as adjunctive agents in depression. The combination of fluoxetine and olanzapine is also approved for use in treatment-resistant depression. Drs Kato and Chang provide a detailed review of the evidence for the efficacy of these agents. Continuing this discussion, Drs Pae and Patkar review a variety of clinical issues that arise during the use of atypical agents ranging from the timing of their use to considerations of the number needed to harm for adverse effects.

Another issue that relates to the treatment of resistant depression is careful monitoring of care. Measurement-based care is increasingly advocated in depression and is particularly helpful for distinguishing between the patient who responds but has residual symptoms and the patient who remits. Drs Han, Yeh, Kato, Sato, Chang and Pae review a variety of scales available for use to monitor care. They provide a clinical case to illustrate measurement-based care.

J. C. Nelson (✉)
Department of Psychiatry, University of California,
San Francisco, CA, USA
e-mail: CraigN@lppi.ucsf.edu

When adjunctive treatments are administered, the clinician needs to consider both pharmacological issues related to the drug added as well as the potential for drug interactions with the initial agent. In addition, ethnic differences may alter the administration of these agents. Drs Han and Pae review pharmacokinetic issues that arise during adjunctive treatment and consider ethnic differences in drug metabolism that may require dose adjustments.

As a group this collection of papers provides an up-to-date review of the use of adjunctive treatments in major depressive disorder with special attention to issues arising in Asian populations. The review ranges from the consideration of randomized controlled evidence to a discussion of practical issues of interest to clinicians. As a consequence, it will be of interest to a broad international audience.

Disclosure This manuscript has been published in a journal supplement that was created with an unrestricted educational grant from Korea Otsuka International Asia and Arab (KOIAA). The author has received lecture honoraria from Eli Lilly (Global), Lundbeck, Otsuka (Asia), and Schering Plough/Merck (Asia). The author has also served

on advisory boards or provided consultation for Avanir, Biovail, Bristol Myers Squibb, Cenestra Health, Corcept, Covidien, Eli Lilly, Forest, Labopharm, Lundbeck, Medtronic, Merck, Mylan (Dey Pharma), Novartis, Orexigen, Otsuka, Pfizer, Sanofi-Aventis, Shire, Sierra Neuropharmaceuticals, Sunovion, and Theracos.

References

1. Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, et al. The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *JAMA*. 2003;289(23):3095–105.
2. Rush AJ, Trivedi MH, Wisniewski SR, Nierenberg AA, Stewart JW, Warden D, et al. Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps: a STAR*D report. *Am J Psychiatry*. 2006;163(11):1905–17.
3. American Psychiatric Association. Practice guideline for the treatment of patients with major depressive disorder (revision 3rd edition). http://www.psychiatryonline.com/pracGuide/pracGuideChapToc_7. Accessed 2010.
4. Nelson JC, Papakostas GI. Atypical antipsychotic augmentation in major depressive disorder: a meta-analysis of placebo-controlled randomized trials. *Am J Psychiatry*. 2009;166:980–91.