ERRATUM

Erratum to: Variation in caries treatment proposals among dentists in Norway: the best interest of the child

A. Rønneberg¹ · A. B. Skaare¹ · B. Hofmann^{2,3} · I. Espelid¹

Published online: 28 August 2017
© European Academy of Paediatric Dentistry 2017

Erratum to: Eur Arch Paediatr Dent DOI 10.1007/s40368-017-0298-4

Owing to an error in typesetting, the results section of the abstract was rendered incorrectly. The correct wording is as follows:

Results The response rate was 65% ($n = 391$) among the GDPs. A majority of the GDPs preferred a new appointment with behaviour management techniques (BMT) to a child presenting pulpitis and pain. Dentists educated outside the Nordic region would use restraint more often as a treatment alternative when the child was in pain than Nordic-educated dentists ($p < 0.05$). Dentists with less

than 10 years of experience preferred BMT and sedation more often when the child was in pain than their older colleagues, who, however, preferred a waiting approach and no immediate treatment if the child was not in pain ($p < 0.05$).

The original publication has now been corrected accordingly.

The online version of the original article can be found under doi:[10.1007/s40368-017-0298-4](https://doi.org/10.1007/s40368-017-0298-4).

✉ A. Rønneberg
anne.ronneberg@odont.uio.no

¹ Department of Paediatric Dentistry and Behavioural Science, Faculty of Dentistry, Institute of Clinical Dentistry, University of Oslo, Oslo, Norway

² Section for Health, Technology and Society, The Norwegian University of Science and Technology (NTNU), Gjøvik, Norway

³ Centre for Medical Ethics, University of Oslo, Oslo, Norway