Conjoined Twins Following Assisted Reproduction Technique

Anuja Thomas¹ · Bijoy K. Balakrishnan¹ · Meenu Batra¹ · Rekha Raichel Kuriakose¹ · Kannoly Karunakaran Gopinathan¹

Received: 2 August 2015 / Accepted: 8 December 2015 / Published online: 21 December 2015
© Society of Fetal Medicine 2015

Abstract The incidence of conjoined twins as reported in a worldwide epidemiological study is 1.47 per 100,000 births. It varies from 1 in 45,000–200,000 births. Conjoined twins following assisted reproduction are even rarer and we report one such case. A 36-year-old woman who conceived by intracytoplasmic sperm injection, was referred to the feto-maternal unit for nuchal translucency scan. A diagnosis of thoracopagus conjoined twins was made on ultrasound at 12 weeks 3 days of gestation. The couple was counseled in detail and they opted for termination. Pregnancy was terminated medically and the ultrasound diagnosis was confirmed. Assisted reproduction techniques involve a risk of conjoined twins and the present case report supports this. Early diagnosis of conjoined twins and determination of the type of fusion is possible. This aids in discussing the prognosis and counseling the couple regarding options of termination or continuation of pregnancy.

Keywords Conjoined twins · Intracytoplasmic sperm injection (ICSI) · Thoracopagus · Omphalocele

Introduction

Conjoined twinning is a rare complication of monozygotic twin pregnancy. Its high rate of morbidity and mortality makes early prenatal diagnosis significant. Prognosis

depends on the type and site of fusion, degree of sharing of the vital organs, and associated anomalies.

Report of Case

A 36-year-old primigravida who conceived by intracytoplasmic sperm injection (ICSI) after 11 years of marriage, was referred to the fetomaternal unit for nuchal translucency scan. A diagnosis of thoracopagus conjoined twin was made on ultrasound at 12 weeks 3 days of gestation.

Ultrasonography showed two heads in the same plane very close to each other with no membrane in between. The fetuses were positioned face to face, bodies fused from upper chest with a single heart, with parallel arrangement of the spines. The fetal bodies were not separable and lacked independent movement. They maintained similar position even after observing for a prolonged time (Fig. 1).

The couple was counseled in detail and they opted for termination. The pregnancy was terminated medically and the ultrasound diagnosis was confirmed. The fetuses were males and they were united at the thoraces. There was an omphalocele and a single umbilical cord (Fig. 2). The couple refused autopsy.

Discussion

The incidence of conjoined twins as reported in a worldwide epidemiological study is 1.47 per 100,000 births [1]. It varies from 1 in 45,000–200,000 births. Conjoined twins following assisted reproduction are even rarer. It has been documented that assisted reproductive technology (ART) might increase the risk of conjoined twins [2]. Multiple factors might play a role in this, like ovulation induction,

✉ Anuja Thomas
dranujathomaz@gmail.com

¹ Department of Feto Maternal Medicine, CIMAR Hospital, Off NH-17, Thykkavu Bus Stop Cheranalloor, Edappally, Kochi, Kerala 682034, India

Fig. 1 Ultrasound image showing fetuses facing each other, two heads in the same plane with bodies fused from upper chest

Fig. 2 Thoracopagus male conjoined twins with omphalocele

ICSI, assisted hatching, in vitro culture condition, etc. [3]. There is a female to male ratio of 3:1 [4].

There are two theories of origin:

Theory of fission: According to this theory, incomplete division of the monozygotic embryonic disc occurs after day 13 of conception.

Theory of fusion: The theory of fusion states that the previously separated embryonic discs undergo secondary fusion. The fusion theory is more accepted recently [5].

It has been reported that the rate of monozygotic twins, after ART, has risen about 2.25 times above the background rate [6]. The occurrence of monozygotic twinning after assisted reproduction has been linked by different authors to ovulation induction [7], gonadotropin treatment [8], maternal age either younger [9] or older [10], culture conditions [11], day 5 embryo transfer [12–15], and prolonged culture due to abnormal hatching [16] or apoptosis in the embryo [17]. Recently, Sobek et al. [18] have proposed that the high incidence of monozygotic twins following assisted reproduction techniques is conditioned by hereditary factors and that good ovarian function only facilitates the expression.

First trimester ultrasound enables accurate determination of chorionicity and amnionity which is of great value in assisted reproduction. Apart from being an effective method of screening for aneuploidies, the 11–14 weeks scan can also identify many major nonchromosomal abnormalities [19]. Though feasible, early first trimester diagnosis of conjoined twins is associated with false positive rates owing to limited fetal movements at this gestation. Also, it has been reported that first trimester 3D imaging does not add any advantage to 2D [20]. A classification of conjoined twins based on the site of fusion was put forward by Winkler et al. [21] (Table 1).

The incidence of major congenital anomalies in conjoined twins is 10–20 %, excluding the defects at the site of fusion [22]. The anomalies include congenital heart defects, spina bifida, cystic hygroma, limb defects, abdominal wall defects, and diaphragmatic hernia [23]. Conjoined twins, especially the ones with major congenital anomalies, have a high mortality [24, 25]. About 60 % die in utero and another 35 % are stillbirth or die within 24 h of birth [21]. The success of separation surgeries depends on the complexity of fusion, degree of sharing of the organs, severity of abnormalities, and the post-operative clinical conditions.

Conclusion

ART has increased the rate of conjoined twins as is evident from this rare case following ICSI. Early diagnosis of conjoined twinning and determination of the type of fusion

Table 1 Classification of conjoined twins modified from reference [21]

Type	Description	Incidence (%)
<i>Ventral union</i>		
Cephalopagus	Fused from head to umbilicus, lower abdomen and pelvis separated. Each twin with two extremities	11
Thoracopagus	Thoraces are fused with a shared heart or a single interatrial vessel	20–40
Omphalopagus	Similar to thoracopagus without shared heart or a single interatrial vessel	18–33
Ischiopagus	Joined at the pelvis. External genitalia and anus always shared	6–11
<i>Dorsal union</i>		
Craniopagus	Joined at the skull except the face and foramen magnum. Skull, brain and meninges shared	2
Pyopagus	Joined at the sacrococcygeal and perineal regions. Anus shared and rectums separate. Spinal cord may be shared	18–28
Rachipagus	Fused dorsally above the level of sacrum	Rare
<i>Lateral union</i>		
Parapagus	Twins have side-by-side connection with shared pelvis and variable cephalad sharing defined as follows:	28
Dithoracic	Separate thoraces and heads	
Dicephalic	Separate heads, fused thoraces	
Diprosopus	Single head with two faces on the same side	

is possible. This aids in discussing the prognosis and counseling the couple regarding options of termination or continuation.

Compliance with Ethical Standards

Conflict of interest None.

References

- Mutchinik OM. A world wide collaborative epidemiological study on conjoined twins. *Am J Med Genet C Semin Med Genet*. 2011;157C(4):274–87.
- Hirata T, Osuga Y, Fujimoto A, Oishi H, Hiori H, Fujiwara T, et al. Conjoined twins in a triplet pregnancy after after Intracytoplasmic sperm injection and blastocyst transfer: case report and review of literature. *Fertil Steril*. 2009;91(3):933.e9–12.
- Mercan R. Conjoined twin after intracytoplasmic sperm injection and transfer of day-3 embryos. *Fertil Steril*. 2011;96(2):e111–4.
- Abossolo T, Dancoisne P, Tuaillon J, Orvain E, Sommer JC, Rivière JP. Early prenatal diagnosis of asymmetric cephalothoracopagus twins. *J Gynecol Obstet Biol Reprod*. 1994;23(1):79–84.
- Spencer R. Theoretical and analytical embryology of conjoined twins: part I: embryogenesis. *Clin Anat*. 2000;13:36–53.
- Vitthala S, Gelbaya TA, Brison DR, Fitzgerald CT, Nardo LG. The risk of monozygotic twins after assisted reproductive technology: a systematic review and meta-analysis. *Hum Reprod Update*. 2009;15(1):45–55.
- Derom C, Vlietnick R, Derom R, Van de Bergh H, Thiery M. Increased monozygotic twinning rate after ovulation induction. *Lancet*. 1987;1:1236–8.
- Schachter M, Raziel A, Friedler S, Strassburger D, Bern O, Ron-El R. Monozygotic twinning after assisted reproductive techniques: a phenomenon independent of micromanipulation. *Hum Reprod*. 2001;16:1264–9.
- Schieve L, Meikle S, Peterson H, Jeng G, Bennett N, Wilcox L. Does assisted hatching pose a risk for monozygotic twinning in pregnancies conceived through in vitro fertilization? *Fertil Steril*. 2000;74:288–94.
- Abusheika N, Salha O, Sharma V, Brinsden P. Monozygotic twinning ad IVF/ICSI treatment: areport of 11 cases and review of literature. *Hum Reprod Update*. 2000;6:393–6.
- Cassuto G, Chavrier M, Menezo Y. Culture conditions and not prolonged culture time are responsible for monozygotic twinning in human invitro fertilization. *Fertil Steril*. 2003;80:462–3.
- da Costa A, Abdelmassih S, de Oliveira F, Abdelmassih V, Abdelmassih R, Nagy Z, Balmaceda J, et al. Monozygotic twinning and transfer at the blastocyst stage after ICSI. *Hum Reprod*. 2001;16:333–6.
- Sheiner E, Har-Vardi I, Potashnik G. The potential association between blastocyst transfer and monozygotic twinning. *Fertil Steril*. 2001;75:217–8.
- Jain J, Boonstanfar R, Slater CM, Francis M, Paulson R. Monozygotic twins and triplets in association with blkasticeyst transfer. *J Assist Reprod Genet*. 2004;21:103–7.
- Wright V, Schieve L, Vahratian A, Reynolds M. Monozygotic twinning associated with day 5 embryo transfer after IVF. *Hum Reprod*. 2004;19:1831–6.
- Malter HE, Cohen J. Blastocyst formation and hatching in vitro following zona drilling of mouse and human embryos. *Gamete Res*. 1989;24(1):67–80.
- Menezo YJ, Sakkas D. Monozygotic twinning: is it related to apoptosis in the embryo? *Hum Reprod*. 2002;17(1):247–8.
- Sobek A Jr, Zborilova B, Prochazka M, Silhanova E, Koutna O, Klaskova E, Tkadlec E, Sobek A. High incidence of monozygotic twinning after assisted reproduction is related to genetic information, but not to assisted reproduction technology itself. *Fertil Steril*. 2015;103:756–60.
- Souka AP, Pilalis A, Kavalakis Y, Kosmas Y, Antsaklis P, Antsaklis A. Assessment of fetal anatomy at the 11–14-week ultrasound examination. *Ultrasound Obstet Gynecol*. 2004;24:730–4.
- Pajkrt E, Jauniau E. First-trimester diagnosis of conjoined twins. *Prenat Diagn*. 2005;25:820–6.

21. Winkler N, Kennedy A, Byrne J, Woodward P. The imaging spectrum of conjoined twins. *Ultrasound Q*. 2008;24:249–55.
22. Jones KL. *Smith's recognizable patterns of human malformation*. 6th ed. Philadelphia: Elsevier; 2006.
23. Brizot ML, Liao AW, Lopes LM, Okumura M, Marques MS, Krebs V, et al. Conjoined twins pregnancies: experience with 36 cases from a single center. *Prenat Diagn*. 2011;31:1120–5.
24. Fernandes GL, Matsubara FK, Marques FK, et al. Three-dimensional prenatal diagnosis of monocephalus diprosopus tetraophthalmos. *J Ultrasound Med*. 2010;29(3):501–3.
25. Spitz L, Kiely EM. Conjoined twins. *J Am Med Assoc*. 2003;289(10):1307–10.